

## Changes of physicochemical indicators during mastitis and the effects of milk ejection on their sensitivity

Rupert M Bruckmaier<sup>1</sup>\*, Daniel Weiss<sup>1</sup>, Martin Wiedemann<sup>2</sup>, Susanne Schmitz<sup>1</sup> and Georg Wendl<sup>2</sup>

<sup>1</sup> Lehrstuhl für Physiologie, Technische Universität München, 85350 Freising, Germany

<sup>2</sup> Bayerische Landesanstalt für Landwirtschaft, Institut für Landtechnik, Bauwesen und Umwelttechnik, 85354 Freising, Germany

Received 29 August 2003 and accepted for publication 28 November 2003

We examined the relationship between physicochemical indicators and somatic cells in the milk of dairy cows during experimentally induced mastitis and their significance as indicators for use in controlling udder health. We were concerned particularly with the effect of alveolar milk ejection on the sensitivity of these indicators. In Expt 1, *Escherichia coli* lipopolysaccharide (*Esch. coli* LPS) was injected into the left rear quarter to induce an inflammatory reaction in one quarter in each of six cows. The contralateral control quarter was injected with a solution of NaCl (9 g/l). Nine milk samples were taken from both quarters until 60 h after injection. In Expt 2, repeated milk samples were taken every 20 s from one quarter during a 120-s teat stimulation in 20 cows with different somatic cell counts (SCC). Quarters were clustered for low (<5·0 log cells/ml), mid (5·0–5·7 log cells/ml) and high (>5·7 log cells/ml) SCC of the sample taken at t=0 s. Samples were analysed for SCC, electrical conductivity (EC) and Na<sup>+</sup> and Cl<sup>-</sup> concentrations. During the experimental inflammation SCC, EC, Na<sup>+</sup> and Cl<sup>-</sup> peaked at 12 h from LPS administration and values in treated quarters (T) at this time were elevated to 7900, 157, 501 and 169% of the values in untreated quarters, respectively. In Expt 2, SCC, EC, Na<sup>+</sup> and Cl<sup>-</sup> in high SCC quarters were 2520, 121, 283 and 141% of low SCC quarters at the start of stimulation (t=0 s), respectively. Highly significant ( $P<0\cdot001$ ) differences in EC, Na<sup>+</sup> and Cl<sup>-</sup> between high and low SCC quarters disappeared owing to the onset of alveolar milk ejection 100 s after the first contact with the teat. In conclusion, SCC in cows' milk provided the strongest amplitude in the case of an intramammary inflammation. EC, Na<sup>+</sup> or Cl<sup>-</sup> were useful tools only if the measurements were performed in cisternal milk before the start of alveolar milk ejection.

**Keywords:** Electrical conductivity, ions in milk, SCC, milk ejection.

Electrical conductivity (EC) in milk has been discussed as an indicator of mammary inflammation for decades (Malcolm et al. 1942; Linzell & Peaker, 1975; Fernando et al. 1982). Results of experimental devices to measure EC were promising (Linzell & Peaker, 1975), although EC is influenced by parameters such as milking interval and milk composition (Wittkowski et al. 1979; Fernando et al. 1981; Woolford et al. 1998; Ontsouka et al. 2003). In-line systems were established to measure EC throughout milking (Gebre-Egziabher et al. 1979; Onyango et al. 1988). However, despite the larger pool of available data the sensitivity in detecting mastitis was not satisfactory for practical use (Maatje et al. 1992; Nielen et al. 1995).

EC is determined by the ions dissolved in milk, mainly Na<sup>+</sup> and Cl<sup>-</sup> (Linzell & Peaker, 1975; Wittkowski et al. 1979; Fernando et al. 1982). The ion concentration in normal milk is much lower than in blood serum. In the case of a mammary infection, the tight junctions lose integrity and the epithelium becomes leaky owing to neutrophil diapedesis and tissue damage by mastitis pathogens (Stelwagen et al. 1997; Nguyen & Neville, 1998). Consequently blood and milk components pass through the leaky epithelium.

The major proportion of milk, which in dairy cows is stored in the alveolar tissue, is available only after the start of milk ejection (Knight et al. 1994; Pfeilsticker et al. 1996). However, as much as 20% of the milk is stored in the cisternal compartment of the udder and is immediately available for milk removal before milk ejection occurs (Bruckmaier & Blum, 1998). Changes in EC in different

\*For correspondence; e-mail: bruckmaier@wzw.tum.de

milk fractions within a specific milking event have been discussed (Linzell & Peaker, 1975; Wittkowski et al. 1979; Hamann & Gyodi, 2000; Ontsouka et al. 2003). However, the specific effect of alveolar milk ejection induced by teat stimulation without milk being removed, even by solely taking milk samples, has not been studied before.

Two experiments were conducted to study the sensitivity of EC,  $\text{Na}^+$  and  $\text{Cl}^-$  measurements in detecting mastitis as compared with SCC. The hypothesis tested was that the expected exponential increase of SCC in response to an experimentally induced mastitis is accompanied by a clearly detectable increase of  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations and EC in the milk, which allows the use of these measures for the detection of mastitis. Most importantly, the influence of the occurrence of alveolar milk ejection on the ability of foremilk samples to detect udder health status via SCC, EC, and concentrations of  $\text{Na}^+$  and  $\text{Cl}^-$  was investigated.

## Materials and Methods

### Animals and milking management

Cows were kept in loose housing and were milked in a milking parlour at 05.00 and 16.00. The diet was maize and grass silage, hay and concentrate given according to individual production levels. Cows were in their first to ninth lactation and were 50–206 d in milk. In Expt 1, Brown Swiss cows were used, whereas seven Red Holstein and thirteen Brown Swiss cows with a daily yield of  $29.4 \pm 1.7$  kg were used in Expt 2.

### Experimental design

**Expt 1.** Experiments were conducted in six clinically healthy cows. Only rear quarters with a log SCC  $<5.2$  and which cultured negative for mastitis pathogens in a sample taken 7 d before the experiment were accepted for the study. *Esch. coli* endotoxin (lipopolysaccharide, LPS; 100 µg) dissolved in 10 ml of NaCl (9 g/l) was injected intramammarily into the left rear quarter (T) after the morning milking. The right rear quarter (C) was injected with 10 ml of NaCl solution only. Cisternal milk samples of 50 ml were collected without any prior stimulation immediately before and at 3, 6, 9 h after LPS treatment and during succeeding milkings at 12, 24, 36, 48 and 60 h after LPS treatment.

**Expt 2.** Repeated milk sampling was performed during pre-stimulation in twenty cows. All teats were manually stimulated for 120 s. Milk samples (40 ml) were collected from the start until the end of stimulation at 20-s intervals from one quarter, while the other quarters were continuously stimulated. Thus, seven consecutive samples were taken. The udder was not touched before the start of

stimulation and sampling. The procedure was repeated for three consecutive milkings in each cow.

Milk samples were analysed for SCC in the laboratory of the Milchprüfung Bayern e.V., 80336 München, Germany, using a MilkoScan 4500 analyser (Foss Electric, 3400 Hillerød, Denmark). Aliquots of each sample were frozen at  $-20^\circ\text{C}$  immediately after sampling for the determination of EC and  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations. EC was measured at  $25^\circ\text{C}$  using the LDM electrode from WTW (LDM 130, Wissenschaftlich-Technische-Werkstaetten GmbH, 82362 Weilheim, Germany).  $\text{Na}^+$  and  $\text{Cl}^-$  were determined in raw milk using the ion selective electrodes, models 9811 and 9617BN respectively (Orion Research, Beverly MA, USA).


### Statistical analyses

Results are presented as means  $\pm$  SEM. Results were analysed using the repeated measures analysis of the MIXED procedure and tested for significance ( $P < 0.05$ ) using Least Significant-Difference Test (LSD) of the SAS (version 8.01) program package. For Expt 2, quarters were classified according to the SCC of the first sample obtained at  $t = 0$  s. Three clusters were used: SCC  $<5.0$  log cells/ml (SCC I),  $5.0\text{--}5.7$  log cells/ml (SCC II) and  $>5.7$  log cells/ml (SCC III). In Expt 1, the mixed model contained the treatment and the time of sampling as fixed effects and the animal was considered as a repeated factor. Likewise in Expt 2, the model included the time of sampling and the SCC cluster as fixed effects, and the animal entered the model as repeated factor.

## Results

### Expt 1

SCC before treatment was not significantly different between T and C quarters although a slight but insignificant rise of SCC was observed as compared with the sample taken 7 d before the experiment in some of the control quarters. Values of EC,  $\text{Na}^+$  and  $\text{Cl}^-$  were low and did not differ between the quarters (Fig. 1). Within 3 h after LPS application, SCC, EC,  $\text{Na}^+$  and  $\text{Cl}^-$  increased in T quarters. In C quarters, EC,  $\text{Na}^+$  and  $\text{Cl}^-$  increased transiently owing to the administration of NaCl (9 g/l) solution and were numerically but not significantly higher at 3 h than at 0 h. Subsequently, SCC and  $\text{Na}^+$  were significantly higher in T quarters than in C quarters between 6 and 24 h after administration of LPS and NaCl, respectively. EC and  $\text{Cl}^-$ , however, were significantly higher in T than in C quarters only at 9 and 12 h after LPS and NaCl treatments, respectively. SCC, EC,  $\text{Na}^+$  and  $\text{Cl}^-$  peaked at 12 h after LPS administration. Their peak values in T quarters, expressed as linear values, were 7900, 157, 501 and 169% of C quarters, respectively. SCC remained elevated in T quarters until the end of the experiment (60 h), whereas all



**Fig. 1.** SCC, EC,  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations in the milk of one treated ( $\blacktriangle$ ) and one control quarter ( $\circ$ ) before and after application of 100  $\mu\text{g}$  lipopolysaccharide (LPS) or NaCl control solutions,  $n=6$  cows;  $a, b, c, d$  means without a common letter throughout the time span are different in treated quarters at corresponding times ( $P<0.05$ ); \* indicates a significant ( $P<0.05$ ) difference between treated and control quarters.

other parameters returned to pre-treatment levels within 24 h after LPS treatment.

### Expt 2

Figure 2 illustrates the results obtained in Expt 2. In the first sample ( $t=0$  s) mean SCC was  $4.62 \pm 0.07$ ,  $5.34 \pm 0.04$  and  $6.00 \pm 0.08$  log cells/ml in SCC I ( $n=23$  samples), SCC II ( $n=14$  samples) and SCC III ( $n=22$  samples), respectively. In the first sample ( $t=0$  s) mean SCC in SCC III quarters, expressed as a linear value, was 2520% of the results in SCC I quarters. SCC decreased within 120 s and was significantly lower at 100 s and 120 s than at the start of sampling ( $t=0$  s) in SCC III and SCC I quarters. However, the SCC group differences between SCC I, SCC II and SCC III quarters remained significant throughout the 120-s sampling period.

From 0 to 60 s, EC in SCC III quarters was significantly higher than in SCC I and SCC II quarters, whereas EC did not differ significantly between SCC I and SCC II quarters. EC in SCC III quarters was 121% of that in SCC I quarters at  $t=0$  s. A dramatic decrease of EC occurred between 40 and 100 s of sampling. In SCC III quarters, EC was significantly lower at  $t=80$  s than at  $t=0$  s. The difference of EC between SCC III and SCC II quarters was no longer significant from  $t=100$  s, i.e., EC was similar in SCC III, SCC II and SCC I quarters 100 s after the start of sampling.

$\text{Na}^+$  was significantly higher in SCC III quarters than in SCC I throughout the 120-s sampling period, whereas for  $\text{Cl}^-$ , the significant difference between SCC I and SCC III quarters disappeared at  $t=100$  s. At  $t=0$  s,  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations in SCC III quarters were 283 and 141% of SCC I quarters, respectively.  $\text{Na}^+$  and  $\text{Cl}^-$  did not differ significantly between SCC I and SCC II quarters throughout the sampling period. Despite the dramatic decrease of  $\text{Na}^+$  and  $\text{Cl}^-$  that occurred during the course of stimulation,  $\text{Na}^+$  was still significantly higher in SCC III than in SCC I quarters at  $t=120$  s.

### Discussion

Intramammary injection of 100  $\mu\text{g}$  LPS resulted in a considerable inflammatory reaction in the respective quarter, as reported earlier (Bruckmaier et al. 1993; Moussaoui et al. 2002; Schmitz et al. 2002; Yagi et al. 2002). The observed peak of SCC at 12 h after administration of LPS demonstrates that the inflammatory and immunological reaction required a latency period of several hours. Most of the injected LPS would be expected to be removed by the milking at 12 h after application of LPS, since the transient reaction of EC,  $\text{Na}^+$  and  $\text{Cl}^-$  returned to basal levels. Surprisingly, SCC remained elevated until 60 h after LPS application, although tight junction integrity seemed to be largely restored as indicated by  $\text{Na}^+$  and  $\text{Cl}^-$  patterns. It may be speculated that only during periods of high somatic cell diapedesis rates do the tight junctions get


leaky enough (Stelwagen et al. 1997; Nguyen & Neville, 1998) to allow considerable movement of ions from blood into milk. This may explain the limited sensitivity of  $\text{Na}^+$ ,  $\text{Cl}^-$  and EC as indicators of mastitis in the subclinical range (as shown in Expt 2).

The constant SCC in C quarters documents that the elevated SCC in T quarters was a local reaction in response to LPS administration. As expected, EC,  $\text{Na}^+$  and  $\text{Cl}^-$  values in C quarters were numerically elevated at 3 h after application of NaCl solution. For the same reason, the results for T quarters seem to be shifted. However, the dramatic increase of SCC from 0 to 3 h documents the considerable amplitude of the SCC rise as compared with the changes in EC,  $\text{Na}^+$  and  $\text{Cl}^-$  during the impairment of tight junction integrity.

In Expt 2, single quarters with different SCC levels, but without clinical mastitis were used for repeated milk sampling during a teat stimulation of 120 s. For SCC, EC,  $\text{Na}^+$  and  $\text{Cl}^-$ , the values decreased during the sampling period but none of these variables changed before the sample taken at 60 s. It is therefore obvious that the continuous teat stimulation during sampling induced alveolar milk ejection. The onset of milk ejection, i.e., the occurrence of alveolar milk in the mammary cistern, caused a mixture of alveolar and cisternal milk and hence an immediate change of milk ion concentrations (Ontsouka et al. 2003). The lag time between start of stimulation and the start of milk ejection was previously found to be 50–100 s, as a function of the degree of udder filling (Bruckmaier & Hilger, 2001). As a consequence of milk ejection the measurements of EC and of  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations lose sensitivity as mastitis indicators. This fact needs to be considered in the case of foremilk sampling for EC,  $\text{Na}^+$  and  $\text{Cl}^-$  measurements. Foremilk can only be expected to be cisternal milk until about 40 s from the first touch of the udder.

With respect to the differences between SCC III and SCC I quarters, SCC was the most sensitive indicator. The dramatic increase during inflammation was the result of a local immune reaction and mediator-linked active transport of cells into the milk of the respective quarter. In contrast, the rise of  $\text{Na}^+$  and  $\text{Cl}^-$ , and consequently of EC, was a result of diffusion through leaky tight junctions during somatic cell influx. Thus the increment of EC,  $\text{Na}^+$  and  $\text{Cl}^-$  during intramammary inflammation was only a linear shift as compared with the exponential increase of SCC. Therefore, the mixture of cisternal milk with alveolar milk resulted in the disappearance of highly significant differences between SCC I and SCC III quarters for all physicochemical indicators investigated.

It has to be pointed out that the sensitivity of EC,  $\text{Na}^+$  and  $\text{Cl}^-$  was not great enough to distinguish between SCC I and SCC II quarters. In automatic milking systems, the sensitivity of the currently used measurement of EC could be slightly improved by inter-quarter ratio and comparison from one milking to the next even though the EC is measured after the milk ejection.



**Fig. 2.** SCC, EC,  $\text{Na}^+$  and  $\text{Cl}^-$  concentrations in the different milk fractions, 0 to 120 s after the first contact with the udder; data are clustered according to the SCC of the first sample ( $t=0$  s), ▼ <5·0  $\log$  cells/ml (SCC I), ● 5·0–5·7  $\log$  cells/ml (SCC II) and ▲ >5·7  $\log$  cells/ml (SCC III);  $n=20$  cows; three milkings each cow; <sup>a,b,c</sup> means with different letters indicate differences between clusters ( $P<0·05$ ); \* indicates a significant ( $P<0·05$ ) difference compared with the sample  $t=0$  s.

In accordance with the results of Expt 1, only a high somatic cell diapedesis rate during acute inflammation seems to allow electrolyte diffusion from blood into milk, while during the influx of cells at a lower rate the electrolyte concentrations return to their basal levels. Recommended SCC thresholds to distinguish between disturbed and undisturbed milk composition range between 5·0 log cells/ml (Hillerton, 1999) and 5·3 log cells/ml (Smith, 1995). Therefore the studied variables were not satisfactory for distinguishing these thresholds by using the current in-line sampling routine. However, physicochemical indicators could detect serious immune reactions during mastitis.

In conclusion, SCC is the most sensitive measurement to detect mastitis in bovine milk, both before and after the occurrence of milk ejection. In contrast, EC, Na<sup>+</sup> or Cl<sup>-</sup> concentrations can be useful tools for in-line measuring but only if quarter milk samples of the cisternal fraction are available. To achieve this goal, foremilk samples must be taken before the occurrence of alveolar milk ejection.

We thank the farms Veitshof and Hirschau, Technical University Munich, for providing the animals and the Milchprüfung Bayern e.V. for analyses of the milk samples. The help of Mr A Knon, Mrs and Mr Ochsendorf during the animal experiments is gratefully acknowledged. The study was partially supported by Bavarian Ministry of Agriculture and Forestry and by a grant from the Hanns-Seidel-Foundation for M Wiedemann.

## References

- Bruckmaier RM & Blum JW** 1998 Oxytocin release and milk removal in ruminants. *Journal of Dairy Science* **81** 939–949
- Bruckmaier RM & Hilger M** 2001 Milk ejection in dairy cows at different degrees of udder filling. *Journal of Dairy Research* **68** 369–376
- Bruckmaier RM, Schaellibaum M & Blum JW** 1993 *Escherichia coli* endotoxin-induced mastitis in dairy cows – changes and importance of insulin like growth factor-1 and oxytocin. *Milchwissenschaft* **48** 374–378
- Fernando RS, Rindsig RB & Spahr SL** 1981 Effect of length of milking interval and fat content on milk conductivity and its use for detecting mastitis. *Journal of Dairy Science* **64** 678–682
- Fernando RS, Rindsig RB & Spahr SL** 1982 Electrical conductivity of milk for detection of mastitis. *Journal of Dairy Science* **65** 656–664
- Gebre-Egziabher AH, Wodd C, Robar JD & Blankenagel G** 1979 Evaluation of automatic mastitis detection equipment. *Journal of Dairy Science* **62** 1108–1114
- Hamann J & Gyodi P** 2000 Somatic cells and electrical conductivity in relation to milking frequency. *Milchwissenschaft* **55** 303–306
- Hillerton JE** 1999 Redefining mastitis based on somatic cell count. *IDF Bulletin* **345** 4–9
- Knight CH, Hirst D & Dewhurst RJ** 1994 Milk accumulation and distribution in the bovine udder during the interval between milkings. *Journal of Dairy Research* **61** 167–177
- Linzell JL & Peaker M** 1975 Efficiency of the measurement of the electrical conductivity of milk for the detection of subclinical mastitis in cows: Detection of infected cows at a single visit. *British Veterinary Journal* **131** 447–461
- Maatje K, Huijsmans PJM, Rossing W & Hogewerf PH** 1992 The efficacy of in-line measurement of quarter milk electrical conductivity, milk yield and milk temperature for the detection of clinical and subclinical mastitis. *Livestock Production Science* **30** 239–249
- Malcolm JF, King CW & Campbell MM** 1942 The value of the cell content and electrical conductivity of milk as criteria of bovine mastitis. In *Proceedings of the Society of Agricultural Bacteriologists*, pp. 30–34
- Moussaoui F, Michelutti I, Le Roux Y & Laurent F** 2002 Mechanisms involved in milk endogenous proteolysis induced by a lipopolysaccharide experimental mastitis. *Journal of Dairy Science* **85** 2562–2570
- Nguyen DD & Neville MC** 1998 Tight junction regulation in the mammary gland. *Journal of Mammary Gland Biology and Neoplasia* **3** 233–273
- Nielen M, Schukken YH, Brand A, Deluyker HA & Maatje K** 1995 Detection of subclinical mastitis from on-line milking parlor data. *Journal of Dairy Science* **78** 1039–1049
- Ontsouka CE, Bruckmaier RM & Blum JW** 2003 Fractionized milk composition during removal of colostrum and mature milk. *Journal of Dairy Science* **86** 2005–2011
- Onyango CMJ, Marchant JA, Lake JR & Stanbridge A** 1988 A low maintenance conductivity sensor for detecting mastitis. *Journal of Agricultural Engineering Research* **40** 215–224
- Pfeilsticker HU, Bruckmaier RM & Blum JW** 1996 Cisternal milk in the dairy cow during lactation and after preceding teat stimulation. *Journal of Dairy Research* **63** 509–515
- Schmitz S, Pfaffl MW, Meyer HHD & Bruckmaier RM** 2002 mRNA expression of immunologically important factors and milk proteins in mammary tissue of dairy cows during LPS-induced mastitis. *Journal of Dairy Science* **85** (Suppl. 1) 8
- Smith KL** 1995 Standards for somatic cells in milk: physiological and regulatory. In *IDF Mastitis Newsletter* **144/21** 7–9
- Stelwagen K, Farr VC, McFadden HA, Prosser CG & Davis SR** 1997 Time course of milk accumulation-induced opening of mammary tight junctions, and blood clearance of milk components. *American Journal of Physiology* **273** R379–R386
- Wittkowski G, Gedek W & Kleinschroth E** 1979 [Significance of several parameters in different milking portions with regard to the diagnosis of bovine mastitis.] *Archiv für Lebensmittelhygiene* **30** 19–22
- Woolford MW, Williamson JH & Henderson HV** 1998 Changes in electrical conductivity and somatic cell count between milk fractions from quarters subclinically infected with particular mastitis pathogens. *Journal of Dairy Research* **65** 187–198
- Yagi Y, Shiono H, Shibahara T, Chikayama Y, Nakamura I & Ohnuma A** 2002 Increase in apoptotic polymorphonuclear neutrophils in peripheral blood after intramammary infusion of *Escherichia coli* lipopolysaccharide. *Veterinary Immunology and Immunopathology* **89** 115–125