

Hirt, C. (2016), Gravity forward modelling. In: Encyclopaedia of Geodesy (Ed. E. Grafarend), doi: 10.1007/978-3-319-02370-0_106-1.

Gravity forward modelling

Christian Hirt ^{1,2}

¹ Department of Spatial Sciences, Curtin University, Perth, WA, Australia

² Institute for Advanced Study, TU Munich, Germany

Definition

Gravity forward modelling. Computation of the gravity field of some given mass distribution

Introduction

Gravity forward modelling (GFM) denotes the computation of the gravitational field generated by some source mass distribution. The foundation of GFM is Newton's law of universal gravitation (1687) which states that the attraction force F between two bodies is proportional to the product of their masses m , M and inversely proportional to the square of their distance r :

$$F = G \frac{mM}{r^2} \quad (1)$$

where $G = 6.67384 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ is the universal gravitational constant (Mohr et al., 2012). In most practical applications of GFM, unit mass is assumed in the computation point, and the second body's mass M is considered the source of the gravitational field (Blakeley 1996). The visible topography, given in form of digital terrain models (DTM), is the most frequently used mass distribution in GFM, other distributions may include, but are not limited to, water masses, ice masses (Grombein et al., 2014), and crustal structures in the Earth's interior (Tenzer et al., 2012).

GFM yields the gravitational field of the mass distribution in terms of any functional of the field, e.g., gravitational potential, gravity disturbance (i.e., radial derivative of the potential), and vertical deflections (cf. Nagy et al., 2000). In the literature, a number of different expressions are in use to denote gravity effects or functionals computed from topography models, e.g., topographic gravity, topography-induced gravity, synthetic gravity, terrain effects and topographic reductions.

GFM is a central task in physical geodesy, e.g., in the context of geoid determination (Tziavos and Sideris, 2013) and gravity prediction (Forsberg and Tscherning, 1981). In potential field geophysics, GFM is relevant for the investigation of the interior structure of the Earth and the planets (Wieczorek, 2007). While GFM as forward method delivers the gravity field from mass models, inversion techniques seek to estimate the mass distribution from gravity observations, which however is inherently non-unique (Oldenburg, 1974). Gravity inversion involves iterative application of GFM based on different mass model assumptions until the forward modelled signal sufficiently fits the gravity observations (e.g., Ebbing et al., 2001).

This article briefly outlines representations of source mass models and describes GFM calculation techniques, grouped according to the domain of evaluation (spatial vs. spectral domain), then summarizes applications for GFM, selected results and directions.




Figure 1. Principle of Gravity forward modelling. A. Digital terrain model (DTM) and digital bathymetry model (DBM) as source of mass information. B. Source mass model based on DTM and DBM geometry information, and assignment of mass-density values. The masses are approximated by rectangular prisms in this example. C. Gravity effect (gravity disturbances) as result of the forward modelling of the source masses using Newtonian integration. Examples show Samothrace Island (40.45° latitude, 25.6° longitude).

Source mass models

As prerequisite for the numerical evaluation of Newton's integral, a model of the source mass distribution must be defined, representing the attracting body. This generally includes (a) the definition of the body's geometry, and (b) assignment of a suitable mass-density value. Often the attracting body is defined as a volumetric layer with some upper and lower bound (Tenzer et al., 2012; Balmino et al., 2012), e.g., the visible topography with the geoid as lower bound and terrain elevations (from a DTM) as upper bound. When ocean water masses are modelled (Fig. 1a,b), bathymetric depths from a digital bathymetry model (DBM) define the seafloor geometry as lower bound and the geoid as upper bound (Tenzer et al., 2011; Tocho et al., 2012). Depending on the purpose of the study, the geometry of various other bodies, e.g., ice-shields (Grombein et al., 2014), sediments (Ebbing et al. 2001), crustal or mantle structures (Tenzer et al., 2012; Tsoulis, 2013), can be defined from digital data sets.

Mass-density values assigned to the body can be based on geological samples (e.g., granite 2.7 g/cm⁻³, basalt 2.9 g/cm⁻³, salt rock 2.4 g/cm⁻³) or – in the absence of individual samples – on assumptions (e.g., 2.67 g cm⁻³ as an average value for rock), cf. Jacoby and Smilde (2009). While in many cases the mass-density is treated as constant value inside the attracting body, lateral (Eshagh, 2009; Götzl and Rummel, 2009) or vertical variations in mass-density (e.g., depth-dependent increase in sea water density, Tenzer et al., 2011) can be taken into account.

Following the superposition principle, gravity effects of a collection of masses (e.g., the visible topography, ocean water, and ice-shields) can be obtained through addition of gravity effects by the individual masses (e.g., Blakely, 1996). For computational reasons, water and ice masses are sometimes numerically compressed into layers of rock (rock-equivalent topography RET, Rummel et al., 1988), allowing to work with a single constant mass-density value.

The source mass model can be extended to accommodate the effect of isostatic compensation (e.g., Rummel et al., 1988; Makhloof and Ilk, 2008; Bagherbandi and Sjöberg, 2012; Grombein et al., 2014), e.g., based on the Airy-Heiskanen hypothesis through modification of the root (lower bound) of the topography, or the Pratt-Hayford hypothesis by using laterally varying mass-densities (Götzl and Rummel, 2009). If the source mass-defining DTM is high-pass filtered, e.g., through subtraction of some mean terrain surface, a residual terrain model (RTM, Forsberg and Tscherning 1981; Forsberg, 1984) is obtained. The RTM can be thought of as a model with oscillating positive and negative (in the sense of deficit) masses. In physical geodesy, the RTM is often used as for gravity field smoothing and prediction (Forsberg, 1984).

Gravity forward modelling in the spatial domain

Spatial domain GFM encompasses all evaluation methods for Newton's integral that yields the gravitational potential V generated by the source mass model M (e.g., Blakeley, 1996, 46)

$$V = G \iiint_M \frac{dm}{r} \quad (2)$$

where dm are elementary mass bodies of M and r is the separation between dm and a computation point. Newton's integral is practically evaluated by decomposing the source mass model M into elementary mass bodies dm , the gravitational potentials of which can be computed through analytical or numerical integration (Kuhn and Seitz, 2005). The gravitational potential of M is obtained through addition of all individual potential values implied by the elementary mass bodies (Newtonian integration). Frequently used elementary bodies are point masses, prisms with flat (Nagy et al., 2000) or inclined tops (Smith, 2000), tesseroids (Grombein et al., 2013), and polyhedrons (d'Urso, 2014), also see Papp (1996), Heck and Seitz (2007), Wild-Pfeiffer (2008) and Tsoulis et al. (2009). The gravitational potential of elementary bodies is usually obtained from closed-form expressions (e.g., Nagy et al., 2000), however, numerical integration methods can be an alternative (Asgharzadeh et al., 2007). A decomposition of the mass model into prisms is shown in Fig. 1b.

Gravity disturbances are routinely computed from DTM data through Newtonian integration (Fig. 1c), and subtracted from observed gravity measurements (topographic reduction), e.g., Forsberg (1984), Tziavos et al. (2010). The topographic reduction is often split into two parts, (i) the Bouguer reduction as the gravity effect induced by a solid mass slab or shell of constant thickness (e.g., Strang van Hees, 2000), and (ii) the terrain correction as the gravity effect of the actual DTM masses residual to the slab or shell (e.g., Forsberg, 1984). Terrain corrections are obtained through numerical evaluation of the terrain correction integral which can be derived from Eq. (2), cf. Nahavandchi and Sjöberg (1998).

Accurate spatial domain GFM requires Newtonian integration over the domain of all source masses, which often extend over the whole of the Earth's surface (e.g., Kuhn et al. 2009). In practice, however, integration over the source masses is sometimes done within some limited radius around the computation point (neglecting remote masses), and in planar approximation (instead of the more rigorous spherical or ellipsoidal approximation). The use of RTM as input source model in GFM keeps truncation and approximation effects small, while allowing limitation of the integration to some local radius (e.g., few 10s of km).

The computational effort for spatial domain GFM increase linearly with the number of computation points at which gravity effects are sought. (Sequential) gravity calculations in terms of densely spaced grids (e.g., 100 m resolution) and continental coverage is computationally expensive because Newton's integral has to be evaluated point by point. Parallel computation of gravity effects, e.g., on supercomputing platforms, significantly reduces computation times (e.g., Hirt and Kuhn, 2014). The limited computation radius in the RTM technique reduces the computational efforts too, however, only the high-frequency gravity signals of a mass distribution are obtained (Forsberg, 1984).

Gravity forward modelling in the spectral domain

These methods evaluate Newton's law of gravitation through integral transformation in the spectral domain (e.g., Rummel et al., 1988; Tenzer, 2005; Wieczorek, 2007). This generally require source mass models of global extent. Following Hirt and Kuhn (2014), (1) the source mass model (mostly a DTM) is raised to integer powers (1, 2, ... , n) and expanded into spherical harmonic series. This requires spherical harmonic analyses of each integer power of the source model. (2) The resulting sets of coefficients are used to obtain the gravitational potential - as a series expansion of the integer powers of the source mass distribution - in spherical harmonics. (3) The gravity field (e.g., in terms of potential, gravity disturbances) in the spatial domain is computed via spherical harmonic synthesis.

Spectral domain GFM is the preferred technique for planetary studies, e.g., calculation of Bouguer gravity fields for the Moon (Zuber et al., 2013) and Mars (Neumann et al., 2004). This is because observed gravity data for the planets is mostly given in spherical harmonics too. On Earth, the technique is increasingly used for applications with global scope. Examples include the generation of global Bouguer gravity maps in spherical harmonics (Balmino et al., 2012), study of the topographic potential (Novák, 2010; Gruber et al., 2013), all with ultra-high resolution in the km-range. While GFM in the spectral domain usually deploys a sphere as underlying reference body, recent refinements use an ellipsoid instead

(Claessens and Hirt, 2013; Wang and Yang 2013), which take into consideration the ellipsoidal shape, e.g., of Earth and Mars.

As a benefit of spectral domain GFM, the gravitational potential induced by the mass-density distribution is directly obtained in spherical harmonics, which allows straightforward computation of various functionals through synthesis, and spectral filtering. In contrast, space domain modelling requires one computation run per functional, which can be tedious.

As a second benefit of the spectral technique over space domain GFM, there is no numerical integration involved, which can be computationally expensive for computation point grids with global coverage. However, depending on the spatial resolution of the source mass model, multiple spherical harmonic analyses are required for convergence in the spectral method (e.g., Wieczorek, 2007; Claessens and Hirt, 2013), which increase the computational costs (e.g., 7 integer powers for a 5 arc-min resolution terrestrial mass model) as the resolution increases. Spectral and spatial domain GFM were shown to be numerically equivalent techniques (agreement at the 10^{-5} level) at least when the source mass model is limited to features at ~ 100 km scales or larger (Hirt and Kuhn, 2014). However, it is currently unclear if the techniques are equivalent for higher-resolution mass models too.

Applications, some results and directions

There are several GFM applications in geodesy and geophysics which rely on gravity functionals implied by mass models, particularly DTMs. Gravity observations reduced by the gravity effect of the topography reveal subsurface mass-anomalies (e.g., Ebbing et al., 2001; Jacoby and Smilde, 2009), and are pivotal in lithosphere studies (Wieczorek, 2007; Zuber et al., 2013). Forward-modelled gravity is effective at smoothing gravity observations before interpolation, particularly in mountainous areas (Forsberg and Tscherning, 1981). The strong correlation between gravity and topography in rugged terrain (Forsberg, 1984) can be used to approximate the short-wavelength gravity field with GFM. This is often exploited to predict a detailed gravity field where gravity measurements are limited (Pavlis et al., 2007), or to enhance spherical harmonic global gravity models at short spatial scales (Hirt, 2010). GFM is suitable to evaluate the quality of observed gravity fields (e.g., from satellite observations, Hirt et al., 2012), to generate synthetic models for geoid algorithm testing (Baran et al., 2006), and is central for topographic reductions in geoid determination (Jekeli and Serpas, 2003).

The availability of remotely-sensed DTMs (particularly from the SRTM mission) and improved supercomputing resources have stimulated the development of ultra-high resolution gravity field models based on GFM. Balmino et al. (2012) applied spectral domain GFM and global mass models to generate 2km resolution global grids of gravity effects for the construction of the World Gravity Map (WGM). Hirt et al. (2013) used the RTM technique along with observed gravity to construct a near-global gravity map with ultra-fine 220 m resolution (GGMplus).

It is foreseeable that future GFM will ultimately yield global gravity maps, and grids of terrain corrections at 100m resolution and finer, commensurate with the mass information available through remotely sensed DTMs. Accurate GFM of the topography will require the DTM data to be largely free of artefacts. In order to approximate the actual gravity field

closely, detailed compilations of mass-density values are important to improve the quality of the DTM-based mass models.

Summary

Gravity forward modelling (GFM) comprises methods for the computation of the gravity field of some mass distribution, very often of the topography, but also from water, ice and crustal masses. GFM in the spatial and spectral domain are important for gravity applications in physical geodesy (e.g., gravity reduction, interpolation and prediction, e.g., for the construction of detailed gravity maps) and potential field or planetary geophysics (reduction and interpretation of gravity observations).

Cross references

Topographic Effects, Digital Terrain Models, Spherical Harmonic Models, The Remove Restore Method, Gravity Field of the Planets, Gravity Anomalies

Bibliography

Asgharzadeh M.F., von Frese, R.B., Kim, H.R., Leftwich, T.E., and Kim, J.W., 2007. Spherical prism gravity effects by Gauss-Legendre quadrature integration. *Geophysical Journal International*: 169, 1-11.

Baran I., Kuhn, M., Claessens, S.J., Featherstone, W.E., Holmes, S.A., Vaníček, P., 2006. A synthetic Earth Gravity Model designed specifically for testing regional gravimetric geoid determination algorithms, *Journal of Geodesy*: 80, 1-16.

Blakeley, R.J. (1996), *Potential Theory in Gravity and Magnetic Applications*. Cambridge: Cambridge University Press.

Bagherbandi M., and Sjöberg L.E., 2012. A synthetic Earth gravity model based on a topographic-isostatic model. *Studia Geophysica et Geodetica*: 56, 935-955.

Balmino, G., Vales, N., Bonvalot S., and Briais, A., 2012. Spherical harmonic modelling to ultra-high degree of Bouguer and isostatic anomalies. *Journal of Geodesy*: 86, 499-520.

Claessens, S.J., and Hirt, C., 2013. Ellipsoidal topographic potential - new solutions for spectral forward gravity modelling of topography with respect to a reference ellipsoid. *Journal of Geophysical Research – Solid Earth*: 118, 5991-6002.

D'Urso, M. G., 2014. Analytical computation of gravity effects for polyhedral bodies. *Journal of Geodesy*: 88, 13-29.

Ebbing, J., Braitenberg, C., Götze, H.-J., 2001. Forward and inverse modelling of gravity revealing insight into crustal structures of the Eastern Alps. *Tectonophysics*: 337, 191-208.

Eshagh, M., 2009. Comparison of two approaches for considering laterally varying density in topographic effect on satellite gravity gradiometric data. *Acta Geoph.*: 58, 661-686.

Forsberg, R., and Tscherning, C.C., 1981. The use of height data in gravity field approximation by collocation. *Journal Geophysical Research* 86(B9): 7843–7854.

Forsberg R., 1984. A study of terrain reductions, density anomalies and geophysical inversion methods in gravity field modelling. Report 355, Depart. Geodetic Science Surveying, Ohio State University, Columbus.

Göttl, F., and Rummel, R., 2009. A Geodetic View on Isostatic Models. *Pure and Applied Geophysics*: 166, 1247-1260.

- Grombein T., Seitz, K., and Heck, B., 2013. Optimized formulas for the gravitational field of a tesseroid. *Journal of Geodesy*: 87, 645–660.
- Grombein, T., Luo, X., Seitz, K., and Heck, B., 2014. A wavelet-based assessment of topographic-isostatic reductions for GOCE gravity gradients. *Surveys in Geophysics*: 35, 959-982.
- Gruber C., Novák P., Flechtner, F., and Barthelmes, F., 2013. Derivation of the topographic potential from global DEM models. In *International Association of Geodesy Symposia Series 139*, 535-542. Springer, Berlin Heidelberg.
- Heck, B., and Seitz, K., 2007. A comparison of the tesseroid, prism and point-mass approaches for mass reductions in gravity field modelling. *Journal of Geodesy*: 81, 121-136.
- Hirt C., and Kuhn, M., 2014. A band-limited topographic mass distribution generates a full-spectrum gravity field – gravity forward modelling in the spectral and spatial domain revisited. *Journal of Geophysical Research – Solid Earth*: 119, 3646–3661.
- Hirt, C., Claessens S.J., Fecher, T., Kuhn, M., Pail, R., and Rexer, M., 2013. New ultra-high resolution picture of Earth's gravity field. *Geophysical Research Letters*: 40, 4279-4283.
- Hirt C., Kuhn M., Featherstone W., and Göttl, F., 2012. Topographic/isostatic evaluation of new-generation GOCE gravity field models. *Journal of Geophysical Research - Solid Earth*: 117, B05407.
- Hirt C., 2010. Prediction of vertical deflections from high-degree spherical harmonic synthesis and residual terrain model data. *Journal of Geodesy*: 84, 179-190.
- Jacoby, W., and Smilde, P.L., 2009. *Gravity interpretation*. New York: Springer.
- Jekeli, C. and Serpas, J.G., 2003. Review and numerical assessment of the direct topographical reduction in geoid determination. *Journal of Geodesy*: 77, 226–239.
- Kuhn, M., and Seitz, K., 2005. Comparison of Newton's Integral in the Space and Frequency Domains. In *International Association of Geodesy Symposia Series 128*, 386-391.
- Kuhn, M., Featherstone, W.E., and Kirby, J.F., 2009. Complete spherical Bouguer gravity anomalies over Australia. *Australian Journal of Earth Sciences*: 56, 213-223.
- Makhloof, A.A., and Ilk, K.-H., 2008. Effects of topographic–isostatic masses on gravitational functionals at the Earth's surface and at airborne and satellite altitudes. *Journal of Geodesy*: 82, 93–111.
- Mohr, P. J., Taylor B.N., and Newell, D.B., (2012), CODATA recommended values of the fundamental physical constants: 2010. *Reviews of Modern Physics*: 84, 1527-1605. [Values available from <http://physics.nist.gov/constants>, last accessed on Sep 09, 2014]
- Nahavandchi, H., and Sjöberg, L.E., 1998. Terrain correction to power H3 in gravimetric geoid determination. *Journal of Geodesy*: 72, 124-135.
- Nagy, D., Papp, G., and Benedek, J., 2000. The gravitational potential and its derivatives for the prism. *Journal of Geodesy*: 74, 552-560, Erratum in *Journal of Geodesy*: 76, 475.
- Neumann, G.A., et al., 2004. Crustal structure of Mars from gravity and topography. *Journal of Geophysical Research: Planets*: 109(E8), 1-18.
- Novák, P., 2010. Direct modelling of the gravitational field using harmonic series. *Acta Geodyn. Geomater.*: 7, 35-47.
- Oldenburg, D.W., 1974. The inversion and interpretation of gravity anomalies. *Geophysics*: 39, 526-536.
- Papp, G., 1996. Gravity field approximation based on volume element model of the density distribution. *Acta Geod. Geoph. Hung.*, 91, 339-358.

- Pavlis, N.K., Factor, J.K., Holmes, S.A., 2007. Terrain-related gravimetric quantities computed for the next EGM. In *Proceed. of the 1st International Symposium of the International Gravity Field Service*, 318-323, Harita Dergisi, Istanbul, Turkey.
- Rummel, R., Rapp, R.H., Sünkel, H. and Tscherning, C.C., 1988. Comparisons of global topographic/isostatic models to the Earth's observed gravity field, Report No 388, Dep. Geodetic Sci. Surv., Ohio State University, Columbus, Ohio.
- Smith, D. A., 2000. The gravitational attraction of any polygonally shaped vertical prism with inclined top and bottom faces, *Journal of Geodesy*: 74, 414– 420.
- Tenzer, R., 2005. Spectral domain of Newton's integral, *Boll Geod. Sci. Affini*, 2, 61-73.
- Tenzer, R., Novák, P., and Gladkikh, V., 2011. On the accuracy of the bathymetry-generated gravitational field quantities for a depth-dependent seawater density distribution. *Studia Geophysica et Geodaetica*: 55, 609-626.
- Tenzer, R., Gladkikh, V., Novák, P., Vajda, P., 2012. Spatial and Spectral Analysis of Refined Gravity Data for Modelling the Crust–Mantle Interface and Mantle–Lithosphere Structure. *Surveys in Geophysics* 33, 817-839.
- Tocho, C., Vergos, G.S., and Sideris, M.G., 2012. Investigation of topographic reductions for marine geoid determination in the presence of an ultra-high resolution reference geopotential model. In *International Association of Geodesy Symposia Series 136*, 419-426, Springer Berlin, New York.
- Tsoulis, D., Novák, P., and Kadlec, M., 2009. Evaluation of precise terrain effects using high-resolution digital elevation, models, *J. Geophys. Res.*, 114, B02404.
- Tsoulis, D., 2013. Geodetic use of global digital terrain and crustal databases in gravity field modeling and interpretation. *J Geod Sci* 1, 1-6.
- Tziavos, I.N., and Sideris, M.G., 2013. Topographic Reductions in Gravity and Geoid Modeling. In *Lecture Notes in Earth System Sciences* 110, 337-400, Springer, Berlin Heidelberg.
- Tziavos, I.N., Vergos, G.S. and Grigoriadis, V.N., 2010. Investigation of topographic reductions and aliasing effects to gravity and the geoid over Greece based on various digital terrain models. *Surveys in Geophysics* 31, 23-67.
- Wang, Y.M, and Yang, X., 2013. On the spherical and spheroidal harmonic expansion of the gravitational potential of the topographic masses. *Journal of Geodesy*: 87, 909-921.
- Wieczorek, M.A., 2007. Gravity and topography of the terrestrial planets. In *Treatise on Geophysics*, 10, 165-206, Elsevier-Pergamon, Oxford, United Kingdom.
- Wild-Pfeiffer F., 2008. A comparison of different mass elements for use in gravity gradiometry, *J. Geod.* 82, 637-653.
- Zuber, M. T., et al., 2013. Gravity field of the Moon from the Gravity Recovery and Interior Laboratory (GRAIL) mission, *Science*, 339, 668-67.