

## CASE REPORT

# Ketogenic diet for treating alopecia in BCS11-related mitochondrial disease (Bjornstad syndrome)

Adela Della Marina<sup>1</sup> | Baerbel Leiendecker<sup>1</sup> | Sebastian Roesch<sup>2</sup> |  
Saskia B. Wortmann<sup>3,4,5,6</sup>

<sup>1</sup>Department of Neuropediatrics, Developmental Neurology and Social Pediatrics, University of Duisburg-Essen, Children's Hospital, Essen, Germany

<sup>2</sup>Department of Otorhinolaryngology, Head and Neck Surgery, Paracelsus Medical University, Salzburg, Austria

<sup>3</sup>University Children's Hospital, Paracelsus Medical University, Salzburg, Austria

<sup>4</sup>Institute of Human Genetics, Technical University München, Munich, Germany

<sup>5</sup>Institute of Human Genetics, Helmholtz Zentrum, Neuherberg, Germany

<sup>6</sup>Radboud Center for Mitochondrial Medicine, Department of Pediatrics, Amalia Children's Hospital, Radboudumc, Nijmegen, The Netherlands

## Correspondence

Saskia B. Wortmann, University Children's Hospital, Paracelsus Medical University (PMU), Salzburg, Austria.

Email: s.wortmann@salk.at

**Communicating Editor:** Avihu Boneh



**FIGURE 1** Patient with BCS11L related mitochondrial disease (Bjornstad syndrome) 2 months before (A), at day of starting (B) and after three (C) and four (D, E) months of modified Atkins diet. (F) Shows the patient 6 months after cessation of modified Atkins diet

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Inherited Metabolic Disease* published by John Wiley & Sons Ltd on behalf of SSIEM.

The human *BCS1L* gene encodes a homolog of *Saccharomyces cerevisiae* *bcs1* protein involved in the assembly of complex III of the mitochondrial respiratory chain. Phenotypes associated with pathogenic biallelic *BCS1L* variants range from growth retardation, aminoaciduria, cholestasis, iron overload, lactic acidosis and early death (GRACILE) syndrome; MIM#603388) to Bjornstad syndrome (MIM#262000), characterized by abnormal flattening and twisting of hair shafts (*pili torti*) and sensori-neural hearing loss without intellectual impairment. No causal treatment for this mitochondrial disease is available.

Ketogenic diets stimulate mitochondrial biogenesis, improve mitochondrial function and decrease oxidative stress<sup>1-4</sup> and have been implemented in some patients with mitochondrial disease and epilepsy.<sup>5-7</sup>

We report on a 7-year-old female (*BCS1L*, NM\_001079866.2, c.232A>G; p.Ser78Gly, c.794G>A; p.Arg265Gln) with very sparse and brittle hair since birth (Figure 1A,B), with progressive sensori-neural hearing loss noted from age 18 months onwards. A 3-day dietary history indicated an adequate diet with no protein deficiency but aminoaciduria, as a possible contributing cause for alopecia, was not confirmed or excluded. We initiated a mild ketogenic diet (modified Atkins diet [MAD] with 30-40 g/day carbohydrates [1500 kcal/day], 10% of energy from carbohydrates, 25% from protein and 65% from fat). This was well tolerated and blood ketone levels of 2-4 mmol/L were achieved.<sup>8</sup> Hair growth improved during treatment (Figure 1C- E). Despite this, the family decided to go back to a regular diet after 4 months of diet because of the child craving for carbohydrates, and lack of improvement of hearing. Figure 1F shows that the hair was lost again 6 months after cessation of MAD.

There was improved hair growth during MAD in this individual with *BCS1L* related mitochondrial disease, We can only speculate if continuation of MAD would have

further increased the hair growth or would have influenced the hearing loss and what effects an earlier initiation of MAD would have had.

## REFERENCES

- Stafford P, Abdelwahab MG, Kim do Y, Preul MC, Rho JM, Scheck AC. The ketogenic diet reverses gene expression patterns and reduces reactive oxygen species levels when used as an adjunct therapy for glioma. *Nutr Metab*. 2010;7:74.
- Ahola-Erkilä S, Carroll CJ, Peltola-Mjösund K, et al. Ketogenic diet slows down mitochondrial myopathy progression in mice. *Hum Mol Genet*. 2010;19:1974-1984.
- Bough KJ, Wetherington J, Hassel B, et al. Mitochondrial biogenesis in the anticonvulsant mechanism of the ketogenic diet. *Ann Neurol*. 2006;60:223-235.
- Daniel NN, Hartman AL, Stafstrom CE, Thio LL. How does the ketogenic diet work? Four potential mechanisms. *J Child Neurol*. 2013;28:1027-1033.
- Santra S, Gilkerson RW, Davidson M, Schon EA. Ketogenic treatment reduces deleted mitochondrial DNAs in cultured human cells. *Ann Neurol*. 2004;56:662-669.
- Joshi CN, Greenberg CR, Mhanni AA, Salman MS. Ketogenic diet in Alpers-Huttenlocher syndrome. *Pediatr Neurol*. 2009;40:314-316.
- Kang HC, Lee YM, Kim HD, Lee JS, Slama A. Safe and effective use of the ketogenic diet in children with epilepsy and mitochondrial respiratory chain complex defects. *Epilepsia*. 2007;48:82-88.
- Kossoff EH, Zupec-Kania BA, Auvin S, et al. Optimal clinical management of children receiving dietary therapies for epilepsy: updated recommendations of the International Ketogenic Diet Study Group. *Epilepsia Open*. 2018;3(2):175-192.

**How to cite this article:** Della Marina A, Leiendecker B, Roesch S, Wortmann SB. Ketogenic diet for treating alopecia in *BCS1L*-related mitochondrial disease (Bjornstad syndrome). *JIMD Reports*. 2020;53:10–11. <https://doi.org/10.1002/jimd.12109>