







# Allergen-free extracts from birch, ragweed, and hazel pollen activate human and guinea-pig submucous and spinal sensory neurons

Sabine Buhner<sup>1</sup>  | Stefanie Schäuffele<sup>1</sup> | Pieter Giesbertz<sup>2</sup>  | Ihsan Ekin Demir<sup>3</sup>  | Florian Zeller<sup>4</sup> | Claudia Traidl-Hoffmann<sup>5,6,7</sup>  | Michael Schemann<sup>1</sup>  | Stefanie Gilles<sup>5,6</sup> 

<sup>1</sup>Chair of Human Biology, Technical University Munich, Freising, Germany

<sup>2</sup>Molecular Nutrition Unit, Technical University Munich, Freising, Germany

<sup>3</sup>University Hospital Rechts der Isar, Technical University Munich, Munich, Germany

<sup>4</sup>Department of Surgery, Academic Hospital Freising, Freising, Germany

<sup>5</sup>Environmental Medicine, Faculty of Medicine, University of Augsburg, Augsburg, Germany

<sup>6</sup>Institute of Environmental Medicine, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

<sup>7</sup>Christine Kühne Center for Allergy Research and Education (CK-Care), Davos, Switzerland

## Correspondence

Michael Schemann, Chair of Human Biology, Technical University Munich, Liesel Beckmann Strasse 4, 85354 Freising, Germany.  
Email: [schemann@mytum.de](mailto:schemann@mytum.de)

## Funding information

Christine Kühne Research Center for Allergy and Education (CK-Care)

Listen to the podcast [Here](#)

## Abstract

**Background:** Non-allergenic, low molecular weight components of pollen grains are suspected to trigger changes in gut functions, sometimes leading to inflammatory conditions. Based on extensive neuroimmune communication in the gut wall, we investigated the effects of aqueous pollen extracts (APE) on enteric and spinal sensory neurons.

**Methods:** Using  $\text{Ca}^{2+}$  and fast potentiometric imaging, we recorded the responses of guinea-pig and human submucous and guinea-pig dorsal root ganglion (DRG) neurons to microinjection of low (<3kDa) and high ( $\geq$ 3kDa) molecular weight APEs of birch, ragweed, and hazel. Histamine was determined pharmacologically and by mass spectrometry (LC-MS/MS).

**Key Results:** Birch  $\text{APE}_{<3\text{kDa}}$  evoked strong  $[\text{Ca}^{+2}]_i$  signals in the vast majority of guinea-pig DRG neurons, and in guinea-pig and human enteric neurons. The effect of birch  $\text{APE}_{\geq 3\text{kDa}}$  was much weaker. Fast neuroimaging in human enteric neurons revealed an instantaneous spike discharge after microinjection of birch, ragweed, and hazel  $\text{APE}_{<3\text{kDa}}$  [median (interquartile range) at 7.0 Hz (6.2/9.8), 5.7 Hz (4.4/7.1), and 8.4 Hz (4.3/12.5), respectively]. The percentage of responding neurons per ganglion were similar [birch 40.0% (33.3/100.0), ragweed 50.8% (34.4/85.6), and hazel 83.3% (57.1/100.0)]. A mixture of histamine receptor (H1-H3) blockers significantly reduced nerve activation evoked by birch and ragweed  $\text{APE}_{<3\text{kDa}}$ , but was ineffective on hazel. Histamine concentrations in ragweed, birch and hazel APE's <3kDa were 0.764, 0.047, and 0.013  $\mu\text{M}$ , respectively.

**Conclusions:** Allergen-free APEs from birch, ragweed, and hazel evoked strong nerve activation. Altered nerve-immune signaling as a result of severe pollen exposure could be a pathophysiological feature of allergic and non-allergic gut inflammation.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](#) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Neurogastroenterology & Motility* published by John Wiley & Sons Ltd.

## KEYWORDS

allergy, enteric nervous system, gastrointestinal tract, inflammation, neurogastroenterology, pollen sensitization

## 1 | INTRODUCTION

The last years we have seen increasing evidence for the relevance of allergic sensitization to pollen in chronic inflammatory conditions in the gastrointestinal tract, for example, eosinophilic esophagitis (EoE)<sup>1-6</sup> for review.<sup>7,8</sup> Similarly in regard to Crohn's disease<sup>9</sup> and the irritable bowel syndrome (IBS), pollen sensitization is frequently discussed as a potential factor in the pathogenesis,<sup>10-12</sup> for review.<sup>13</sup> The underlying mechanisms are complex and only partly understood. They may include hypersensitivity responses to pollen cross-reacting foods, the so-called pollen-food-allergy or oral-allergy-syndrome,<sup>14,15</sup> as well other pro-inflammatory effects of pollen exposure.<sup>2,4,5,16,17</sup> Pollen grains, per se, trigger local inflammatory actions in esophageal and/or gastrointestinal mucosa, very similar to effects in the nasal mucosa.<sup>1</sup> This is supported by studies using radiolabeled pollen grains.<sup>18</sup> It was shown that the vast majority (nearly 100%) of pollen grains and high molecular weight pollen extracts in humans do not reach the thoracic respiratory tract but were swallowed and had direct contact with the mucosa of the digestive tract.<sup>18</sup> This is in line with findings from a recent study demonstrating *in situ* pollen germination in 65.6% of esophageal mucosal biopsies obtained from patients suffering from eosinophilic esophagitis.<sup>17</sup> The authors hypothesized that alteration of the mucosal barrier may facilitate the penetration of pollen grains into deeper layers of the gut wall, thus emphasizing the role of pollen in local inflammation.<sup>17</sup>

The intestinal and respiratory lining are packed with immune cells, such as mast cells, which functionally interact with enteric and spinal sensory nerves. Recently, we have demonstrated in detail signaling between nerves and mast cells at the cellular level in intact human intestinal mucosa.<sup>19</sup>

The complexity of pollen effects is not only a simple function of its allergen content triggering allergic hypersensitivity reactions.<sup>20,21</sup> Along with allergens, pollen grains release multiple other substances that are not allergenic themselves, but act as essential adjuvants, facilitating and enhancing allergic immune responses.<sup>22</sup> These are mainly water-soluble components of low molecular weight (MW) of which only a limited number have been chemically and functionally identified to date. One of these substances is histamine. It was suspected early on that in the case of non-immunological effects of pollen extracts, histamine in particular could be responsible for false-positive reactions in skin tests.<sup>23,24</sup> In an extensive 1992 study, the authors detected histamine at concentrations up to 7.4 µg/mL in 48 of 49 aqueous or phenolic pollen extracts.<sup>25</sup> From a neurophysiological point of view, histamine is interesting because, as we have shown, it has strong nerve excitatory properties in enteric neurons in both humans and rodents.<sup>26</sup> This is one reason why we initially focused on this substance in the present study. Other components

### Key points

- Non-allergenic components of pollen grains are suspected to trigger changes in gut functions, sometimes leading to inflammatory conditions. We investigated the effects of low molecular aqueous pollen extracts on enteric and spinal sensory neurons from humans and guinea-pigs.
- Ca<sup>2+</sup> and fast potentiometric imaging techniques were used.
- Low molecular extracts of birch, ragweed, and hazel pollen strongly excite neurons in humans and guinea-pigs; nerve excitation was mediated by histamine in birch and ragweed, but not in hazel pollen extracts.
- Altered nerve signaling as a result of severe pollen exposure may be a pathophysiological feature to allergic and non-allergic challenges.

are, for example, adenosine,<sup>27</sup> which is known to modulate sensory signaling<sup>28</sup> or phytoprostanes which are structurally related to prostaglandins.<sup>29-32</sup> In these studies, 3 kDa was used as the cutoff value for the separation of the allergenic protein fractions. This was due to the fact that proteins, in order to act as allergens, that is, to form an antibody bond, must have a certain minimum size of about 30 amino acid residues (molecular weight about 3 kDa).<sup>33</sup> For comparison, the molecular weights of the common main pollen allergens of birch and hazel are about 17 kDa,<sup>34,35</sup> those of ragweed allergens are 21–31 kDa.<sup>36</sup>

In human airway epithelial cells, low MW fractions of birch, ragweed, and mugwort pollen were found to modulate immune responses and barrier functions.<sup>32,37-39</sup> Additionally, activation of dendritic cell function had been shown.<sup>30,31,40</sup> In a recent study, low MW extract from a.o. birch pollen was shown to compromise the innate antiviral defense of respiratory epithelia during infection with human rhinovirus.<sup>39</sup> Taken together, these results lead to the suggestion that low MW non-allergenic pollen components are responsible for a bundle of symptoms and modulation of innate immune defenses by pollen exposure in both allergic and non-allergic humans.<sup>39</sup>

Whether and, if so, which components of the pollen extracts might play a role in the activation of neurons, particularly in the gastrointestinal wall, is not yet known. Therefore, we performed experiments using different neuronal settings, that is, cell cultures of guinea-pig spinal sensory neurons as well as whole mount preparations of guinea-pig and human intestinal submucous plexus. We recorded on single-cell level slow and fast neuronal responses to

aqueous extracts of birch, ragweed, and hazel pollen. We separated the extracts according to their molecular weight and found that the low MW (protein-free) fractions of all pollen species strongly activated neurons. Using pharmacological methods and mass spectrometry, we identified histamine as a neuroactive component in low MW pollen extracts.

## 2 | METHODS

### 2.1 | Ethics statement

Procedures for the work with human gut samples were approved by the local ethics committee of the Technische Universität München (project approval 5242/11). Informed written consent was obtained from all subjects, and the studies conformed to the standards set by the Declaration of Helsinki except for registration in a database.

All animal work with guinea-pigs was approved by the Animal Welfare Commissioner of TUM School of Life Sciences (reference number 32-568-2) and are in line with methods stated in Annex IV of Directive 2010/63/EU and German animal welfare law and conform to the principles and regulations as described in the "Principles and standards for reporting animal experiments".<sup>41</sup>

### 2.2 | Human tissue samples

Studies were performed using macroscopically normal surgical intestinal specimens of human colon ( $n=19$ ), ileum ( $n=11$ ), jejunum ( $n=3$ ), or duodenum ( $n=4$ ), obtained from 37 patients (19 male, 18 females, mean age  $67 \pm 14$ ) undergoing abdominal surgery at the Medical Clinic in Freising and the Medical Clinic of the Technische Universität München. Diagnoses that led to the surgery were as follows: colon carcinoma/adenoma (16/1), diverticular disease (4), pancreas carcinoma/pancreatitis (3/2), stenosis (3), Crohn's Disease (2), and one each for stoma reversal, small intestinal carcinoma, leukemia, gangrene, ischemia, and unknown diagnosis. Samples were taken from macroscopically normal, unaffected areas as determined by visual inspection by the pathologists. After removal, the surgical specimens were placed in cold aerated sterile HEPES-Krebs solution containing in mM: 135 NaCl, 5.4 KCl; 1.0 Mg Cl<sub>2</sub>·6H<sub>2</sub>O, 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 1.25 CaCl<sub>2</sub>·2H<sub>2</sub>O, 12.2 Glucose, and 3 HEPES (all from Sigma-Aldrich Chemie GmbH) and 10 mL/L antibiotic-antimycotic mix (mg/L: 25 amphotericin B, 107 U/L penicillin G, 10,000 streptomycin in physiological saline; CCPro). They were immediately transported to the laboratory for experiments. Under continuous superfusion with ice-cold carbogen-aerated Krebs solution (pH 7.4) containing in mM: 117 NaCl, 4.7 KCl, 1.2 MgCl<sub>2</sub>·6H<sub>2</sub>O, 1.2 NaH<sub>2</sub>PO<sub>4</sub>, 25 NaHCO<sub>3</sub>, 2.5 CaCl<sub>2</sub>·2H<sub>2</sub>O, and 11 glucose (all from Sigma), the surgical specimens were microscopically dissected by removing the mucosa and the muscular layers to obtain a whole mount preparation of the inner submucosal plexus. The final preparations (10×20 mm) were pinned on silicone rings.<sup>42</sup>

### 2.3 | Guinea-pig specimens

Eleven male "Dunkin-Hartley" guinea-pigs (strain HsdDhl:DH, Envigo GmbH; strain 051, Charles River) of an average weight of  $390 \pm 49$  g were used. Animals were killed by a percussive blow to the head followed by exsanguination. From two animals, DRGs were isolated under a stereomicroscope and collected in a small glass bottle filled with HEPES buffer containing in mM: 1 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 150 NaCl, 10 HEPES, 10 Glucose, and 5 KCl. They were taken from thoracic segments T2–L2 because in rodents, these ganglia provide the major splanchnic afferent innervation to bronchia, lung, and intestine.<sup>43</sup> The DRGs were digested in protease type I (8 U/mL, Gibco), collagenase type II (516.2 U/mL, Gibco) and bovine serum albumin (BSA fraction V 0.37%, Serva). The incubation step lasted 40–50 min. After several washing steps, the pellet was suspended in 600–1200  $\mu$ L culture medium (M199 GlutaMAX, Gibco/Invitrogen) containing 1% Penicillin/Streptomycin (Gibco/Invitrogen), 30 mM glucose, and 10% fetal bovine serum (Gibco/Invitrogen). Each culture dish was inoculated with 300  $\mu$ L ganglia suspension. For the present experiments, we used three culture dishes (two from one guinea-pig and one from the other; ibidi dish 35 mm with ibiTreatment, ibidi). Freshly dissociated and short-term cultured DRG neurons often became dislocated after local application of substances. DRGs were therefore maintained in culture for at least 7 days during which the neurons adhered to the optical bottom of the dish. Medium was changed every 2–3 days and contained 2  $\mu$ M cytosine  $\beta$ -D-arabinofuranoside (Ara-C; Sigma) to suppress proliferation of other cell types.

Submucous plexus activity was assessed on segments of the proximal part of the distal colon (length 2 cm), which were quickly removed after killing the animals. The segments were dissected in ice-cold carbogen-aerated Krebs solution similarly to the procedure described above for human tissue. The mucosa and the muscle layers were carefully removed for preparation of the submucous plexus. The final preparations (10×20 mm) were pinned on silicone rings.

### 2.4 | Ca<sup>2+</sup> sensitive dye imaging

To initially screen for neuronal effects of the aqueous pollen extracts, we used Ca<sup>2+</sup> imaging technique as a common method to assess long-lasting changes in intracellular Ca<sup>2+</sup> levels ([Ca<sup>2+</sup>]<sub>i</sub>). The methods and techniques have been previously described in the enteric nervous system,<sup>19,44</sup> as well as for DRG neurons.<sup>45</sup> Briefly, the primary culture dishes of DRG neurons were incubated for 20 min at room temperature in the dark with the fluorescent calcium indicator Fluo 4-acetoxymethyl (AM) (Invitrogen) at a final concentration of 10  $\mu$ M diluted in HEPES buffer containing 1.25 mM probenecid (Sigma) to prevent dye leakage. The staining solution was washed out for 10 min, and the preparation was mounted in a self-made recording chamber and continuously perfused with 37°C HEPES buffer. For human and guinea-pig submucous plexus preparations, the staining procedure was similar to that of the DRG neurons,

except a longer incubation period (45 min) and the use of Krebs buffer containing 500  $\mu\text{M}$  probenecid instead of HEPES buffer for incubation, washing, and perfusion during the experiments.

The recording chambers were mounted on an inverted epifluorescence microscope (Zeiss Axio Observer A1, Carl Zeiss) equipped with a high-speed monochrome camera (Zeiss AxioCam HSm) and software (Zeiss Axio Vision 4.8) for acquisition and analysis. Fluo-4AM was excited using a blue light emitting diode (LED) Luxeon III (3W, 470 nm dominant wavelength, Phillips Lumiled, Phillips), and the signals were detected with a filter cube F26-514 Bright Line FITC BP (excitation: HC475/35, dichroic: 499, emission: HC530/43, AHF Analysentechnik) using 20 $\times$  objective (A-Plan, NA=0.25, Zeiss).<sup>46</sup> The system measured relative changes in fluorescence ( $\Delta F/F$ ) of Fluo-4AM by monitoring changes in  $[\text{Ca}^{2+}]_i$ . Depending on the stimulus,  $[\text{Ca}^{2+}]_i$  transients were recorded for 35 s up to 60 s using a frame rate of 0.5 or 1 Hz, in selected cases 6.2 Hz. Under basal condition, the variation in background fluorescent was  $\pm 1\% \Delta F/F$ . A threefold standard deviation (3%) was defined as threshold for genuine cell activation.<sup>19</sup> Basal, non-stimulated  $[\text{Ca}^{2+}]_i$  signals were recorded for 35 s.

Aliquots of low and high MW fractions of aqueous birch pollen extract (diluted 1:1 with buffer) were locally perfused on DRG neuronal clusters or submucous neurons, respectively, via a micro perfusion pump (Micro4, WPI Sarasota FL) lasting for 30 s at a speed of 0.1  $\mu\text{L/s}$ . According to our previous studies using spritz pulse application, a 1:10 dilution of the applied substance has to be considered for the local perfusion as well.<sup>26</sup>

To check the vitality of submucous neurons, we stimulated the entire preparation via transmural electrical field stimulation (EFS<sub>TM</sub>).<sup>19</sup> The EFS<sub>TM</sub> of the nerves was achieved by two platinum electrodes and a constant-voltage stimulator (Typ 215/I, Hugo Sachs Elektronik, Hayard-Apparatus GmbH). The stimulus parameters were 50 Hz pulse train for 10 s with pulse duration of 0.5 ms at 25 V. Cultured cell vitality was tested at the end of each experiment using the calcium ionophore ionomycin (15  $\mu\text{M}$ , Sigma) which was applied for 2 s followed by 60 s recording period. Those cells showing no  $\text{Ca}^{2+}$  signals in response to ionomycin were excluded from further analysis.

## 2.5 | Fast neuroimaging technique

Fast neuroimaging allowed detection of action potential discharge at a single-cell level and was used in preparations of human submucous plexus only. This multisite optical recording technique (MSORT) was previously described in detail.<sup>47-49</sup> Briefly, individual ganglia were stained with the fluorescent voltage-sensitive dye 1-(3-sulfonatopropyl)-4- $[\beta$ -(2-(di-*n*-octylamino)-6-naphthyl)vinyl]pyridinium betaine (di-8-ANEPPS, 20  $\mu\text{M}$  dissolved in DMSO (0.135%) and Pluronic F-127 (0.0135%) containing Krebs buffer; Molecular Probes Mobitec). The dye was excited by a green LED (LET S2W, Osram), and the recordings from the stained neurons were performed using an epifluorescence Olympus IX 50 microscope (Olympus Corporation)

equipped with a 40 $\times$  oil immersion objective (UApo 340 nm, Olympus) and an appropriate filter cube (AHF Analysentechnik AG). The technique allows for measurement of the relative changes in fluorescence intensity ( $\Delta F/F$ ), which is linearly related to changes in the membrane potential.<sup>47</sup> Signals were acquired with a frequency of 1 kHz, detected using a cooled charge coupled device (CCD) camera (80 $\times$ 80 pixels; Neuro CCD-SMQ imaging system, Red Shirt Imaging LLC Decatur) and processed by the Neuroplex 10.1.2 software (RedShirt Imaging). With the 40 $\times$  oil objective, the spatial resolution of the CCD camera was 24  $\mu\text{m}^2/\text{pixel}$ .

The viability and the staining of the neurons were tested either by recording spontaneous neuronal activity or by evoking fast excitatory postsynaptic potentials with a single electrical stimulus applied to an interganglionic fiber tract via<sup>42</sup> a teflon-coated platinum electrode (diameter 25  $\mu\text{m}$ , Science products GmbH) connected to a constant-voltage stimulator (A360, WPI; 400  $\mu\text{s}$  pulse duration, amplitudes ranging from 10 to 100  $\mu\text{A}$ ). Aqueous pollen extracts (diluted 1:1 with Krebs buffer) were applied onto a single submucous ganglion by pressure pulse ejection (20 PSI, ejection speed  $55 \pm 27 \text{ nL/s}$  400–800 ms duration) from glass microejection pipettes that were placed close to the ganglion (approximate distance: 100–200  $\mu\text{m}$ ). Signals were recorded for a 2000 ms period. For statistical analysis and comparison, only experiments with an application duration of 800 ms were used.

## 2.6 | Aqueous pollen extracts and pharmacology

Pollen from birch (*Betula pendula*) and hazel (*Corylus avellana*) was isolated from flowering catkins in the early pollen seasons of 2018 (birch) and 2019 (hazel). Pollen from Ragweed (*Ambrosia artemisiifolia*) was isolated from inflorescences of flowering *Ambrosia* plants grown in the greenhouse at the Helmholtz Center Munich (kindly provided by Dr. Ulrike Frank, Institute of Biochemical Plant Pathology). Aqueous pollen extracts (APEs) were prepared as described before.<sup>39,50</sup> Briefly, pollen grains were suspended in HEPES-Krebs buffer without supplements at 30 mg pollen per ml. After incubation in a shaking water-bath (250 rpm) at 37 $^{\circ}\text{C}$  with agitation for 30 min and vigorous vortexing every 10 min, the suspensions were centrifuged (10 min, 3000 $\times g$ ), sterile filtered (0.2  $\mu\text{m}$  pore-size) and the pollen-free supernatant (=aqueous pollen extract) stored in aliquots at  $-80^{\circ}\text{C}$  until use. As previously described in detail,<sup>22,37</sup> aliquots of the pollen extracts were separated by ultrafiltration using centrifugal filter units (Amicon Ultra) into two fractions: a low MW, protein-free fraction (APE<sub><3kDa</sub>), which is rich in small molecules, such as lipids, sugars, nucleosides, and secondary plant metabolites,<sup>40</sup> and a high MW fraction (APE <sub>$\geq$ 3kDa</sub>), which contains the pollen allergens and other proteins, but also residual small molecules that remain bound to proteins. Before use, the samples were diluted 1:1 with Krebs buffer and adjusted to pH 7.4, if necessary. The osmolarity was 291 mosmol/kg for birch APE<sub><3kDa</sub>, 286 mosmol/kg for birch APE <sub>$\geq$ 3kDa</sub>, 298 mosmol/kg for ragweed APE<sub><3kDa</sub>, and 296 mosmol/kg for hazel APE<sub><3kDa</sub>; the osmolarity of Krebs buffer

was 296 mosmol/kg. In some experiments, we tested the effects of APE<sub><3kDa</sub> in the presence of a mixture of specific histamine receptor antagonists, that is, the H1 receptor antagonist pyrilamine (1 μmol/L; Sigma), the H2 antagonist ranitidine (10 μmol/L; Sigma), and the H3 antagonist clobenpropit (10 μmol/L; Biotrend)<sup>26</sup> or in the presence of the neurotoxin tetrodotoxin (TTX 0.5 μM). Antagonists were added to the perfusing Krebs solution 20 mins before reapplication of the respective pollen extract. The washout periods were at least 60 mins.

## 2.7 | Histamine analysis in the aqueous pollen extracts

Histamine concentrations in the different fractions of the APEs of birch, ragweed, and hazel pollen (APE<sub><3kDa</sub>, APE<sub>2-3kDa</sub> and total APE) were quantified using targeted liquid chromatography tandem-mass spectrometry (LC-MS/MS). Therefore, 50 mL of the extracts was mixed with 10 μL of internal standard mixture, consisting of 0.5 μM Histamine-d4 (internal standards, Santa Cruz Biotechnology). Samples were subsequently dried in a SpeedVac vacuum concentrator. Our method is described in detail in.<sup>19</sup> Briefly, pre-column derivatization of all samples was done using phenylisothiocyanate (PITC). A 200 μL mixture consisting of 5% PITC dissolved in pyridine, ethanol, and 0.1% NH<sub>3</sub> in a 1/1/1 ratio was added to the dried samples. Samples were shaken at 25°C for 20 min and were subsequently dried and then resuspended in methanol containing 5 mM ammonium acetate and finally diluted with water to reach a methanol/water ratio of 70:30. The analysis was performed on a triple quadrupole QTRAP 5500 LC-MS/MS system operating in positive ESI mode (AB Sciex) equipped with a 1200 series binary pump (Agilent) and coupled to an HTC pal autosampler (CTC Analytics). Chromatographic separation was achieved using a Zorbax Eclipse XDB-C18 column (length 150 mm, internal diameter 3.0 mm, particle size 3.5 μm; Agilent). Eluent A consisted of 0.2% formic acid in water. Eluent B consisted of 0.2% formic acid in acetonitrile. The histamine concentrations were determined in scheduled multiple reaction monitoring (sMRM). For quantification, a 7-point calibration was performed (0–2 μM) using pure histamine. Data analysis was done using Analyst 1.5.1® software (AB Sciex).

## 2.8 | Immunohistochemistry

Our method for immunohistochemically staining of cultured DRG neurons was previously described in detail.<sup>51</sup> Briefly after the experiments, the DRG neurons were fixed overnight at 4°C in 4% phosphate-buffered formaldehyde containing 0.2% picric acid, washed (3 × 10 min) with phosphate buffer, and permeabilized for 1 h at room temperature in phosphate-buffered saline (PBS) containing 0.5% Triton X-100, 0.1% NaN<sub>3</sub>, and 4% horse serum. After further washing (3 × 10 min with phosphate buffer), tissues were incubated with the primary antibody sheep anti-PGP 9.5 (1:20,000;

The Binding Site) for selectively labeling enteric neurons. This was followed by incubation with the secondary antibody Cy5 (1:500; Dianova). The fluorescence was examined by an epifluorescence microscope (BX61W1; Olympus).

## 2.9 | Data analysis and statistics

As described previously, the [Ca<sup>2+</sup>]<sub>i</sub> imaging technique enables the detection of cell outlines during the rise of intracellular [Ca<sup>2+</sup>]<sub>i</sub> accompanied by the increase in Fluo-4AM fluorescence.<sup>44</sup> In the DRG cell culture, the neurons were identified by their response to nicotine (Sigma; 100 μM, 600 ms pressure pulse application)<sup>42</sup> and/or by their PGP 9.5 immunoreactivity. As parameter for cell activation the maximum intracellular [Ca<sup>2+</sup>]<sub>i</sub> increase relative to resting light level (%ΔF/F) was determined for each stimulus. Further, the time interval (s) from beginning of a stimulus to the peak [Ca<sup>2+</sup>]<sub>i</sub> responses and the relative proportion of responding neurons in the field of view using a ×20 objective (0.08 mm<sup>2</sup>) was determined taking the number of those cells responding to ionomycin as 100%.

For the [Ca<sup>2+</sup>]<sub>i</sub> imaging experiments in submucous plexus preparations, we used EFS<sub>TM</sub> to identify the neuronal area since only neurons respond directly to electrical stimulation.<sup>19,44,52</sup> As a parameter for evoked activation, the percentage of responding neuronal area per ganglion (taking the neuronal area responding to EFS<sub>TM</sub> as 100%) as well as the amplitude of the [Ca<sup>2+</sup>]<sub>i</sub> peak per ganglion and the time to peak were calculated.

The analysis of the fast neuroimaging experiments was performed as described before in detail.<sup>44,53</sup> Briefly, individual neurons can be visualized since the voltage-sensitive dye (Di-8-ANEPPS) incorporates into the membrane revealing the outline of individual cell bodies. The overlay of signals with the ganglion image allowed us to analyze the response of individual neurons. As parameters for the substance-evoked activation of the submucous neurons, we calculated the percentages of responding neurons per ganglion (setting the number of neurons per ganglion which either responded to electrical stimulation or were spontaneously active as 100%) together with their average spike discharge frequency [Hz] per ganglion.

Numbers of tissues (equivalent to number of patients or guinea-pigs), ganglia, and neurons are given in sequence without further specification, for example, a result based on experiments from four tissues, five ganglia, and 20 neurons is presented as (4/5/20). Data are expressed as median values with the 25th and 75th quartiles given in brackets. Statistical analysis and graphs were performed using SigmaPlot 12.5 (Systat Software Inc.). Data were tested for normal distribution using the Shapiro-Wilk test. To determine whether an effect produced by a substance was statistically significantly different from zero, a one-sample signed rank was performed. Data derived from the [Ca<sup>2+</sup>]<sub>i</sub> imaging experiments were compared using the Mann-Whitney rank sum test. Data derived from the voltage-sensitive dye imaging experiments in the human submucous plexus were tested with the Kruskal-Wallis one-way analysis of variance on ranks (comparing neuronal

effects evoked by different pollen species) or by the Friedman repeated measures analysis of variance on ranks combined with the Student–Newman–Keuls Method as pairwise multiple comparison procedure, (comparing results from pharmacology experiments, which were obtained in a paired experimental design). For all tests, a  $p$  value of  $<0.05$  was considered significant. Data availability statement: All data  $n \leq 30$  are presented and available in the manuscript.

### 3 | RESULTS

#### 3.1 | Low MW aqueous birch pollen extracts activate cultured guinea-pig dorsal root ganglia

In a first set of experiments, we used cultured guinea-pig DRG's and  $Ca^{2+}$  imaging to screen for neuroactive properties of the low MW aqueous pollen fractions ( $APE_{<3kDa}$ ), first focusing on the birch pollen (Figure 1A). Results were based on DRGs from two guinea-pigs, three primary cultures, 10 fields of view containing clusters of in total 35 DGR neurons (guinea-pig no.1/one culture/five fields of view/19 DRG neurons; guinea-pig no. 2/two cultures/five fields of view/16 DRG neurons). The size of the DRG neurons ranged from 14 to 34  $\mu m$ . According to Lawson et al.<sup>54</sup> DRGs were divided into small ( $<25 \mu m$ ) and medium/large cells ( $25-35/>35 \mu m$ ). Thirty-one DRG neurons responded to the application of birch  $APE_{<3kDa}$  and showed a strong  $[Ca^{+2}]_i$  transit of 77.5 % $\Delta F/F$  (48.7/128.7) which was significantly different from zero ( $p < 0.001$ ); one-sample signed rank test; 81% ( $n=25$ ) belonging to the small type and 19% ( $n=6$ ) to the medium/large type. However, both types responded equally [DRG neurons  $<25 \mu m$ : 79.1% $\Delta F/F$  (42.7/133.3) vs. DRG neurons  $\geq 35 \mu m$ : 51.0% $\Delta F/F$  (49.2/101.6;  $p=0.382$ , Mann–Whitney rank sum test)]. The signal from DRG neurons peaked shortly after stimulus onset [6.0s (4.0/10.3),  $n=31$ ]. In six small-sized DRG neurons, we tested the involvement of TTX sensitive sodium channels. Neurons still responded to birch  $APE_{<3kDa}$  in TTX (0.5  $\mu M$ ), but the  $[Ca^{+2}]_i$  peak was greatly reduced from 59.8% $\Delta F/F$  (37.8/117.2) to 2.9 (0.0/6.3), thus indicating additional TTX insensitive mechanisms underlying the  $Ca^{2+}$  signaling in response to birch pollen extracts.

#### 3.2 | Aqueous birch pollen extracts activate neurons in freshly dissected submucous plexus preparations

Using  $Ca^{2+}$  imaging technique, we checked whether the same sample would also activate neurons in intact tissue, that is, in fresh, whole mount preparations of the enteric submucous plexus of guinea-pig intestine and also in human intestinal preparations. As shown in Figure 1B, birch  $APE_{<3kDa}$  evoked a strong  $[Ca^{+2}]_i$  transient in all neurons of guinea-pig submucous ganglia. In this approach, we additionally tested for effects of the corresponding high MW fraction ( $APE_{\geq 3kDa}$ ) containing the allergen fraction. Although the  $APE_{\geq 3kDa}$



fraction also activated a majority of neurons per ganglion [100.0% (83.3/100.0);  $n=4$  tissues and 4 ganglia], the peak response was clearly lower compared to the effects of  $APE_{<3kDa}$  (Figure 1B). The time to peak, however, did not differ [ $APE_{<3kDa}$ : 28.7s (25.2/32.3) vs.  $APE_{\geq 3kDa}$ : 30.0s (28.5/32.8),  $p=0.648$ , Mann–Whitney rank sum test]. Also, in human submucous plexus preparations ( $n=8$  tissues, 11 ganglia) the low MW birch pollen fraction evoked a clear  $[Ca^{+2}]_i$  signal [46.7% $\Delta F/F$  (31.5/66.0)] in almost all vital neurons per ganglion [100.0% (40.8/100.0) peaking at 21.4s (16.0/27.5)]. All  $[Ca^{+2}]_i$  signals elicited by pollen extracts were significantly different from zero ( $p < 0.02$ ; one-sample signed rank test).

#### 3.3 | Low MW ragweed, birch, and hazel pollen extracts induce fast neuronal responses

To check for fast onset responses, we continued our experiments using an ultra-fast neuroimaging approach with a voltage-sensitive dye as a reporter. Here, we focused exclusively on human submucous plexus preparations and included the low MW fractions of two further pollen taxa (ragweed and hazel). The spritz application of  $APE_{<3kDa}$  of all three pollen types evoked a fast onset action potential discharge in human submucous neurons (Figure 2). The spike discharge was comparable between the three pollen extracts [median for birch: 7.0Hz (6.2/9.8), ragweed: 5.7Hz (4.4/7.1), and hazel  $APE_{<3kDa}$  8.4Hz (4.3/12.5), respectively]; all effects were significantly different from zero,  $p < 0.016$ ; one-sample signed rank test. Likewise, the percentage of neurons responding per ganglion to the extracts were similar [median birch: 40.0% (33.3/100.0), ragweed: 50.8% (34.4/85.6), and hazel:  $APE_{<3kDa}$  83.3% (57.1/100.0), respectively; Figure 2]. Next, we investigated which substance might be involved in the nerve activation. We focused on histamine as the most likely candidate. As seen in Figure 3A,B, the spike discharge of those neurons that responded to birch and ragweed  $APE_{<3kDa}$  was significantly reduced by a mixture of H1, H2, and H3 receptor antagonists and recovered after wash out. The response to corresponding hazel  $APE_{<3kDa}$  was only marginally affected by histamine receptor antagonists.

#### 3.4 | Aqueous ragweed, birch, and hazel pollen extracts contain histamine

Using targeted liquid chromatography-mass spectrometry, we identified histamine in the extracts of the different pollen species (Table 1). The analysis revealed a higher histamine concentration for ragweed  $APE$  in all fractions compared to birch and hazel  $APE$ , respectively. In  $APE_{<3kDa}$ , the histamine concentration was highest for ragweed (0.764  $\mu M$ ), lowest for hazel pollen (0.013  $\mu M$ ). Regarding the histamine concentration in the non-fractionated total extracts, it was found that the deviations between the measured values and the calculated data (concentrations in the  $APE_{<3kDa}$  plus  $APE_{\geq 3kDa}$ ) were small and ranged from 5.6 to 13.4%. This underlines the reliability of the quantitative analysis.

**(A) Guinea pig DRG culture: Response to birch APE<sub><3kDa</sub> fraction****(B) Guinea pig submucous neurons: Response to birch APE<sub><3kDa</sub> and APE<sub>≥3kDa</sub> fractions**

**FIGURE 1** (A) Effects of birch low MW aqueous pollen fractions (APE<sub><3kDa</sub>) on cultured guinea-pig DRG neurons. Identification of anti-PGP 9.5 positive neurons (red) in the cell culture (left image; epifluorescence microscope Olympus BX61W1; objective  $\times 40$ ). Calcium image of the cell culture stained with Ca<sup>2+</sup> sensitive dye Fluo-4 AM brightfield image in the middle, fluorescence image on the right panel [epifluorescence microscope Zeiss Axio Observer A1; objective  $\times 20$ ; DRG neuron (blue arrow)]. Cell activation was reported by imaging changes in intracellular calcium [Ca<sup>2+</sup>]<sub>i</sub>. The graph (right panel) illustrates the traces of the [Ca<sup>2+</sup>]<sub>i</sub> signals in response to the APE<sub><3kDa</sub> fraction of birch pollen. The signal of the DRG neuron appeared instantaneously after the onset of the stimulus. The sample frequency in this example was 6.2 Hz. (B) Effects of low (APE<sub><3kDa</sub>) and high (APE<sub>≥3kDa</sub>) MW birch pollen fractions on guinea-pig submucous neurons. The traces on the left panel illustrate that the birch APE<sub><3kDa</sub> fraction, containing no protein allergens, evoked a stronger neuronal activation compared to the corresponding APE<sub>≥3kDa</sub> fraction. The amplitudes of the [Ca<sup>2+</sup>]<sub>i</sub> signals (right panel) differed significantly. The bars below the traces indicate the duration of the stimulus. Statistics were based on the number of ganglia. *p*-values are based on Mann-Whitney rank sum test; numbers in parentheses indicate number of tissues/ganglia studied.



**4 | DISCUSSION**

In the present study, we demonstrated for the first time that low MW aqueous pollen extracts of birch, ragweed, and hazel contain neuroactive components that strongly activate spinal sensory

neurons (in guinea-pig DRGs) and enteric neurons of the submucosal plexus of guinea-pig and human. Nerve activation was independent of classic allergic hypersensitivity reactions. Pharmacologically, a significant histaminergic component was shown to mediate the nerve-activating effects of birch and ragweed pollen extracts, but not for







(5/5/39)

(5/5/49)

(4/4/17)

Pollen type	Histamine concentration ( $\mu\text{M}$ )		
	Low molecular weight fraction ( $\text{APE}_{<3\text{kDa}}$ )	High molecular weight fraction ( $\text{APE}_{\geq 3\text{kDa}}$ )	Total APE
Ragweed	0.764	0.168	1.076
Birch	0.047	0.004	0.055
Hazel	0.013	0.004	0.018

TABLE 1 Histamine concentration in different fractions of the aqueous pollen extracts (APE).

signaling between nerves and immune cells, most importantly mast cells.<sup>19,58–60</sup> However, as we have shown in the present study for different pollen types, there appears to be an effective mechanism that ensures the activation of neurons independent of the allergic response of the host. Here, neuronal stimulation is mediated via low MW substances released directly from pollen grains.

To look for neuronal activation, we first performed calcium imaging experiments. This is the method of choice to visualize slow, long-lasting changes in the intracellular calcium  $[\text{Ca}^{2+}]_i$  levels and to also detect low-threshold activation.<sup>44</sup> The technique is considered as a very sensitive method minimizing the risk to miss responses or effects. In our study, we primarily focused on extracts from birch pollen, as birch pollen is the most potent aeroallergen in Central and Northern Europe with high seasonal peak concentrations,<sup>61</sup> sensitization rates of 6.4–22.6%, and considerable clinical importance.<sup>62</sup> Birch pollen, in particular, is frequently discussed in the context of the pathophysiology of eosinophilic esophagitis,<sup>1,14,16</sup> (for review),<sup>7,8,63</sup> We have studied two different types of neurons. One is primary cultured guinea-pig DRG neurons in order to study the ability of pollen-derived substances to affect sensory nerves. The DRGs were taken from thoracic segments T2–L2 because these ganglia represent the major splanchnic afferent innervation of the bronchi, lungs, and intestines in rodents.<sup>43</sup> DRGs contain the cell bodies of afferent sensory neurons. The receptors found on peripheral terminals of these neurons *in vivo* are most often expressed on their cell bodies as well.<sup>64</sup> As we could show in the present study, the  $\text{APE}_{<3\text{kDa}}$  of birch pollen strongly activate DRG sensory neurons. About 80% of the responding neurons belonged to the small size population, which mainly mediate pain-, itch-, and somato-sensation (for review).<sup>56,57</sup> We have previously reported a very similar response of small-sized DRG neurons after application of mucosal biopsy supernatants of patients with irritable bowel syndrome.<sup>45</sup> The evoked calcium peaks were strongly reduced by preincubation of the neurons with TTX, suggesting that opening of voltage-sensitive fast Na-channels are required for initiating intracellular Ca-signals.

Our study showed that similarly to the DRG neurons, guinea-pig and human enteric neurons responded with strong calcium signals to a stimulation with the  $\text{APE}_{<3\text{kDa}}$  of birch pollen. Therefore, nerve activation is a rather general hallmark of the pollen extracts and may not reflect the specificity of a particular neuron type or species. Guinea-pig intestinal plexus preparations were also used to test the effect of the  $\text{APE}_{\geq 3\text{kDa}}$  fraction of birch pollen extracts, which, unlike the  $\text{APE}_{<3\text{kDa}}$  fraction, contain the allergenic proteins. The  $\text{APE}_{\geq 3\text{kDa}}$

fraction also evoked a calcium signal in guinea-pig neurons, but the response was much smaller compared with the effects of the corresponding  $\text{APE}_{<3\text{kDa}}$  birch pollen fraction. A hypersensitivity reaction can be excluded because the guinea-pigs were not sensitized to the pollen, and, moreover, the neuronal response occurred much more rapidly than would be expected from a classical mast cell-triggered event.<sup>19</sup> Thus, either direct effects of proteins or other, as yet unidentified, substances need to be considered. However, some residual low MW substances that may be retained in the high MW fraction, for example, due to binding to filter matrix or protein, could also be responsible for the corresponding neuronal activation. We will come back to this point in detail later in the discussion.

Because of species differences in terms of receptor pharmacology between rodents and humans (for review),<sup>58</sup> we focused exclusively on human ENS preparations as matrix for all further targeted studies on signaling and pharmacology of pollen-induced nerve activation, despite the limited availability of human tissues. We also switched our recording to fast neuroimaging technology. By using a voltage-sensitive dye, this technique allows reliable measurement of action potentials at the level of single cells.<sup>49</sup> Furthermore, we wanted to clarify whether the nerve-activating effect of the low molecular aqueous pollen extracts is specific to birch, or represents a general property of other pollen species as well. To clarify this, we included two more pollen taxa. One of them was hazel, characterized by a marked homology of the major allergen (Cor a 1) to that of birch (Bet v 1),<sup>61,65</sup> and ragweed pollen as a representative of herbaceous plants. Ragweed is in the focus of many studies due to its extreme allergenic potential.<sup>66</sup> The  $\text{APE}_{<3\text{kDa}}$  from all three pollen types elicited an immediate discharge of action potentials with broadly similar frequency and pattern, suggesting that nerve-activating effects may not be a species-specific but a general feature of the water-soluble, non-allergenic fraction of pollen extracts. We identified histamine as an important mediator for two of the pollen extracts based on the finding that the nerve activation was strongly reduced by a combination of H1, H2, and H3 receptor blockers, which primarily mediated histamine effects in human enteric neurons.<sup>26</sup> The histaminergic component was only observed for ragweed and birch pollen extracts. Other, yet to be identified, substances must mediate the effects of hazel pollen extract.

Further, we quantified the histamine concentration in the pollen extracts using LC-MS/MS analysis. A high level was detected in the  $\text{APE}_{<3\text{kDa}}$  fraction of ragweed pollen, which is very consistent with the histamine-mediated nerve activation of this extract. The corresponding concentration in the birch pollen extract was lower,

however, still about 4x higher than in the APE<sub><3kDa</sub> extracts from hazel pollen. This could explain the lack of effects of histamine antagonists on nerve activation in hazel APE<sub><3kDa</sub>. The quantitative analysis confirmed our pharmacological findings and highlights the importance of histamine as a neuroactive low molecular weight component of pollen.

We extended the analyses also to the high molecular weight fractions. It turned out that only for ragweed pollen a significant concentration of histamine could be detected. The level suggests that a pharmacological proof of histamine would also work. This means that in the case of ragweed pollen, a histaminergic neuroactive component may also be involved for the APE<sub>≥3kDa</sub> fraction. In contrast, in the APE<sub>≥3kDa</sub> fraction of birch (and equally of hazel) pollen, the level of histamine was so low making a pharmacological detection of histamine by antagonists in our setup rather unlikely. Therefore, we believe that mainly non-histaminergic mechanisms are responsible for the observed nerve-activating effect of the birch pollen APE<sub>≥3kDa</sub> fraction. In addition, we measured the histamine concentration also in the total extract (not fractionated). The values are comparable in magnitude to the histamine concentrations determined in the study by Williams et al.<sup>25</sup> in aqueous and phenolic pollen extracts of a variety different species. Importantly, the dose of pollen extract used in our experiments is within the dose range applied in nasal allergen challenges (e.g., for birch: 1–40 µg/mL Bet v)<sup>67,68</sup> and can thus be considered clinically relevant.

In the present study, we provided evidence that aqueous pollen extracts of birch, ragweed, and hazel contain neuroactive components which strongly activate spinal sensory neurons and enteric neurons. This nerve activation is independent of the classical allergic hypersensitivity responses. We suggest that the rapid release of bioactive mediators from pollen during contact with the aqueous phase of mucosal membranes may represent a pathophysiological relevant feature contributing to the manifestation or aggravation of acute or chronic inflammatory conditions of the mucosa.

## AUTHOR CONTRIBUTIONS

SB designed and drafted the work, conducted the experiments, analyzed and interpreted the data, and wrote the manuscript. SS conducted part of the Ca imaging experiments and analyzed the data. PG performed the targeted liquid chromatography tandem-mass spectrometry, analyzed and interpreted the data, and wrote the corresponding parts of the manuscript. FZ and IED provided human samples and medical council. They were accountable for correct patient characterization and material handling. MS and CT-H critically reviewed the manuscript for important intellectual content. They were involved in the conception, design, and funding of the work. SG designed and drafted the work, interpreted the data, and edited and revised the manuscript. She had the original idea and was involved in the conception and design of the work. All authors have read and approved the final version of this manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are

appropriately investigated and resolved. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed. This work has been previously presented as a Poster at the 3rd Meeting of the Federation of Neurogastroenterology and Motility 2018 in Amsterdam. Buehner S, Schäuffele S, Gilles-Stein S, Demir IE, Ceyhan G, Zeller F, Traidl-Hoffmann C, Schemann M. Pollen extracts of birch and ragweed activate human and guinea-pig submucous neurons and guinea-pig DRG neurons. P014.

## ACKNOWLEDGMENT

Open Access funding enabled and organized by Projekt DEAL.

## FUNDING INFORMATION

The study was supported by the Christine Kühne Research Center for Allergy and Education (CK-Care).

## CONFLICT OF INTEREST STATEMENT

No competing interests declared.

## ORCID

Sabine Buhner  <https://orcid.org/0000-0002-9115-0463>

Pieter Giesbertz  <https://orcid.org/0000-0001-7461-8602>

Ihsan Ekin Demir  <https://orcid.org/0000-0001-8241-1871>

Claudia Traidl-Hoffmann  <https://orcid.org/0000-0001-5085-5179>

Michael Schemann  <https://orcid.org/0000-0003-1007-9843>

Stefanie Gilles  <https://orcid.org/0000-0002-5159-2558>

## REFERENCES

- Magnusson J, Lin XP, Dahlman-Höglund A, et al. Seasonal intestinal inflammation in patients with birch pollen allergy. *J Allergy Clin Immunol*. 2003;112(1):45-50. doi:10.1067/mai.2003.1604
- Slae M, Persad R, Leung AJT, Gabr R, Brocks D, Huynh HQ. Role of environmental factors in the development of pediatric eosinophilic esophagitis. *Dig Dis Sci*. 2015;60(11):3364-3372. doi:10.1007/s10620-015-3740-7
- Fahey L, Robinson G, Weinberger K, Giambrone AE, Solomon AB. Correlation between aeroallergen levels and new diagnosis of eosinophilic esophagitis in New York City. *J Pediatr Gastroenterol Nutr*. 2017;64(1):22-25. doi:10.1097/MPG.0000000000001245
- Armentia A, Martín-Armentia S, Martín-Armentia B, et al. Is eosinophilic esophagitis an equivalent of pollen allergic asthma? Analysis of biopsies and therapy guided by component resolved diagnosis. *Allergol Immunopathol (Madr)*. 2018;46(2):181-189. doi:10.1016/j.aller.2017.11.001
- Anderson J, Moonie S, Hogan MB, Scherr R, Allenback G. Eosinophilic esophagitis: comorbidities and atopic disease in Nevada. *Dis Esophagus*. 2020;33(5):doz105. doi:10.1093/dote/doz105
- Agulló-García A, Cubero JL, Lezaun A, Boné J, Guallar I, Colás C. Clinical and anatomopathological features of eosinophilic oesophagitis in children and adults. *Allergol Immunopathol (Madr)*. 2020;48(6):560-567. doi:10.1016/j.aller.2020.03.009
- Ram G, Lee J, Ott M, et al. Seasonal exacerbation of esophageal eosinophilia in children with eosinophilic esophagitis and allergic rhinitis. *Ann Allergy Asthma Immunol*. 2015;115(3):224-228.e1. doi:10.1016/j.anai.2015.07.004

8. Capucilli P, Hill DA. Allergic comorbidity in eosinophilic esophagitis: mechanistic relevance and clinical implications. *Clin Rev Allergy Immunol*. 2019;57(1):111-127. doi:10.1007/s12016-019-08733-0
9. Johnson E, Therkelsen SP, Nentwich I, Nissen-Meyer LSH, Hetland G. IgE-sensitization to food and inhalant allergens in IBD patients compared with normal blood donors at Oslo university hospital, Norway. *Scand J Gastroenterol*. 2019;54(9):1107-1110. doi:10.1080/00365521.2019.1663445
10. Tobin MC, Moparty B, Farhadi A, DeMeo MT, Bansal PJ, Keshavarzian A. Atopic irritable bowel syndrome: a novel subgroup of irritable bowel syndrome with allergic manifestations. *Ann Allergy Asthma Immunol*. 2008;100(1):49-53. doi:10.1016/S1081-1206(10)60404-8
11. Koloski N, Jones M, Walker MM, et al. Population based study: atopy and autoimmune diseases are associated with functional dyspepsia and irritable bowel syndrome, independent of psychological distress. *Aliment Pharmacol Ther*. 2019;49(5):546-555. doi:10.1111/apt.15120
12. Patel K, Vila-Nadal G, Shah J, et al. Is pollen-food syndrome a frequent comorbidity in adults with irritable bowel syndrome? *Allergy*. 2020;75(7):1780-1783. doi:10.1111/all.14209
13. Nybacka S, Öhman L, Störstrud S, et al. Neither self-reported atopy nor IgE-mediated allergy are linked to gastrointestinal symptoms in patients with irritable bowel syndrome. *Neurogastroenterol Motil*. 2018;30(10):e13379. doi:10.1111/nmo.13379
14. van Rhijn BD, van Ree R, Versteeg SA, et al. Birch pollen sensitization with cross-reactivity to food allergens predominates in adults with eosinophilic esophagitis. *Allergy*. 2013;68(11):1475-1481. doi:10.1111/all.12257
15. Rentzos G, Johanson L, Goksör E, Telemo E, Lundbäck B, Ekerljung L. Prevalence of food hypersensitivity in relation to IgE sensitisation to common food allergens among the general adult population in West Sweden. *Clin Transl Allergy*. 2019;9:22. doi:10.1186/s13601-019-0261-z
16. Rentzos G, Lundberg V, Stotzer PO, Pullerits T, Telemo E. Intestinal allergic inflammation in birch pollen allergic patients in relation to pollen season, IgE sensitization profile and gastrointestinal symptoms. *Clin Transl Allergy*. 2014;4:19. doi:10.1186/2045-7022-4-19
17. Armentia A, Martín-Armentia S, Álvarez-Nogal R, Armentia BM, Gayoso MJ, Fernández-González D. Germination of pollen grains in the oesophagus of individuals with eosinophilic oesophagitis. *Clin Exp Allergy*. 2019;49(4):471-473. doi:10.1111/cea.13312
18. Wilson AF, Novey HS, Berke RA, Surprenant EL. Deposition of inhaled pollen and pollen extract in human airways. *N Engl J Med*. 1973;288(20):1056-1058. doi:10.1056/NEJM197305172882006
19. Buhner S, Barki N, Greiter W, et al. Calcium imaging of nerve-mast cell signaling in the human intestine. *Front Physiol*. 2017;8:971. doi:10.3389/fphys.2017.00971
20. Gilles S, Behrendt H, Ring J, Traidl-Hoffmann C. The pollen enigma: modulation of the allergic immune response by non-allergenic, pollen-derived compounds. *Curr Pharm des*. 2012;18(16):2314-2319.
21. Pointner L, Bethanis A, Thaler M, et al. Initiating pollen sensitization – complex source, complex mechanisms. *Clin Transl Allergy*. 2020;10:36. doi:10.1186/s13601-020-00341-y
22. Gilles-Stein S, Beck I, Chaker A, et al. Pollen derived low molecular compounds enhance the human allergen specific immune response in vivo. *Clin Exp Allergy*. 2016;46(10):1355-1365. doi:10.1111/cea.12739
23. Guerin B, Tioulong S. Analysis of the non-immunological activity of allergen extracts in cutaneous tests. *Clin Allergy*. 1979;9(3):283-291. doi:10.1111/j.1365-2222.1979.tb01555.x
24. Marquardt P, Vogg G. Pharmacological and chemical investigations of the active substances of bee pollens. *Arzneimittelforschung*. 1952;2(6):267-271.
25. Williams PB, Nolte H, Dolen WK, Koepke JW, Selner JC. The histamine content of allergen extracts. *J Allergy Clin Immunol*. 1992;89(3):738-745. doi:10.1016/0091-6749(92)90382-c
26. Breunig E, Michel K, Zeller F, Seidl S, Weyhern CW, Schemann M. Histamine excites neurones in the human submucous plexus through activation of H1, H2, H3 and H4 receptors. *J Physiol (Lond)*. 2007;583(Pt 2):731-742. doi:10.1113/jphysiol.2007.139352
27. Wimmer M, Alessandrini F, Gilles S, et al. Pollen-derived adenosine is a necessary cofactor for ragweed allergy. *Allergy*. 2015;70(8):944-954. doi:10.1111/all.12642
28. Raybould HE, Cooke HJ, Christofi FL. Sensory mechanisms: transmitters, modulators and reflexes. *Neurogastroenterol Motil*. 2004;16(Suppl 1):60-63. doi:10.1111/j.1743-3150.2004.00477.x
29. Traidl-Hoffmann C, Mariani V, Hochrein H, et al. Pollen-associated phytoprostanes inhibit dendritic cell interleukin-12 production and augment T helper type 2 cell polarization. *J Exp Med*. 2005;201(4):627-636. doi:10.1084/jem.20041065
30. Gilles S, Mariani V, Bryce M, et al. Pollen-derived E1-phytoprostanes signal via PPAR-gamma and NF-kappaB-dependent mechanisms. *J Immunol*. 2009;182(11):6653-6658. doi:10.4049/jimmunol.0802613
31. Gilles S, Beck I, Lange S, Ring J, Behrendt H, Traidl-Hoffmann C. Non-allergenic factors from pollen modulate T helper cell instructing notch ligands on dendritic cells. *World Allergy Organ J*. 2015;8(1):2. doi:10.1186/s40413-014-0054-8
32. Soh WT, Aglas L, Mueller GA, et al. Multiple roles of bet v 1 ligands in allergen stabilization and modulation of endosomal protease activity. *Allergy*. 2019;74(12):2382-2393. doi:10.1111/all.13948
33. Huby RD, Dearman RJ, Kimber I. Why are some proteins allergens? *Toxicol Sci*. 2000;55(2):235-246. doi:10.1093/toxsci/55.2.235
34. Breiteneder H, Ferreira F, Hoffmann-Sommergruber K, et al. Four recombinant isoforms of Cor a I, the major allergen of hazel pollen, show different IgE-binding properties. *Eur J Biochem*. 1993;212(2):355-362. doi:10.1111/j.1432-1033.1993.tb17669.x
35. Mogensen JE, Wimmer R, Larsen JN, Spangfort MD, Otzen DE. The major birch allergen, bet v 1, shows affinity for a broad spectrum of physiological ligands. *J Biol Chem*. 2002;277(26):23684-23692. doi:10.1074/jbc.M202065200
36. Léonard R, Wopfner N, Pabst M, et al. A new allergen from ragweed (*Ambrosia artemisiifolia*) with homology to art v 1 from mugwort. *J Biol Chem*. 2010;285(35):27192-27200. doi:10.1074/jbc.M110.127118
37. Blume C, Swindle EJ, Gilles S, Traidl-Hoffmann C, Davies DE. Low molecular weight components of pollen alter bronchial epithelial barrier functions. *Tissue Barriers*. 2015;3(3):e1062316. doi:10.1080/15476286.2015.1062316
38. Bergougnan C, Dittlein DC, Hümmel E, et al. Physical and immunological barrier of human primary nasal epithelial cells from non-allergic and allergic donors. *World Allergy Organ J*. 2020;13(3):100109. doi:10.1016/j.waojou.2020.100109
39. Gilles S, Blume C, Wimmer M, et al. Pollen exposure weakens innate defense against respiratory viruses. *Allergy*. 2020;75(3):576-587. doi:10.1111/all.14047
40. Gilles S, Fekete A, Zhang X, et al. Pollen metabolome analysis reveals adenosine as a major regulator of dendritic cell-primed T(H) cell responses. *J Allergy Clin Immunol*. 2011;127(2):454-461.e1-9. doi:10.1016/j.jaci.2010.12.1082
41. Grundy D. Principles and standards for reporting animal experiments in the journal of physiology and experimental physiology. *Exp Physiol*. 2015;100(7):755-758. doi:10.1113/EP085299
42. Buhner S, Li Q, Vignali S, et al. Activation of human enteric neurons by supernatants of colonic biopsy specimens from patients with irritable bowel syndrome. *Gastroenterology*. 2009;137(4):1425-1434. doi:10.1053/j.gastro.2009.07.005
43. Robinson DR, McNaughton PA, Evans ML, Hicks GA. Characterization of the primary spinal afferent innervation of the mouse colon using retrograde labelling. *Neurogastroenterol Motil*. 2004;16(1):113-124. doi:10.1046/j.1365-2982.2003.00456.x

44. Michel K, Michaelis M, Mazzuoli G, Mueller K, Vanden Berghe P, Schemann M. Fast calcium and voltage-sensitive dye imaging in enteric neurones reveal calcium peaks associated with single action potential discharge. *J Physiol (Lond)*. 2011;589(Pt 24):5941-5947. doi:10.1113/jphysiol.2011.219550
45. Buhner S, Braak B, Li Q, et al. Neuronal activation by mucosal biopsy supernatants from irritable bowel syndrome patients is linked to visceral sensitivity. *Exp Physiol*. 2014;99:1299-1311. doi:10.1113/expphysiol.2014.080036
46. Nemethova A, Michel K, Gomez-Pinilla PJ, Boeckxstaens GE, Schemann M. Nicotine attenuates activation of tissue resident macrophages in the mouse stomach through the  $\beta$ 2 nicotinic acetylcholine receptor. *PLoS One*. 2013;8(11):e79264. doi:10.1371/journal.pone.0079264
47. Neunlist M, Peters S, Schemann M. Multisite optical recording of excitability in the enteric nervous system. *Neurogastroenterol Motil*. 1999;11(5):393-402.
48. Buhner S, Li Q, Berger T, et al. Submucous rather than myenteric neurons are activated by mucosal biopsy supernatants from irritable bowel syndrome patients. *Neurogastroenterol Motil*. 2012;24(12):1134-e572. doi:10.1111/nmo.12011
49. Schemann M, Michel K, Peters S, Bischoff SC, Neunlist M. Cutting-edge technology. III. Imaging and the gastrointestinal tract: mapping the human enteric nervous system. *Am J Physiol Gastrointest Liver Physiol*. 2002;282(6):G919-G925. doi:10.1152/ajpgi.00043.2002
50. Rauer D, Gilles S, Wimmer M, et al. Ragweed plants grown under elevated CO<sub>2</sub> levels produce pollen which elicit stronger allergic lung inflammation. *Allergy*. 2021;76(6):1718-1730. doi:10.1111/all.14618
51. Schemann M, Kugler EM, Buhner S, et al. The mast cell degranulator compound 48/80 directly activates neurons. *PLoS One*. 2012;7(12):e52104. doi:10.1371/journal.pone.0052104
52. Ostertag D, Buhner S, Michel K, et al. Reduced responses of submucous neurons from irritable bowel syndrome patients to a cocktail containing histamine, serotonin, TNF $\alpha$ , and tryptase (IBS-cocktail). *Front Neurosci*. 2015;9:465. doi:10.3389/fnins.2015.00465
53. Buhner S, Hahne H, Hartwig K, et al. Protease signaling through protease activated receptor 1 mediate nerve activation by mucosal supernatants from irritable bowel syndrome but not from ulcerative colitis patients. *PLoS One*. 2018;13(3):e0193943. doi:10.1371/journal.pone.0193943
54. Lawson SN, Crepps BA, Perl ER. Relationship of substance P to afferent characteristics of dorsal root ganglion neurones in Guinea-pig. *J Physiol*. 1997;505(Pt 1):177-191. doi:10.1111/j.1469-7793.1997.00177.x
55. van Wanrooij SJM, Wouters MM, Van Oudenhove L, et al. Sensitivity testing in irritable bowel syndrome with rectal capsaicin stimulations: role of TRPV1 upregulation and sensitization in visceral hypersensitivity? *Am J Gastroenterol*. 2014;109(1):99-109. doi:10.1038/ajg.2013.371
56. Udem BJ, Taylor-Clark T. Mechanisms underlying the neuronal-based symptoms of allergy. *J Allergy Clin Immunol*. 2014;133(6):1521-1534. doi:10.1016/j.jaci.2013.11.027
57. Voisin T, Bouvier A, Chiu IM. Neuro-immune interactions in allergic diseases: novel targets for therapeutics. *Int Immunol*. 2017;29(6):247-261. doi:10.1093/intimm/dxx040
58. Buhner S, Schemann M. Mast cell-nerve axis with a focus on the human gut. *Biochim Biophys Acta*. 2012;1822(1):85-92. doi:10.1016/j.bbdis.2011.06.004
59. Schemann M, Neunlist M. The human enteric nervous system. *Neurogastroenterol Motil*. 2004;16(Suppl 1):55-59. doi:10.1111/j.1743-3150.2004.00476.x
60. Tikoo S, Barki N, Jain R, et al. Imaging of mast cells. *Immunol Rev*. 2018;282(1):58-72. doi:10.1111/imr.12631
61. Biedermann T, Winther L, Till SJ, Panzner P, Knulst A, Valovirta E. Birch pollen allergy in Europe. *Allergy*. 2019;74(7):1237-1248. doi:10.1111/all.13758
62. Ziemianin M, Waga J, Czarnobilska E, Myszkowska D. Changes in qualitative and quantitative traits of birch (*Betula pendula*) pollen allergenic proteins in relation to the pollution contamination. *Environ Sci Pollut Res Int*. 2021;28(29):39952-39965. doi:10.1007/s11356-021-13483-8
63. Pineton de Chambrun G, Desreumaux P, Cortot A. Eosinophilic enteritis. *Dig Dis*. 2015;33(2):183-189. doi:10.1159/000369540
64. Drew LJ, Wood JN, Cesare P. Distinct mechanosensitive properties of capsaicin-sensitive and -insensitive sensory neurons. *J Neurosci*. 2002;22(12):RC228.
65. Darnhofer B, Tomin T, Liesinger L, Schittmayer M, Tomazic PV, Birner-Gruenberger R. Comparative proteomics of common allergenic tree pollens of birch, alder, and hazel. *Allergy*. 2021;76(6):1743-1753. doi:10.1111/all.14694
66. Chen KW, Marusciac L, Tamas PT, Valenta R, Panaitescu C. Ragweed pollen allergy: burden, characteristics, and management of an imported allergen source in Europe. *Int Arch Allergy Immunol*. 2018;176(3-4):163-180. doi:10.1159/000487997
67. Godnic-Cvar J, Susani M, Breiteneder H, et al. Recombinant bet v 1, the major birch pollen allergen, induces hypersensitivity reactions equal to those induced by natural bet v 1 in the airways of patients allergic to tree pollen. *J Allergy Clin Immunol*. 1997;99(3):354-359. doi:10.1016/s0091-6749(97)70053-8
68. Niederberger V, Stübner P, Spitzauer S, et al. Skin test results but not serology reflect immediate type respiratory sensitivity: a study performed with recombinant allergen molecules. *J Invest Dermatol*. 2001;117(4):848-851. doi:10.1046/j.0022-202x.2001.01470.x

**How to cite this article:** Buhner S, Schäuffele S, Giesbertz P, et al. Allergen-free extracts from birch, ragweed, and hazel pollen activate human and guinea-pig submucous and spinal sensory neurons. *Neurogastroenterology & Motility*. 2023;35:e14559. doi:10.1111/nmo.14559