

# Multiple Genome Sequences of Important Beer-Spoiling Lactic Acid Bacteria

Andreas J. Geissler,<sup>a</sup> Jürgen Behr,<sup>b</sup> Rudi F. Vogel<sup>a</sup>

Technische Universität München, Lehrstuhl für Technische Mikrobiologie, Freising, Germany<sup>a</sup>; Technische Universität München, Bavarian Center for Biomolecular Mass Spectrometry, Freising, Germany<sup>b</sup>

**Seven strains of important beer-spoiling lactic acid bacteria were sequenced using single-molecule real-time sequencing. Complete genomes were obtained for strains of *Lactobacillus paracollinoides*, *Lactobacillus lindneri*, and *Pediococcus clausenii*. The analysis of these genomes emphasizes the role of plasmids as the genomic foundation of beer-spoiling ability.**

Received 9 August 2016 Accepted 17 August 2016 Published 6 October 2016

Citation Geissler AJ, Behr J, Vogel RF. 2016. Multiple genome sequences of important beer-spoiling lactic acid bacteria. *Genome Announc* 4(5):e01077-16. doi:10.1128/genomeA.01077-16.

Copyright © 2016 Geissler et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jürgen Behr, [juergen.behr@wzw.tum.de](mailto:juergen.behr@wzw.tum.de).

Beer is a harsh environment with which bacteria have to cope, as antibacterial hurdles, such as the low pH and the presence of antibacterial hops, have to be taken. Some lactic acid bacteria (LAB) are capable of growing in and spoiling beer (1, 2). To gain insights into the genomic adaptation of LAB to beer, we sequenced the complete genomes of seven different brewery isolates, comprising three different species.

Beer spoilage ability was tested as described previously (3). DNA isolation, sequencing, assembly, annotation, and genome analysis were done as described previously for *Lactobacillus backii* (4). High-molecular-weight DNA was purified from de Man-Rogosa-Sharpe (MRS) liquid cultures using the Genomic-tip 100/G kit (Qiagen). Single-molecule real-time sequencing (SMRT) (PacBio RS II) was carried out at GATC Biotech (Konstanz, Germany) (5). An insert size of 8 to 12 kb was selected for library creation, resulting in at least 200 Mb of raw data from one to two SMRT cells (1 × 120-min movies) applying P4-C2 chemistry. Assembly was done with SMRT Analysis version 2.2.0.p2, using the Hierarchical Genome Assembly Process (HGAP) (6), and completed by manual curation (<https://github.com/PacificBiosciences/Bioinformatics-Training/wiki/Finishing-Bacterial-Genomes>). Genomes were annotated using the NCBI Prokaryotic Genome Annotation Pipe-

line (PGAP) and Rapid Annotations using Subsystems Technology (RAST) (7–9). Genomes were further analyzed with CMG-BioTools, BADGE, and PSORTb (3, 10, 11).

Strain characteristics, sequencing statistics, genome information, and accession numbers are listed in Table 1. Chromosome sizes range from 1.39 Mbp for *Lactobacillus lindneri* to 3.48 Mbp for *Lactobacillus paracollinoides*, with G+C contents ranging from 34.3 to 47.2%. We found zero to eight plasmids with G+C contents from 34.7 to 44.5% and sizes from 13,353 bp to 57,063 bp. The analysis of RAST-annotated genomes resulted in the following core genomes: *Lactobacillus paracollinoides* with 2,503 gene families, *L. lindneri* with 1,305 gene families, and *Pediococcus clausenii* with 1,687 gene families, while *P. clausenii* ATCC BAA-344 was also included in the latter calculation (12). Chromosomes encode four complete rRNA operons in case of *P. clausenii*, five in case of *L. paracollinoides*, and six for *L. lindneri*.

All seven genomes were compared to each other as well as to the genomes of 17 strains with relevance for beer spoiling, including one *P. clausenii*, five *L. backii*, five *P. damnosus*, and six *L. brevis* genomes (3, 4, 12–16). We found that the investigated LAB species with relevance for beer spoiling are characterized by different genomic preconditions regarding chromosome size, num-

TABLE 1 Strain characteristics, sequencing statistics, genome information, and accession numbers

Species	Strain	Source	Beer spoilage ability <sup>a</sup>	Biosample no. <sup>b</sup>	Accession no. <sup>c</sup>	Avg coverage of HGAP assembly (×)	Size (Mbp)	No. of contigs <sup>d</sup>	G+C content (%)	PEG <sup>e</sup>	CDSs <sup>f</sup>
<i>L. paracollinoides</i>	TMW 1.1979 (DSM 20197)	Beer	NB	SAMN04505735	CP014912–CP014914	107	3.32	3	47.0	2,953	2,872
<i>L. paracollinoides</i>	TMW 1.1994	Brewery environment	B	SAMN04505736	CP014915–CP014923	102	3.66	9	46.8	3,363	3,219
<i>L. paracollinoides</i>	TMW 1.1995	Pilsner beer	B	SAMN04505737	CP014924–CP014932	88	3.75	9	46.6	3,378	3,286
<i>L. lindneri</i>	TMW 1.481	Brewery environment	B	SAMN04505733	CP014907–CP014911	143	1.45	5	34.4	1,429	1,387
<i>L. lindneri</i>	TMW 1.1993	Beer	B	SAMN04505734	CP014872	262	1.39	1	34.3	1,347	1,305
<i>P. clausenii</i>	TMW 2.53	Brewery environment	NB-B	SAMN04505731	CP014933–CP014935	192	1.95	3	37.1	1,895	1,849
<i>P. clausenii</i>	TMW 2.54	Brewery environment	B	SAMN04505732	CP014936–CP014939	146	1.99	4	37.1	1,940	1,884

<sup>a</sup> NB, nonspoilage; B, spoiler; NB-B, unstable, mostly B.

<sup>b</sup> All BioSamples are part of BioProject PRJNA290141.

<sup>c</sup> Accession numbers are listed for all contigs of each whole genome (as range).

<sup>d</sup> In chromosome plus plasmids and partial plasmids.

<sup>e</sup> PEG, number of protein-encoding genes based on RAST annotation.

<sup>f</sup> CDSs, number of coding sequences (total) based on NCBI PGAP.

ber of proteins, G+C content, coding density, codon usage, amino acid usage, proteome similarity, chromosome (DNA) similarity, functional pattern (SEED/COG), and subcellular localization of proteins. Regarding chromosomal properties, beer-spoiling LAB species cover the whole diversity within the genus *Lactobacillus*. In contrast, brewery isolates of the abovementioned species, although to a different extent, share a number of highly homologous plasmid-carried genes, including the important lifestyle markers *horA*, *horC*, and *fabZ* (3, 17, 18). This emphasizes the role of plasmids in beer-spoiling ability (3, 19, 20).

**Accession number(s).** The seven complete genomes have been deposited in DDBJ/EMBL/GenBank under the accession numbers listed in Table 1.

## FUNDING INFORMATION

This work, including the efforts of Rudi F. Vogel, was funded by Allianz Industrie Forschung (AiF) (AiF 17576N).

This research project was funded by the German Ministry of Economics and Technology (via AiF) and the Wifoe (Wissenschaftsförderung der Deutschen Brauwirtschaft e.V., Berlin), project AiF 17576N. None of the funding sources had any influence on the study design, the collection, analysis, and interpretation of data, the writing of the report, or the decision to submit the article for publication.

## REFERENCES

- Vriesekoop F, Krahl M, Hucker B, Menz G. 2012. 125th anniversary review: bacteria in brewing: the good, the bad and the ugly. *J Inst Brew* 118:335–345. <http://dx.doi.org/10.1002/jib.49>.
- Suzuki K. 2011. 125th anniversary review: microbiological instability of beer caused by spoilage bacteria. *J Inst Brew* 117:131–155. <http://dx.doi.org/10.1002/j.2050-0416.2011.tb00454.x>.
- Behr J, Geissler AJ, Schmid J, Zehe A, Vogel RF. 2016. The identification of novel diagnostic marker genes for the detection of beer spoiling *Pediococcus damnosus* strains using the BLAST diagnostic gene finder. *PLoS One* 11:e0152747. <http://dx.doi.org/10.1371/journal.pone.0152747>.
- Geissler AJ, Behr J, Vogel RF. 2016. Multiple genome sequences of the important beer spoiling species *Lactobacillus backii*. *Genome Announc* 4. <http://dx.doi.org/10.1128/genomeA.00826-16>.
- Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <http://dx.doi.org/10.1126/science.1162986>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpidis N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of Standard Operating Procedures (SOPs) for (meta)genomic annotation. *Omic* 12:137–141. <http://dx.doi.org/10.1089/omi.2008.0017>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
- Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, Dao P, Sahinalp SC, Ester M, Foster LJ, Brinkman FS. 2010. PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes. *Bioinformatics* 26:1608–1615. <http://dx.doi.org/10.1093/bioinformatics/btq249>.
- Vesth T, Lagesen K, Acar Ö, Ussery D. 2013. CMG-BioTools, a free workbench for basic comparative microbial genomics. *PLoS One* 8:e60120. <http://dx.doi.org/10.1371/journal.pone.0060120>.
- Pittet V, Abegunde T, Marfleet T, Haakensen M, Morrow K, Jayaprakash T, Schroeder K, Trost B, Byrns S, Bergsveinson J, Kusalik A, Ziola B. 2012. Complete genome sequence of the beer spoilage organism *Pediococcus clausenii* ATCC BAA-344T. *J Bacteriol* 194:1271–1272. <http://dx.doi.org/10.1128/JB.06759-11>.
- Makarova K, Slesarev A, Wolf Y, Sorokin A, Mirkin B, Koonin E, Pavlov A, Pavlova N, Karamychev V, Polouchine N, Shakhova V, Grigoriev I, Lou Y, Rohksar D, Lucas S, Huang K, Goodstein DM, Hawkins T, Plengvidhya V, Welker D, Hughes J, Goh Y, Benson A, Baldwin K, Lee JH, Diaz-Muniz I, Dosti B, Smeianov V, Wechter W, Barabote R, Lorca G, Altermann E, Barrangou R, Ganesan B, Xie Y, Rawsthorne H, Tamir D, Parker C, Breidt F, Broadbent J, Hutkins R, O'Sullivan D, Steele J, Unlu G, Saier M, Kleenhammer T, Richardson P, Kozyavkin S, Weimer B, Mills D. 2006. Comparative genomics of the lactic acid bacteria. *Proc Natl Acad Sci U S A* 103:15611–15616. <http://dx.doi.org/10.1073/pnas.0607171103>.
- Behr J, Geissler AJ, Preissler P, Ehrenreich A, Angelov A, Vogel RF. 2015. Identification of ecotype-specific marker genes for categorization of beer-spoiling *Lactobacillus brevis*. *Food Microbiol* 51:130–138. <http://dx.doi.org/10.1016/j.fm.2015.05.015>.
- Bergsveinson J, Pittet V, Ewen E, Baecker N, Ziola B. 2015. Genome sequence of rapid beer-spoiling isolate *Lactobacillus brevis* BSO 464. *Genome Announc* 3(6):e01411-15. <http://dx.doi.org/10.1128/genomeA.01411-15>.
- Fukao M, Oshima K, Morita H, Toh H, Suda W, Kim SW, Suzuki S, Yakabe T, Hattori M, Yajima N. 2013. Genomic analysis by deep sequencing of the probiotic *Lactobacillus brevis* KB290 harboring nine plasmids reveals genomic stability. *PLoS One* 8:e60521. <http://dx.doi.org/10.1371/journal.pone.0060521>.
- Suzuki K, Iijima K, Ozaki K, Yamashita H. 2005. Isolation of a hop-sensitive variant of *Lactobacillus lindneri* and identification of genetic markers for beer spoilage ability of lactic acid bacteria. *Appl Environ Microbiol* 71:5089–5097. <http://dx.doi.org/10.1128/AEM.71.9.5089-5097.2005>.
- Sakamoto K, Margolles A, van Veen HW, Konings WN. 2001. Hop resistance in the beer spoilage bacterium *Lactobacillus brevis* is mediated by the ATP-binding cassette multidrug transporter *HorA*. *J Bacteriol* 183:5371–5375. <http://dx.doi.org/10.1128/JB.183.18.5371-5375.2001>.
- Bergsveinson J, Baecker N, Pittet V, Ziola B. 2015. Role of plasmids in *Lactobacillus brevis* BSO 464 hop tolerance and beer spoilage. *Appl Environ Microbiol* 81:1234–1241. <http://dx.doi.org/10.1128/AEM.02870-14>.
- Suzuki K. 2015. Gram-positive spoilage bacteria in brewing, p 141–174. *In* Hill AE (ed), *Brewing microbiology managing microbes, ensuring quality and valorising waste*. Woodhead publishing series in food science, technology, and nutrition. Woodhead Publishing, Elsevier, Cambridge, United Kingdom.