



The wild tomato species *Solanum chilense* shows variation in pathogen resistance between geographically distinct populations

Remco Stam, Daniela Scheikl and Aurélien Tellier

Section of Population Genetics, Technical University of Munich, Freising, Germany

ABSTRACT

Wild tomatoes are a valuable source of disease resistance germplasm for tomato (*Solanum lycopersicum*) breeders. Many species are known to possess a certain degree of resistance against certain pathogens; however, evolution of resistance traits is yet poorly understood. For some species, like *Solanum chilense*, both differences in habitat and within species genetic diversity are very large. Here we aim to investigate the occurrence of spatially heterogeneous coevolutionary pressures between populations of *S. chilense*. We investigate the phenotypic differences in disease resistance within *S. chilense* against three common tomato pathogens (*Alternaria solani*, *Phytophthora infestans* and a *Fusarium* sp.) and confirm high degrees of variability in resistance properties between selected populations. Using generalised linear mixed models, we show that disease resistance does not follow the known demographic patterns of the species. Models with up to five available climatic and geographic variables are required to best describe resistance differences, confirming the complexity of factors involved in local resistance variation. We confirm that within *S. chilense*, resistance properties against various pathogens show a mosaic pattern and do not follow environmental patterns, indicating the strength of local pathogen pressures. Our study can form the basis for further investigations of the genetic traits involved.

Submitted 20 September 2016
Accepted 8 December 2016
Published 18 January 2017

Corresponding author
Remco Stam, stam@wzw.tum.de

Academic editor
Erica Goss

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.2910

© Copyright
2017 Stam et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Ecology, Evolutionary Studies, Plant Science

Keywords Wild tomato, Phytopathology, *Fusarium*, *Phytophthora*, *Alternaria*, General linearised mixed models

INTRODUCTION

In nature, plants are exposed to a wide range of pathogens and pests. While in most cases the plants appear non-specifically resistant against these threats, drastic or recurrent epidemics do occur (*Thrall, Burdon & Bock, 2001; Soubeyrand et al., 2009*) and variability in specific resistance to pathogens is observed (*Thrall, Burdon & Young, 2001; Salvaudon, Giraud & Shykoff, 2008*). Understanding how reciprocal co-adaptation of hosts and pathogens maintains such diversity has been a key question in theoretical and empirical evolutionary biology. Theoretically, negative direct frequency-dependent selection (ndFDS) is shown to be a necessary condition to maintain long-term stable diversity for resistance in plants and infectivity in pathogens (*Tellier & Brown, 2007*). Seed banking, perenniality or polycyclic disease can generate ndFDS, while costs of resistance and infectivity (virulence) are

necessary but not sufficient for stable long term polymorphism to occur ([Tellier & Brown, 2009](#); [Brown & Tellier, 2011](#)). Another factor often suggested to maintain diversity is the spatial structure of host and pathogen populations. Spatial structure and migration of hosts and pathogens as well as population sizes and genetic drift generate patterns of local adaptation over space and time ([Thrall & Burdon, 2002](#); [Gandon & Michalakis, 2002](#)). However, a spatial structure with homogeneous environment does not generate ndFDS ([Thrall, Burdon & Bever, 2002](#); [Tellier & Brown, 2011](#)). Stable long term polymorphism is favoured by spatially heterogeneous environments across which the prevalence and severity of disease or the costs of resistance and infectivity may differ ([Gavrilets & Michalakis, 2008](#); [Moreno-Gamez, Stephan & Tellier, 2013](#)).

From an ecological perspective, and based on the classic disease triangle from plant pathology ([Agrios, 2005](#)) the outcome of species interactions are mediated by the abiotic and biotic environment. The influence of the environment therefore generates spatial and temporal variation in evolutionary and coevolutionary dynamics ([Thompson, 2005](#)), and increasing evidence for geographical variation in coevolutionary dynamics and patterns of local adaptation are found in microcosm experiments ([Forde, Thompson & Bohannan, 2004](#); [Vogwill et al., 2009](#); [Lopez-Pascua, Brockhurst & Buckling, 2010](#)).

Nevertheless, few field systems exist to study and document the coevolution of plants and their pathogens occurring at short time scales and across several populations. One example is the wild flax—flax rust pathosystem, where local adaptations have been observed and the most resistant varieties of flax generally harboured more virulent strains of rust ([Thrall, Burdon & Bever, 2002](#); [Thrall & Burdon, 2003](#)). Similarly, the local adaptation of powdery mildew *Podosphaera plantaginis* to *Plantago lanceolata* populations spread over different islands off the coast of Sweden showed virulent strains to co-occur with more resistant plants ([Laine, 2005](#); [Soubeyrand et al., 2009](#)). In the latter plant-pathogen system, several theoretically proposed mechanisms to generate ndFDS have been shown to originate from the environmental heterogeneity across populations: (1) GxGxE interactions (host genotype x pathogen genotype x environment, (for example [Laine, 2006](#)), (2) heterogeneity in disease incidence and prevalence determining epidemiological pressures ([Soubeyrand et al., 2009](#)) and co-infection ([Susi et al., 2015](#)) and (3) different strength of connectivity between populations accelerating or decelerating the speed of coevolution across the landscape ([Jousimo et al., 2014](#)). Thus, these factors are expected to promote and facilitate long term polymorphism at resistance and infectivity loci without unrealistic costs of these alleles. Here we aim to investigate the variation of disease resistance properties amongst populations of *S. chilense*, a wild tomato species, against several pathogens.

Wild *Solanum* species are in general particularly good model species to study between and within species variation, because they occur in diverse geographic and climatic habitats and have a very well studied demography and known evolutionary history ([Städler, Roselius & Stephan, 2005](#); [Städler, Arunyawat & Stephan, 2008](#); [Tellier et al., 2011](#)). Additionally, several studies exist suggesting that bacterial resistance-associated genes are under selective pressure ([Rose et al., 2005](#); [Rose et al., 2011](#); [Rose, Michelmore & Langley, 2007](#)). *S. chilense* is native in South America, ranging from southern Peru to central Chile, and colonised a broad range of habitats. *S. chilense* populations are found from coastal regions, even in

slightly alkaline environments, all the way to high altitude (>3,000 m) mountain regions. It also occurs in extreme dry habitats on the border of the Atacama desert, as well as near rivers and creeks (Peralta, Spooner & Knapp, 2008).

S. chilense most likely originated with its sister species *S. peruvianum*, in south Peru and then migrated south (Städler, Arunyawat & Stephan, 2008). A study of the species' demography found four genetically distinct groups; one in the north of the range, one in the central region and two in the south (one on the coast and one at high altitudes). Interestingly, the two southern groups are, even though geographically close to each other, more related to the central group than to each other, possibly due to the separating effect of the extremely arid Atacama desert (Böndel et al., 2015). In addition, *S. chilense* shows clear climatic adaptations. Populations from drier regions are responding faster to drought (Fischer et al., 2013) and individual populations found at high altitudes (>3,000 m) show higher freezing tolerance (Nosenko et al., 2016). *S. chilense* has also been the source of resistance loci against the fungus *Verticillium dahliae* (Tabaeizadeh et al., 1999) and against various viruses (Griffiths & Scott, 2001; Ji et al., 2007; Verlaan et al., 2013). Seeing that *S. chilense* occurs in such a wide range of habitats and that the species shows specific signs of local climatic adaptations, we wondered whether we could find variation for pathogen resistance as well.

Since no detailed data exist about the co-occurrence of wild pathogens and *S. chilense*, we chose to test *S. chilense* disease resistance properties with three widely studied and economically relevant pathogens, *Alternaria solani*, *Phytophthora infestans* and a *Fusarium* sp.

A. solani causes early blight and is amongst the most destructive diseases of tomato in tropical and subtropical regions, leading to yield losses of up to 80% in certain regions. *A. solani* has been found in central Peru and is known to cause disease not only on potato—its main host—but also on many other nightshades, including tomato (Song et al., 2011; Kumar et al., 2013). In addition, previous work has shown that *A. solani* resistance can be studied using detached leaf assays (Chaerani & Voorrips, 2006; Chaerani et al., 2007).

Fusarium spp. are pathogens that cause very severe disease symptoms on a very wide range of host plants that span almost the entire globe (Agrios, 2005). Two *Fusarium* spp are in the top ten most important fungi in plant pathology (Dean et al., 2012). The *F. oxysporium* species complex comprises over 100 formae specialis that all infect specific hosts, including tomato (Michielse & Rep, 2009). The *F. oxysporium* species complex is widely used to study molecular and genetic mechanisms involved in plant pathogen interactions (Houterman, Cornelissen & Rep, 2008; Ma, Cornelissen & Takken, 2013) and even though it is generally reported to be a vascular pathogen, it has regularly been successfully deployed in detached leaf infection assays (e.g., Kavroulakis et al., 2007).

Phytophthora infestans is an oomycete that causes late blight on potato and tomato. In potato alone the damage amounts up to \$1 bn annually (Haverkort et al., 2009). Due to its economic value and the vast amount of molecular and genetic research performed on it, it is considered the most important oomycete plant pathogen (Kamoun et al., 2015). Like the other two pathogens used in this study, *P. infestans* is known to occur in the geographic range of various wild *Solanum* species (Perez et al., 2001). Specifically, the strain EC1 that we used here has its origin in Ecuador and is particularly relevant for agriculture as it is

a rather aggressive strain that is capable of overcoming certain novel genetic resistances (Foster et al., 2009; Nowicki et al., 2011).

Here we test the resistance of different *S. chilense* populations. We selected seven populations that represent three previously described genotype groups from Peru and Chile (Böndel et al., 2015). Two populations originate from the central range (LA1958, LA3111), two from the coastal regions (LA2932, LA4107) and two from the southern mountainous region (LA4117, LA4330). A seventh population is geographically in the middle between the southern mountain and the central group (LA2931). Böndel et al. group it with the central populations, but assign properties of both groups to it. Figure 1A shows the species distribution and highlights the selected populations. These groups correspond to very distinctive habitats and can thus be used to investigate whether we see differences in infection rate throughout the range of the species. We also test whether these differences show a linear pattern when tested against geographical and climatic variables (e.g., correlation between latitude or precipitation and resistance) or whether a multitude of factors leads to specific geographical variations in resistance to each of the three pathogens. Our study will help inform on the occurrence of spatially heterogeneous coevolutionary pressures between the populations of *Solanum chilense*, a solanaceous wild species, and its pathogens.

METHODS

Plant growth

Seed batches were obtained from the tomato genomics resource centre (TGRC, Davis, USA). We grew seven different *Solanum chilense* populations (accession numbers LA1963, LA2931, LA2932, LA3111, LA4107, LA4117 and LA4330) consisting of 10 different plants each and one *Solanum pennellii* (LA0716) population in our glasshouse from randomly chosen seeds. The plants were grown with 16 h light and a minimum temperature of 18 °C. Mature plants were cut back at a biweekly interval to assure young leaves of similar age were available at all times for all populations.

Pathogen propagation and spore production

Alternaria solani

A. solani strains B055 and St108 were obtained from the chair of Phytopathology at the TUM (Munich, Germany) (Leiminger et al., 2016). The strain was originally collected from potato, though *A. solani* strains are known to be generalists with strains capable of infection both potato and tomato. *A. solani* and cultivated on SNA (Salt Nutrient Agar) plates (at 22 °C, 12 h UV-A light, 12 h darkness and 85% humidity) for three weeks to induce sporulation. We harvested the spores by submerging the cultures in with ddH₂O and scratching the mycelium off the agar with a microscope slide. The solution was filtered through four layers of mesh (grade 60, approximately 12 × 12 threads per cm²) and diluted to a concentration of 5,000 spores per ml. Each leaflet was infected with a 10 µl droplet.

Phytophthora infestans

We obtained late blight pathogen *P. infestans* strain EC1 from the James Hutton Institute (Dundee, UK). Although originally obtained from potato, samples belonging to the EC1




Figure 1 *S. chilense* populations and phenotypic observations. (A) A map showing the populations used in this study, belonging to the central (red), southern mountainous (blue) or southern coastal (green) region. The geographic range of the whole species is depicted in the background (grey dots). (B) The phenotypic observations after infection range from no visible symptoms (No Symptoms row) and small black necrotic lesions resembling the Hypersensitive Response (HR), both scored as ‘not infected,’ to intermediate and strong infection (third and fourth rows: infection and infection, respectively), both scored as ‘infected.’ Columns indicate infection with *Alternaria*, *Fusarium* and *Phytophthora*. We could not observe HR in the *Alternaria* infections. (C) Experimental set-up. Detached leaves were randomised for each population (P#) and spread over various boxes. Each leaflet was drop inoculated and scored. Here only the first five boxes are drawn.

lineage have been found on other solanaceous hosts (Vargas *et al.*, 2008). It was cultivated on RyeB agar, incubated six days at RT in darkness, three days at RT in daylight. We scratched the sporangia from the plate with ice cold water using a pipette tip and stored it at 4 °C until further use (up to three hours). The solution was diluted to 2,000–3,000 sporangia per ml and the leaflets were infected with 5 µl of this solution.

Fusarium sp

Fusarium infected lesions were identified on a few detached *S. chilense* leaves from our glasshouse. These lesions were extracted and re-cultivated for several rounds on Potato-Dextrose-Agar (PDA) for clean-up. Microscopic observations and sequence analysis of cloned Beta-Tubulin genes from several individual cultures confirmed that the cultures were axenic (Data S7). We used BLAST with the said Tubulin sequences to search against the fungidb.org database and to confirm the genus. The clean *Fusarium* was grown on PDA for a minimum of four days at RT. Spores were harvested by adding ddH₂O and aspirating the liquid. The spores were diluted to 2×10^5 – 5×10^5 spores per ml and we infected the individual leaflets with 5 µl of this solution.

All protocols for pathogen cultivation, including ingredients for the growth media can be found in more detail at <https://www.protocols.io/view/Plant-Pathogen-Cultivation-fmkbk4w>.

Leaf pre-treatment

In some wild tomato species (e.g., *S. pennellii*), thick and sticky surface coatings have a dramatic effect on pathogen ingress. In *S. chilense*, surface coatings are notably less thick, and resemble those of cultivated tomato. However, the selected *S. chilense* populations have slightly different leaf morphology. Different amounts of wax could be present and could cause differences in infection rates between the populations. We aimed to look at differences in molecular defence rather than physical barriers preventing hyphal ingress. Hence, to minimise the effect of differences in surface coating, we washed the leaves briefly in 70% ethanol, followed by rinsing with distilled water and a brief drying period before infection. This ethanol treatment has the added effect of surface sterilisation of the leaves, preventing bacterial contamination. The effects of *S. chilense* surface sterilisation and washing off the wax is noticeable during infection, but not as dramatic as with *S. pennellii* (Data S1).

Infection assays

To assess the infection rate of the different pathogens, we performed spot inoculations on detached leaves. Each spot was scored as either negative (no symptoms or only a small hypersensitive response lesion) or positive (larger (expanding) lesions or full proliferation of the pathogen) (Fig. 1B). To minimise the effect of variation between plants within one population, we randomly collected leaves with the same age from eight to ten plants per population and shuffled them. We then drew the leaves from that mix to distribute them over up to nine boxes for each infection experiment. Each box contained 16 leaves, in four columns with four rows. Each column corresponding to a different population. Each box contained different combinations of populations (Fig. 1C). Box number and leaf position

were marked to later rule out possible effects. For each pathogen 16–24 leaves—about 100 leaflets—were infected for each population. Over the course of several months, we performed four biological replicates for each population, accumulating to about 450–500 infected leaflets per population per pathogen. The *Alternaria* infections were done on the adaxial side of the leaves, *Phytophthora* and *Fusarium* infections were done on the abaxial side of the leaves. The leaves were incubated at RT and scored after six to eight days, dependent seasonal variability of temperature and light conditions in the lab, which slightly affected the rate of symptom development.

Data analysis

All data analysis was done using R (version 3.2.3) (R foundation for statistical computing). Generalised Linear Mixed Models (GLMM) were made using the glmer option from the package lme4 (Bates et al., 2015, p. 4). To construct GLMM we used a binomial variable (y) consisting of the number of successful and unsuccessful infection events per leaf. The GLMM were constructed taking the leaf position in the box (leaf) and a combination of the box number and experimental date (exp:box) into account as random effects. For our first model populations names were used as fixed effects. (model 1 = $y \sim$ accession + (1|leaf) + (1|exp : box)). For the next models, we hierarchically tested different climatic and geographical parameters (e.g., model 2 = $y \sim$ geographic1 + climatic1 + climatic2 + (1|leaf) + (1|exp : box)). Pairwise comparisons were examined using an implementation of Tukey Honest Significant Difference test as provided by function glht from the R package multcomp (Hothorn, Bretz & Westfall, 2008). glht allows post-hoc hypothesis testing, similar to THSD, but is more suitable for general linearised (mixed) models. The boxplots were drawn using the package ggplot2 (Wickham, 2009, p. 2). All packages are available through CRAN.

Distribution map and geographical characteristics

Geographical data for all populations were obtained from the Tomato Genome Resource Centre. Climatic data were extracted from the <http://worldclim.org/> database. The species distribution map was drawn using the maps package (Becker & Wilks, 1995) in R. All geographic and climate data used can be found in [Data S2](#).

RESULTS

S. chilense populations show different resistant properties against different pathogens

To test infection rates, we infected individual leaflets for up to 16 leaves of each *S. chilense* population with *Alternaria solani* (st108) and counted the occurrence of infected leaflets per leaf, as this represents the success rate of the pathogen to establish itself and overcome genetic resistance. We scored infection events as either negative (no infection or clear small necrotic lesions, indicating a hypersensitive response) or positive (ranging from growth just outside the droplet area up to full infection of the leaflet) (Fig. 1B) (All raw data can be found in [Data S3](#)). We observed variation within each population. In almost all instances at least one leaf was fully infected whereas another was completely resistant. These outliers




Figure 2 Infected leaf fraction for different *S. chilense* populations. The boxplots show the median and 1st and 3rd quartile of the infected fractions per leaf for (A) *Alternaria*, (B) *Fusarium* and (C) *Phytophthora*. The black dots represent the mean value for the infections. The Y axis ranges from 0 (no infected leaflets on a leaf) to 1 (all leaflets show infection). On the X axis, each population is represented. The colours correspond to the geographic regions as depicted in Fig. 1.

have large effect on the calculated mean fraction. To allow good judgement we report the 1st and 3rd quartile, the median value as well as the mean value for each population (Fig. 2). The mean and median of the infected fractions range from 0.35 and 0.42 for LA3111 to 0.74 and 0.81 for LA4330 or 0.67 and 0.82 for LA2932.

To test the robustness of our method, we did an additional infection with a second strain of *Alternaria* (B055). The overall infection rates were lower in this set of experiments (median of 0.54 compared to 0.62), however Data S4 shows that just like for strain st108, LA3111 is the least infected population with a mean of 0.40 and LA4330 and LA2932 have a high median, with an infected fraction of 0.70 or 0.73 respectively.

With *Fusarium* we also saw differences in the infected fraction amongst populations. Interestingly LA3111 was in this case the most infected population (mean: 0.72, median: 0.82) whereas LA4107 was the least susceptible (mean = 0.28, median = 0.11).

Finally, for *P. infestans*, the infected fractions again showed a different pattern. There was a larger spread of the data as can be seen by the increased distance between the 1st and 3rd quartile and the lowest and highest mean and median fraction were closer together, ranging from 0.30 and 0.21 for LA3111 to 0.60 and 0.70 for LA4330 (Fig. 2C). LA3111, one population that seemed particularly resistant against *Alternaria* and *Phytophthora* seems to be the most susceptible to *Fusarium*.

To test the significance of the differences and the effect of the different populations on infection, we constructed a general linearised mixed model (GLMM). We assigned experimental parameters (data, box and leaf number) as random effects and tested whether there were significant differences between the populations for each infecting species by looking at the infection counts (y) per leaf. These models showed that indeed there are highly significant differences ($p < 0.00001$) in infection rates between some populations for all three pathogens tested (Data S2).

Pairwise comparisons reveal individual differences between different pathogens

To further determine which populations were different from each other, we performed pairwise comparisons using a variant of Tukey's Honest Significant Difference test (see

Alternaria	Fusarium	Phytophthora	
-0.55183	0.10738	-0.26040	LA1963 – LA2931
-0.87557	-0.78253	-0.73009	LA1963 – LA2932
0.29545	-1.11537	0.26290	LA1963 – LA3111
-0.85150	0.71985	-0.26015	LA1963 – LA4107
-0.46488	0.88139	-0.42327	LA1963 – LA4117
-1.37044	0.02266	-1.23225	LA1963 – LA4330
-0.32373	-0.88991	-0.46969	LA2931 – LA2932
0.84728	-1.22275	0.52330	LA2931 – LA3111
-0.29967	0.61247	0.00025	LA2931 – LA4107
0.08695	0.77401	0.16287	LA2931 – LA4117
-0.81860	-0.08472	-0.97185	LA2931 – LA4330
1.17102	-0.33285	0.99300	LA2932 – LA3111
0.02407	1.50237	0.46994	LA2932 – LA4107
0.41069	1.66391	0.30681	LA2932 – LA4117
0.49487	0.80518	-0.50215	LA2932 – LA4330
-1.14695	1.83522	-0.52305	LA3111 – LA4107
-0.76033	1.99676	-0.68618	LA3111 – LA4117
-1.66588	1.13803	-1.49516	LA3111 – LA4330
0.38662	0.16154	-0.16312	LA4107 – LA4117
-0.51894	-0.69719	-0.97210	LA4107 – LA4330
-0.90556	-0.85873	-0.80897	LA4117 – LA4330

Figure 3 Populations with significant different infected fractions. Graph depicting whether a pairwise difference shows a significant result for *Alternaria solani*, *Fusarium sp.* and *Phytophthora infestans*. Each row represents a pairwise comparison. Green cells represent a significant difference ($p < 0.001$ after multiple testing correction) and the numbers represent the estimated effect, with negative numbers indicating that the population mentioned on the left is less resistant than the one on the right.

methods). The observed pairwise differences were clearly distinct between the three pathogens. [Figure 3](#) shows a summary of the pairwise differences, with corresponding difference estimates for each comparison. Cells with significant differences ($p < 0.001$) are highlighted in green. All pairwise differences with their 95% confidence intervals are plotted in [Data S5](#). Of the 63 pairwise comparisons, 32 showed a significant difference in infection ratio. Overall, there were more significant differences between populations when it came to *Fusarium* infection (15) than to *Alternaria* infection (10) or *Phytophthora* (7). Interestingly, some populations showed the same result for all pathogens: there are no differences between LA1963 and LA2931 (both central) nor for LA2931 and LA4107 (central and south coast) or LA4107 and LA4117 (south coast and south mountain). Also, LA1963 was always more susceptible than LA2932, and LA4117 was always more susceptible than LA4330. In some cases a population in a pair was more resistant to one pathogen and more susceptible to another. LA4330 was more resistant than LA3111 to *Fusarium*, but less resistant to *Alternaria* and *Phytophthora*.

A mix of climatic and geographic variables affect pathogen resistance

To see whether a change in certain geographic and climatic conditions could be linked to an increase or decrease of resistance rates between populations, we built new GLMM using such data. First we made a simple model for resistance to *Alternaria*, testing the infection counts (y) against either latitude or longitude, a combination of both or an interaction of both. This showed that both latitude and longitude had a significant effect ($p < 0.001$,

Table 1 Summary of GLMM results. AIC is a measure of relative quality of the model, with a lower AIC (within one species) indicating a better model. AICs in bold represent models that are significant ($p < 0.001$). TempA denotes the temperature in the wettest quarter, TempB in the coldest quarter and TempC the annual mean temperature.

	Model	<i>Alternaria</i>	<i>Fusarium</i>	<i>Phytophthora</i>
1	y ~accession	2641.8	2307.6	1893.3
2	y ~Lat	2708.6	2431.3	1958.3
3	y ~Long	2815.1	2490.8	1965.8
4	y ~Long + Lat	2703.9	2420.6	1945.4
5	y ~Long*Lat	2705.8	2419.1	1947.4
6	y ~Long + Lat + Alt + AnnPrecip + TempA + TempB	2641.8	2307.6	1893.3
6b	y ~Long + AnnPrecip + TempA + TempB	2753.2	2303.6	1927.7
7	y ~Altitude	2843.3	2503.5	1985.1
8	y ~TempA	2842.2	2506.5	1987.0
9	y ~TempB	2842.2	2506.5	1981.3
10	y ~TempC	2842.2	2506.5	1984.1
11	y ~AnnPreci	2784.2	2457.1	1968.4
12	y ~group	2705.0	2489.7	1930.2

Table 1). The quality of the models is reported by the statistical software using the Akaike Information Criterion (AIC), where a lower AIC, indicates a better relative quality of the model. A model with both parameters showed a better AIC. However, a model with an interaction only shows significance of the latitude parameter. We extended the model to include both parameters (longitude + latitude) as well as several environmental parameters. We obtained the best AIC (2641.8) for a model containing altitude, annual precipitation, the temperature in the wettest and the temperature in the coldest quarter. Additions of other climatic data did not yield an improvement of the model. A selection of tested models with their AIC and significance is shown in Table 1. When closely investigating the best fitting model for resistance to *Alternaria* (Table 1, model 6), we noted that whereas all variables contribute significantly to the model, the estimated effect size differs greatly. Table 2 shows that of all effects, longitude was the strongest effect, followed by the mean minimum temperature in winter (TempB), the annual precipitation and altitude. It should be noted that models that only take temperature effects into account did not account for significance (Table 1). A GLMM with the infection counts set against the previously identified genetic groups (y ~group), yielded a higher and thus worse AIC (2705). The model with the populations yields an as good AIC as the one with six climatic variables. This suggested that no single variable has a strong, exclusive correlation to infection rate and that each population represented its own micro environment with specific geographic and climate parameters that are all of influence.

Similar to resistance to *Alternaria*, we tested all variables for resistance to *Phytophthora* and resistance to *Fusarium*. The pattern seen for resistance to *Phytophthora* is almost identical to that of *Alternaria* resistance. The AIC values are generally lower, but the trends are the same. Interestingly, *Fusarium* resistance showed a different picture. Whereas longitude was still the strongest effect, its effect and significance were lower and the temperature in the coldest quarter of the year had a relatively larger effect. The effect of

Table 2 Effects and significance of environmental variables, estimated size of the effect and the significance for all variables used in model 6 (Table 1). *P*-values in bold are deemed significant. TempA denotes the temperature in the wettest quarter, TempB in the coldest quarter.

Variable	<i>Alternaria</i>		<i>Fusarium</i>		<i>Phytophthora</i>	
	Effect estimate	<i>P</i> -value	Effect estimate	<i>P</i> -value	Effect estimate	<i>P</i> -value
Latitude	−0.5	<0.001	−0.1	0.8	−0.3	<0.001
Longitude	17.4	<0.001	7.9	0.007	17.3	<0.001
Altitude	−7.4	<0.001	0.0	0.99	−6.5	<0.001
Precipitation	9.1	<0.001	3.8	0.02	8.3	<0.001
TempA	3.2	<0.001	1.5	0.004	3.1	<0.001
TempB	10.2	<0.001	7.8	<0.001	10.6	<0.001

altitude was not significant and differences in annual precipitation had a nearly negligible effect as well. In fact, a simplified model (Table 1, model 6b) that excludes the insignificant parameters had a better fit. As with resistance to *Alternaria*, the model testing for the genetic group effect showed a lesser fit than the model per population (results for selected models can be found in Data S2).

DISCUSSION

The wild tomato *Solanum chilense* grows in a variety of habitats in Chile and Peru, ranging from lower coastal areas to very high altitudes (>3,000 m). These populations experience considerable variation in geographic parameters like precipitation and temperatures. It is known that *S. chilense* has a clear demographic pattern and signs of adaptations to climatic differences between different populations (Fischer et al., 2011; Fischer et al., 2013; Nosenko et al., 2016). A demographic pattern of North–South colonisation is observable with larger and more diverse populations in the north of the range and smaller and less diverse populations in the south. In addition, there is little to no genetic exchange between some of the southern most populations that are separated by the extremely dry Atacama desert. This leads to the conclusion that *S. chilense* can be divided in a northern, a central and two southern genotype groups (Böndel et al., 2015).

We hypothesised that pathogen pressures must differ a lot between such diverse geographical locations and as such *S. chilense* should show variation in resistance to common plant pathogens between the different populations. The underlying assumption is that at the evolutionary time scale of thousands of years, climatic conditions generate different selective pressures and presence/absence of pathogens, which then drives the evolution of the plant immune genes along different evolutionary paths in different regions. This heterogeneity in space may be seen as variation in functionality or duplication of resistance genes (so-called R-genes) and different levels of quantitative resistance across populations depending on pathogen availability and trade-offs for costs of resistance (Moreno-Gamez, Stephan & Tellier, 2013). The result of this evolutionary process was tested by performing infection assays in a common environment to assess the degree of resistance to three *Solanum* pathogens. We observed clear differences between the infection success rates of the three pathogens on the different *S. chilense* populations, indeed suggesting local adaptations to pathogen pressure. We could only observe a clear separation between

three regional *S. chilense* genotype groups for resistance to *Alternaria*, where the central populations are more susceptible than those from groups in the south. This shows that the genotype groups can be seen as an indicator of resistance against *Alternaria*. For resistance to the other pathogens large within-group differences exist and the genotype groups cannot be considered an indicator of resistance. Pairwise comparisons confirmed that outcomes differ within groups and between pathogens. For example, a pair that shows significant differences for resistance to *Phytophthora* and *Alternaria* (LA1963–LA4330) does not show differences in infection rate for *Fusarium*. Very strong pairwise differences can even be seen within the previously identified genotype groups (e.g., LA2932–LA4107 with *Fusarium*). We also showed that there are no populations that are in general more resistant (e.g., against all pathogens). For example, LA3111 is particularly resistant against *Fusarium*, but the most susceptible to *Phytophthora* and *Alternaria*.

We used a GLMM to test which factors might contribute to these differences. Interestingly, whereas the species *S. chilense* as a whole, shows a strong north–south demography, our analyses show that in models with multiple variables not latitude, but longitude has a very strong effect on infection rates. This could on the one hand be explained by the absence of the northern-most group in our analysis. On the other hand, a more likely explanation is the bigger geographic and associated climatic difference in the east–west gradient of the species, with low altitude coastal areas in the west, and high mountains in the east. Temperature differences can have large effects on the prevalence of pathogen populations as shown for wild plant-pathosystems (Laine, 2008) and also on crops, pathogens show adaptation to different climates (Mboup et al., 2012; Stefansson, McDonald & Willi, 2013). The mountainous areas in our study have particularly cold winters and relatively low mean temperatures in summer, which could be detrimental for pathogen survival or slow its growth and thus reduce pathogen pressure, which in turn might lead to loss of resistance in the plants.

Our results show indeed that inclusion of climatic variables for “temperature in winter” as well as “temperature in the wettest quarter” greatly improve our models and have a significant effect on infection rate for both *A. solani* and *P. infestans*. The importance of overwintering inoculum has previously been shown to be a main predictor for *Podosphaera plantaginis* epidemics on *Plantago lanceolata* in the next growing season (Soubeyrand et al., 2009). However, it must be noted that models that only incorporate winter temperature or indeed any other single climatic variables effects did not show any significance. This is in line with a between species comparison for wild potato (Spooner et al., 2009) and might be related to the fact that some higher altitude locations also have the highest annual precipitation rates. For example for *P. infestans* a relative high humidity has large positive effects on successful sporulation (Harrison & Lowe, 1989), which could counter the negative temperature effects.

Our climate data were extracted from <http://worldclim.org/> and might not provide the whole picture. For example, precipitation data might be accurate, but do not take into account a common sea-fog phenomenon that can be observed along the coast of Chile and Peru (Cereceda & Schemenauer, 1991; Schemenauer & Cereceda, 1992). This fog increases the local humidity for several hours up to several days in certain “fog basins”. Similarly, no

data is available on any nearby streams, rivers or irrigation canals for any of the populations. For some populations, a note is available for the state of the site at the time of collection (e.g., “dry quebrada”), but it remains unknown whether these features are constant or changed shortly before collection.

The best fitted models incorporate five climatic and geographic variables. Adding more variables did not improve the model, mainly due to the correlations between the available climate data. The strongest effects were observed for combinations of longitude and latitude together with climatic variables, indicating that one or two variables alone do not determine pathogen resistance. The latitude effect, which can be observed in the evolution of the species as a whole, seems to be less strong in our analyses, where longitude plays a larger role. Overall, our results indicate that indeed *S. chilense* populations show variations in pathogen resistance, which are possibly the result of adaptations to local pathogen pressures. The mosaic-like structure of our results indicates that these resistances are likely caused by a multitude of factors. These findings are in line with several inter-species studies in wild potato, where no correlation could be found between geographical location of the species and resistance against *P. infestans* (Khuutti et al., 2015) or *A. solani* (Jansky, Simon & Spooner, 2008). To further unravel the combination of factors contributing to local variations, new sampling excursions would be required, that not just collect plant and pathogen, but also measure local geographic and climatic parameters.

Several mechanisms have been theoretically proposed to generate stable long term polymorphism at host resistance and pathogen infectivity loci. Notably, many of those are linked to the environmental heterogeneity across populations. Here we show that (1) multiple environmental factors are required to improve the GLMM (2) infection rates differ between populations and even within previously genetic groups and (3) *S. chilense* populations show individual patterns in response to the three different pathogens. Therefore we conclude on (1) the existence of possible GxGxE interactions for given host-pathogen interactions, (2) heterogeneity in disease incidence and prevalence across habitats, and most interestingly (3) a likely geographic mosaic of exposure to different pathogens. However, additional experiments with locally collected pathogens are required to confirm these conclusions.

The presence-absence of different above- and below-ground pathogens on the same plants may be a key component of wild systems generating scenarios such as co-infection (Susi et al., 2015), cross-immunity or facilitation (Tack et al., 2015), with consequences for the genomics of pathogens (McMullan et al., 2015). Our research did not yet focus on any genetic differences underlying the variation in infection rates and linking phenotype to genotype should be one of the follow-up projects. Identification of the genes involved in these resistance variations could also help to identify which plant defence mechanisms are affected between populations and if there are indeed evolutionary differences between defence pathways in non-host resistance compared to resistance variation within or in closely related species (Schulze-Lefert & Panstruga, 2011; Stam et al., 2014). We have recently shown that targeted resequencing of genes of interest (e.g., R-genes) can be a potent tool to calculate evolutionary parameters of gene families of interest in wild tomato

(Stam, Scheikl & Tellier, 2016). Such resequencing studies could thus help to pinpoint how molecular mechanisms are affected by different pathogens as well as climatic variables.

CONCLUSIONS

Differences in pathogen disease resistance have been well described between many wild crop relatives. Here we presented a phenotypic study that shows large variation in resistance against three common tomato pathogens between populations of the wild tomato species *S. chilense*. We showed that there are clear differences between individual populations. Using generalised linear mixed models, we show that this variation does not follow a simple geographical cline, that multiple climatic factors are needed to explain parts of the variation and that even within previously identified genotype groups resistance properties can differ dramatically. Our study confirms a mosaic pattern in resistance properties within one species and can form the starting point for studies unravelling environmental effects on said properties as well as the genetic and molecular mechanisms involved in plant-pathogen coevolution.

ACKNOWLEDGEMENTS

We would like to thank Hannes Rief and Giulia Schiavoni for help with the scoring of the infection assays, Hans Hausladen, Michael Heß and Regine Ditteman (TUM Phytopathology) for providing the *Alternaria* strains and help with *Fusarium* isolation, and Brian Harrower (The James Hutton Institute, Dundee, UK) for providing *Phytophthora* and all reviewers and the editor for useful comments on the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Remco Stam was funded by the Alexander von Humboldt Stiftung. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Alexander von Humboldt Stiftung.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Remco Stam conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.

- Daniela Scheikl performed the experiments, wrote the paper, reviewed drafts of the paper.
- Aurélien Tellier conceived and designed the experiments, analyzed the data, reviewed drafts of the paper.

Data Availability

The following information was supplied regarding data availability:

The raw data has been supplied as a [Supplementary File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2910#supplemental-information>.

REFERENCES

- Agrios GN. 2005. *Plant pathology*. San Diego: Elsevier.
- Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67(1):1–48 DOI 10.18637/jss.v067.i01.
- Becker RA, Wilks AR. 1995. Constructing a geographical database. AT&T Bell Laboratories statistics research report [95.2]. Available at <http://ect.bell-labs.com/sl/doc/95.2.p>.
- Böndel KB, Lainer H, Nosenko T, Mboup M, Tellier A, Stephan W. 2015. North–south colonization associated with local adaptation of the wild tomato species *solanum chilense*. *Molecular Biology and Evolution* 32:2932–2943 DOI 10.1093/molbev/msv166.
- Brown JKM, Tellier A. 2011. Plant-parasite coevolution: bridging the gap between genetics and ecology. *Annual Review of Phytopathology* 49:345–367 DOI 10.1146/annurev-phyto-072910-095301.
- Cereceda P, Schemenauer RS. 1991. The occurrence of fog in Chile. *Journal of Applied Meteorology* 30:1097–1105 DOI 10.1175/1520-0450(1991)030<1097:TOOFIC>2.0.CO;2.
- Chaerani R, Groenwold R, Stam P, Voorrips RE. 2007. Assessment of early blight (*Alternaria solani*) resistance in tomato using a droplet inoculation method. *Journal of General Plant Pathology* 73(2):96–103 DOI 10.1007/s10327-006-0337-1.
- Chaerani R, Voorrips RE. 2006. Tomato early blight (*Alternaria solani*): the pathogen, genetics, and breeding for resistance. *Journal of General Plant Pathology* 72(6):335–347 DOI 10.1007/s10327-006-0299-3.
- Dean R, Van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ, Dickman M, Kahmann R, Ellis J, Foster GD. 2012. The top 10 fungal pathogens in molecular plant pathology. *Molecular Plant Pathology* 13(4):414–430 DOI 10.1111/j.1364-3703.2011.00783.x.
- Fischer I, Camus-Kulandaivelu L, Allal F, Stephan W. 2011. Adaptation to drought in two wild tomato species: the evolution of the Asr gene family. *New Phytologist* 190(4):1032–1044 DOI 10.1111/j.1469-8137.2011.03648.x.

- Fischer I, Steige KA, Stephan W, Mboup M. 2013.** Sequence evolution and expression regulation of stress-responsive genes in natural populations of wild tomato. *PLOS ONE* **8**:e78182 DOI [10.1371/journal.pone.0078182](https://doi.org/10.1371/journal.pone.0078182).
- Forde SE, Thompson JN, Bohannan BJM. 2004.** Adaptation varies through space and time in a coevolving host-parasitoid interaction. *Nature* **431**:841–844 DOI [10.1038/nature02906](https://doi.org/10.1038/nature02906).
- Foster SJ, Park TH, Pel M, Brigneti G, Sliwka J, Jagger L, Van der Vossen E, Jones JDG. 2009.** Rpi-vnt1.1, a Tm-22 homolog from *Solanum venturii*, confers resistance to potato late blight. *Molecular Plant-Microbe Interactions* **22**(5):589–600 DOI [10.1094/MPMI-22-5-0589](https://doi.org/10.1094/MPMI-22-5-0589).
- Gandon S, Michalakis Y. 2002.** Local adaptation, evolutionary potential and host-parasite coevolution: interactions between migration, mutation, population size and generation time. *Journal of Evolutionary Biology* **15**(3):451–462 DOI [10.1046/j.1420-9101.2002.00402.x](https://doi.org/10.1046/j.1420-9101.2002.00402.x).
- Gavrilets S, Michalakis Y. 2008.** Effects of environmental heterogeneity on victim–exploiter coevolution. *Evolution* **62**(12):3100–3116 DOI [10.1111/j.1558-5646.2008.00513.x](https://doi.org/10.1111/j.1558-5646.2008.00513.x).
- Griffiths PD, Scott JW. 2001.** Inheritance and linkage of tomato mottle virus resistance genes derived from *Lycopersicon chilense* accession LA 1932. *Journal of the American Society for Horticultural Science* **126**:462–467.
- Harrison JG, Lowe R. 1989.** Effects of humidity and air speed on sporulation of *Phytophthora infestans* on potato leaves. *Plant Pathology* **38**(4):585–591 DOI [10.1111/j.1365-3059.1989.tb01455.x](https://doi.org/10.1111/j.1365-3059.1989.tb01455.x).
- Haverkort AJ, Struik PC, Visser R.GF, Jacobsen E. 2009.** Applied biotechnology to combat late blight in potato caused by *phytophthora infestans*. *Potato Research* **52**(3):249–264 DOI [10.1007/s11540-009-9136-3](https://doi.org/10.1007/s11540-009-9136-3).
- Hothorn T, Bretz F, Westfall P. 2008.** Simultaneous inference in general parametric models. *Biometrical Journal* **50**(3):346–363 DOI [10.1002/bimj.200810425](https://doi.org/10.1002/bimj.200810425).
- Houterman PM, Cornelissen BJC, Rep M. 2008.** Suppression of plant resistance gene-based immunity by a fungal effector. *PLOS Pathogens* **4**:e1000061 DOI [10.1371/journal.ppat.1000061](https://doi.org/10.1371/journal.ppat.1000061).
- Jansky SH, Simon R, Spooner DM. 2008.** A test of taxonomic predictivity: resistance to early blight in wild relatives of cultivated potato. *Phytopathology* **98**(6):680–687 DOI [10.1094/phyto-98-6-0680](https://doi.org/10.1094/phyto-98-6-0680).
- Ji Y, Scott JW, Hanson P, Graham E, Maxwell DP. 2007.** Sources of resistance, inheritance, and location of genetic loci conferring resistance to members of the tomato-infecting begomoviruses. In: Czosnek H, ed. *Tomato Yellow Leaf Curl Virus Disease*. Netherlands: Springer, 343–362.
- Jousimo J, Tack AJM, Ovaskainen O, Mononen T, Susi H, Tollenaere C, Laine A-L. 2014.** Ecological and evolutionary effects of fragmentation on infectious disease dynamics. *Science* **344**:1289–1293 DOI [10.1126/science.1253621](https://doi.org/10.1126/science.1253621).
- Kamoun S, Furzer O, Jones JDG, Judelson HS, Ali GS, Dalio RJD, Roy SG, Schena L, Zambounis A, Panabières F, Cahill D, Ruocco M, Figueiredo A, Chen X-R,**

- Hulvey J, Stam R, Lamour K, Gijzen M, Tyler BM, Grünwald NJ, Mukhtar MS, Tomé DFA, Tör M, Van den Ackerveken G, McDowell J, Daayf F, Fry WE, Lindqvist-Kreuze H, Meijer HJG, Petre B, Ristaino J, Yoshida K, Birch PRJ, Govers F. 2015. The top 10 oomycete pathogens in molecular plant pathology. *Molecular Plant Pathology* **16**(4):413–434 DOI [10.1111/mpp.12190](https://doi.org/10.1111/mpp.12190).
- Kavroulakis N, Ntougias S, Zervakis GI, Ehaliotis C, Haralampidis K, Papadopoulou KK. 2007. Role of ethylene in the protection of tomato plants against soil-borne fungal pathogens conferred by an endophytic *Fusarium solani* strain. *Journal of Experimental Botany* **58**(14):3853–3864 DOI [10.1093/jxb/erm230](https://doi.org/10.1093/jxb/erm230).
- Khiutti A, Spooner DM, Jansky SH, Halterman DA. 2015. Testing taxonomic predictivity of foliar and tuber resistance to *phytophthora infestans* in wild relatives of potato. *Phytopathology* **105**(9):1198–1205 DOI [10.1094/PHYTO-02-15-0046-R](https://doi.org/10.1094/PHYTO-02-15-0046-R).
- Kumar S, Singh R, Kashyap PL, Srivastava AK. 2013. Rapid detection and quantification of *Alternaria solani* in tomato. *Scientia Horticulturae* **151**:184–189 DOI [10.1016/j.scienta.2012.12.026](https://doi.org/10.1016/j.scienta.2012.12.026).
- Laine AL. 2005. Spatial scale of local adaptation in a plant-pathogen metapopulation. *Journal of Evolutionary Biology* **18**(4):930–938 DOI [10.1111/j.1420-9101.2005.00933.x](https://doi.org/10.1111/j.1420-9101.2005.00933.x).
- Laine A-L. 2006. Evolution of host resistance: looking for coevolutionary hotspots at small spatial scales. *Proceedings of the Royal Society B: Biological Sciences* **273**:267–273 DOI [10.1098/rspb.2005.3303](https://doi.org/10.1098/rspb.2005.3303).
- Laine A-L. 2008. Temperature-mediated patterns of local adaptation in a natural plant-pathogen metapopulation. *Ecology Letters* **11**(4):327–337 DOI [10.1111/j.1461-0248.2007.01146.x](https://doi.org/10.1111/j.1461-0248.2007.01146.x).
- Leiminger JH, Auinger H-J, Wenig M, Bahnweg G, Hausladen H. 2016. Genetic variability among *Alternaria solani* isolates from potatoes in Southern Germany based on RAPD-profiles. *Journal of Plant Diseases and Protection* **120**(4):164–172 DOI [10.1007/BF03356470](https://doi.org/10.1007/BF03356470).
- Lopez-Pascua LDC, Brockhurst MA, Buckling A. 2010. Antagonistic coevolution across productivity gradients: an experimental test of the effects of dispersal. *Journal of Evolutionary Biology* **23**(1):207–211 DOI [10.1111/j.1420-9101.2009.01877.x](https://doi.org/10.1111/j.1420-9101.2009.01877.x).
- Ma L, Cornelissen BJC, Takken FLW. 2013. A nuclear localization for Avr2 from *Fusarium oxysporum* is required to activate the tomato resistance protein I-2. *Frontiers in Plant Science* **4**:Article 94 DOI [10.3389/fpls.2013.00094](https://doi.org/10.3389/fpls.2013.00094).
- Mboup M, Bahri B, Leconte M, De Vallavieille-Pope C, Kaltz O, Enjalbert J. 2012. Genetic structure and local adaptation of European wheat yellow rust populations: the role of temperature-specific adaptation. *Evolutionary Applications* **5**(4):341–352 DOI [10.1111/j.1752-4571.2011.00228.x](https://doi.org/10.1111/j.1752-4571.2011.00228.x).
- McMullan M, Gardiner A, Bailey K, Kemen E, Ward BJ, Cevik V, Robert-Seilaniantz A, Schultz-Larsen T, Balmuth A, Holub E, Van Oosterhout C, Jones JDG. 2015. Evidence for suppression of immunity as a driver for genomic introgressions and host range expansion in races of *Albugo candida*, a generalist parasite. *eLife* **4**:e04550 DOI [10.7554/eLife.04550](https://doi.org/10.7554/eLife.04550).

- Michielse CB, Rep M. 2009.** Pathogen profile update: *Fusarium oxysporum*. *Molecular Plant Pathology* **10**(3):311–324 DOI [10.1111/j.1364-3703.2009.00538.x](https://doi.org/10.1111/j.1364-3703.2009.00538.x).
- Moreno-Gamez S, Stephan W, Tellier A. 2013.** Effect of disease prevalence and spatial heterogeneity on polymorphism maintenance in host-parasite interactions. *Plant Pathology* **62**(S1):133–141 DOI [10.1111/ppa.12131](https://doi.org/10.1111/ppa.12131).
- Nosenko T, Böndel KB, Kumpfmüller G, Stephan W. 2016.** Adaptation to low temperatures in the wild tomato species *Solanum chilense*. *Molecular Ecology* **25**(12):2853–2869 DOI [10.1111/mec.13637](https://doi.org/10.1111/mec.13637).
- Nowicki M, Foolad MR, Nowakowska M, Kozik EU. 2011.** Potato and tomato late blight caused by *phytophthora infestans*: an overview of pathology and resistance breeding. *Plant Disease* **96**(1):4–17 DOI [10.1094/PDIS-05-11-0458](https://doi.org/10.1094/PDIS-05-11-0458).
- Peralta IE, Spooner DM, Knapp S. 2008.** The taxonomy of tomatoes: a revision of wild tomatoes (*Solanum* section *Lycopersicon*) and their outgroup relatives in sections *Juglandifolium* and *Lycopersicoides*. *Systematic Botany Monographs* **84**:1–186.
- Perez WG, Gamboa JS, Falcon YV, Coca M, Raymundo RM, Nelson RJ. 2001.** Genetic structure of Peruvian populations of *Phytophthora infestans*. *Phytopathology* **91**(10):956–965 DOI [10.1094/PHTO.2001.91.10.956](https://doi.org/10.1094/PHTO.2001.91.10.956).
- Rose LE, Grzeskowiak L, Hörger AC, Groth M, Stephan W. 2011.** Targets of selection in a disease resistance network in wild tomatoes. *Molecular Plant Pathology* **12**(9):921–927 DOI [10.1111/j.1364-3703.2011.00720.x](https://doi.org/10.1111/j.1364-3703.2011.00720.x).
- Rose LE, Langley CH, Bernal AJ, Michelmore RW. 2005.** Natural variation in the Pto pathogen resistance gene within species of wild tomato (*Lycopersicon*). I. Functional analysis of Pto alleles. *Genetics* **171**(1):345–357 DOI [10.1534/genetics.104.039339](https://doi.org/10.1534/genetics.104.039339).
- Rose LE, Michelmore RW, Langley CH. 2007.** Natural variation in the Pto disease resistance gene within species of wild tomato (*Lycopersicon*). II. Population genetics of Pto. *Genetics* **175**(3):1307–1319 DOI [10.1534/genetics.106.063602](https://doi.org/10.1534/genetics.106.063602).
- Salvaudon L, Giraud T, Shykoff JA. 2008.** Genetic diversity in natural populations: a fundamental component of plant–microbe interactions. *Current Opinion in Plant Biology* **11**(2):135–143 DOI [10.1016/j.pbi.2008.02.002](https://doi.org/10.1016/j.pbi.2008.02.002).
- Schemenauer R, Cereceda P. 1992.** Meteorological conditions at a coastal fog collection site in Peru. *Atmosfera* **6**:175–188.
- Schulze-Lefert P, Panstruga R. 2011.** A molecular evolutionary concept connecting nonhost resistance, pathogen host range, and pathogen speciation. *Trends in Plant Science* **16**:117–125 DOI [10.1016/j.tplants.2011.01.001](https://doi.org/10.1016/j.tplants.2011.01.001).
- Song W, Ma X, Tan H, Zhou J. 2011.** Abscisic acid enhances resistance to *Alternaria solani* in tomato seedlings. *Plant Physiology and Biochemistry: PPB/Société Française de Physiologie Végétale* **49**(7):693–700 DOI [10.1016/j.plaphy.2011.03.018](https://doi.org/10.1016/j.plaphy.2011.03.018).
- Soubeyrand S, Laine A-L, Hanski I, Penttinen A. 2009.** Spatiotemporal structure of host-pathogen interactions in a metapopulation. *The American Naturalist* **174**(3):308–320 DOI [10.1086/603624](https://doi.org/10.1086/603624).
- Spooner DM, Jansky SH, Simon R, Spooner DM, Jansky SH. 2009.** Tests of taxonomic and biogeographic predictivity: resistance to disease and insect pests in wild relatives of cultivated potato. *Crop Science* **49**:1367–1376.

- Städler T, Arunyawat U, Stephan W. 2008.** Population genetics of speciation in two closely related wild tomatoes (*Solanum* section *Lycopersicon*). *Genetics* **178**(1):339–350 DOI [10.1534/genetics.107.081810](https://doi.org/10.1534/genetics.107.081810).
- Städler T, Roselius K, Stephan W. 2005.** Genealogical footprints of speciation processes in wild tomatoes: demography and evidence for historical gene flow. *Evolution; International Journal of Organic Evolution* **59**(6):1268–1279 DOI [10.1111/j.0014-3820.2005.tb01777.x](https://doi.org/10.1111/j.0014-3820.2005.tb01777.x).
- Stam R, Mantelin S, McLellan H, Thilliez G. 2014.** The role of effectors in nonhost resistance to filamentous plant pathogens. *Frontiers in Plant Science* **5**:Article 582 DOI [10.3389/fpls.2014.00582](https://doi.org/10.3389/fpls.2014.00582).
- Stam R, Scheikl D, Tellier A. 2016.** Pooled enrichment sequencing identifies diversity and evolutionary pressures at NLR resistance genes within a wild tomato population. *Genome Biology and Evolution* **8**:1501–1515 DOI [10.1093/gbe/evw094](https://doi.org/10.1093/gbe/evw094).
- Stefansson TS, McDonald BA, Willi Y. 2013.** Local adaptation and evolutionary potential along a temperature gradient in the fungal pathogen *Rhynchosporium commune*. *Evolutionary Applications* **6**(3):524–534 DOI [10.1111/eva.12039](https://doi.org/10.1111/eva.12039).
- Susi H, Barrès B, Vale PF, Laine A-L. 2015.** Co-infection alters population dynamics of infectious disease. *Nature Communications* **6**:Article 5975 DOI [10.1038/ncomms6975](https://doi.org/10.1038/ncomms6975).
- Tabaeizadeh Z, Agharbaoui Z, Harrak H, Poysa V. 1999.** Transgenic tomato plants expressing a *Lycopersicon chilense* gene demonstrate improved resistance to *Verticillium dahliae* race 2. *Plant Cell Reports* **19**(2):197–202 DOI [10.1007/s002990050733](https://doi.org/10.1007/s002990050733).
- Tack A.JM, Laine A-L, Burdon JJ, Bissett A, Thrall PH. 2015.** Below-ground abiotic and biotic heterogeneity shapes above-ground infection outcomes and spatial divergence in a host–parasite interaction. *New Phytologist* **207**(4):1159–1169 DOI [10.1111/nph.13408](https://doi.org/10.1111/nph.13408).
- Tellier A, Brown JKM. 2007.** Stability of genetic polymorphism in host–parasite interactions. *Proceedings of the Royal Society B: Biological Sciences* **274**:809–817 DOI [10.1098/rspb.2006.0281](https://doi.org/10.1098/rspb.2006.0281).
- Tellier A, Brown J.KM. 2009.** The influence of perenniality and seed banks on polymorphism in plant–parasite interactions. *The American Naturalist* **174**(6):769–779 DOI [10.1086/646603](https://doi.org/10.1086/646603).
- Tellier A, Brown JKM. 2011.** Spatial heterogeneity, frequency-dependent selection and polymorphism in host–parasite interactions. *BMC Evolutionary Biology* **11**:319 DOI [10.1186/1471-2148-11-319](https://doi.org/10.1186/1471-2148-11-319).
- Tellier A, Fischer I, Merino C, Xia H, Camus-Kulandaivelu L, Städler T, Stephan W. 2011.** Fitness effects of derived deleterious mutations in four closely related wild tomato species with spatial structure. *Heredity* **107**:189–199 DOI [10.1038/hdy.2010.175](https://doi.org/10.1038/hdy.2010.175).
- Thompson JN. 2005.** *The geographic mosaic of coevolution*. Chicago: University of Chicago Press.
- Thrall PH, Burdon JJ. 2002.** Evolution of gene-for-gene systems in metapopulations: the effect of spatial scale of host and pathogen dispersal. *Plant Pathology* **51**:169–184 DOI [10.1046/j.1365-3059.2002.00683.x](https://doi.org/10.1046/j.1365-3059.2002.00683.x).

- Thrall PH, Burdon JJ. 2003.** Evolution of virulence in a plant host-pathogen metapopulation. *Science* **299**(5613):1735–1737 DOI [10.1126/science.1080070](https://doi.org/10.1126/science.1080070).
- Thrall PH, Burdon JJ, Bever JD. 2002.** Local adaptation in the *Linum marginale*-*Melampsora lini* host-pathogen interaction. *Evolution; International Journal of Organic Evolution* **56**:1340–1351.
- Thrall PH, Burdon JJ, Bock CH. 2001.** Short-term epidemic dynamics in the *Cakile maritima*-*Alternaria brassicicola* host-pathogen association. *Journal of Ecology* **89**(5):723–735 DOI [10.1046/j.0022-0477.2001.00598.x](https://doi.org/10.1046/j.0022-0477.2001.00598.x).
- Thrall PH, Burdon JJ, Young A. 2001.** Variation in resistance and virulence among demes of a plant host-pathogen metapopulation. *Journal of Ecology* **89**(5):736–748 DOI [10.1046/j.0022-0477.2001.00597.x](https://doi.org/10.1046/j.0022-0477.2001.00597.x).
- Vargas AM, Ocampo LMQ, Céspedes MC, Carreño N, González A, Rojas A, Zuluaga AP, Myers K, Fry WE, Jiménez P, Bernal AJ, Restrepo S. 2008.** Characterization of *phytophthora infestans* populations in Colombia: first report of the A2 mating type. *Phytopathology* **99**(1):82–88 DOI [10.1094/PHYTO-99-1-0082](https://doi.org/10.1094/PHYTO-99-1-0082).
- Verlaan MG, Hutton SF, Ibrahim RM, Kormelink R, Visser RGF, Scott JW, Edwards JD, Bai Y. 2013.** The tomato yellow leaf curl virus resistance genes Ty-1 and Ty-3 are allelic and code for DFDGD.-class RNA-dependent RNA polymerases. *PLOS Genetics* **9**:e1003399 DOI [10.1371/journal.pgen.1003399](https://doi.org/10.1371/journal.pgen.1003399).
- Vogwill T, Fenton A, Buckling A, Hochberg ME, Brockhurst MA. 2009.** Source populations act as coevolutionary pacemakers in experimental selection mosaics containing hotspots and coldspots. *The American Naturalist* **173**(5):E171–E176 DOI [10.1086/597374](https://doi.org/10.1086/597374).
- Wickham H. 2009.** *ggplot2: elegant graphics for data analysis*. New York: Springer-Verlag.