

# The 'heart' of preventive cardiology: Lifestyle medicine for the treatment of cardiometabolic diseases

Katharina Lechner<sup>1,2</sup>, Elke Lorenz<sup>3</sup> and Jonathan A Drezner<sup>4</sup>

European Journal of Preventive  
Cardiology  
0(00) 1–2  
© The European Society of  
Cardiology 2020



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/2047487319899107  
journals.sagepub.com/home/cpr



With great interest we read the paper by Arne Janssen and colleagues.<sup>1</sup> We commend the authors for raising awareness within the preventive cardiology community for non-alcoholic fatty liver disease (NAFLD) and its emerging role as a risk indicator for atherosclerotic cardiovascular disease (ASCVD). The implication of this new understanding is to pursue a more multidisciplinary approach to address the unacceptable global disease burden attributable to the twin pandemic of metabolic and cardiovascular disease, and nicely aligns with a recently published *JACC Council Perspective* arguing in favour of implementing preventive cardiology as a subspecialty of cardiovascular medicine.<sup>2</sup>

We share the authors' view that NAFLD is linked to and/or might even constitute a pathogenic component of ASCVD.<sup>1</sup> It is now widely accepted that the accumulation of dysfunctional, ectopic adipose tissue in the liver and other organs instigates a cluster of local and systemic coordinated pathophysiological events that have been linked to high-risk atherosclerosis. These include but are not limited to atherogenic dyslipidaemia, endothelial dysfunction, oxidative stress, insulin resistance, compensatory hyperinsulinaemia and diabetes.<sup>3</sup> It is therefore of little surprise, as eloquently outlined by the authors, that the presence of NAFLD deserves a thoughtful cardiovascular risk assessment and evaluation for subclinical atherosclerosis to detect individuals who might benefit from lifestyle and therapeutic interventions aimed at prevention and management of ASCVD.<sup>1</sup>

We would like to extend the discussion by adding to Figure I, which nicely depicts the array of subspecialties and diseases in which preventive cardiology and management of cardiometabolic disease should be included in the optimal care path for the patient.<sup>1</sup> While we agree that all of the mentioned subspecialties and associated diseases are of great importance and strongly associated with ASCVD, we believe that sports and exercise medicine, nutritional medicine, sleep medicine and public health medicine also should be included as subspecialties contributing to preventive cardiology that play a crucial role in ASCVD prevention. It is

important to note that lifestyle strongly modifies the genetically driven risk in ASCVD and that management of cardiometabolic diseases indeed must involve public health and lifestyle medicine.<sup>4</sup> Ensuring that every individual and/or patient receives access to comprehensive information and support for the implementation of physical exercise,<sup>5</sup> protective eating patterns (i.e. a low carb Mediterranean diet<sup>6</sup> and diets rich in long-chain n-3 fatty acids eicosapentaenoic acid and docosahexaenoic acid<sup>4</sup>), restorative sleep patterns and learning strategies to mitigate the physiologic response to distress<sup>4</sup> is of utmost importance. Furthermore, public health interventions aimed at decreasing sedentary time at work and the burden of environmental stressors such as tobacco smoke in public buildings, non-physiological light exposure (i.e. blue light) and noise and air pollution are necessary to control risk factors that are beyond the control of each individual.<sup>4</sup>

In conclusion, we would like to build upon the valuable framework presented by Arne Janssen and colleagues<sup>1</sup> by highlighting the paramount role of sports and exercise medicine, nutritional biochemistry, sleep medicine and public health medicine as important subspecialties in preventive cardiology that should be included in the care paths of cardiometabolic diseases. We posit that the metabolic signature of the high-risk phenotype is in large part an orchestrated and coordinated response to high-risk lifestyle risk factors.<sup>4</sup> Therefore, lifestyle medicine, both on an individual and on a public health level,

<sup>1</sup>Technical University of Munich, School of Medicine, Department of Prevention, Rehabilitation and Sports Medicine, Germany

<sup>2</sup>DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Germany

<sup>3</sup>Deutsches Herzzentrum München, Technische Universität München, Germany

<sup>4</sup>Department of Family Medicine and the Center for Sports Cardiology, University of Washington, USA

#### Corresponding author:

Katharina Lechner, Technical University Munich, German Heart Centre, Department of Cardiology, Lazarettstraße 36, 80636 Munich, Germany. Email: [contact@katharinalechner.net](mailto:contact@katharinalechner.net) Twitter: @KLechnerMD

has an indispensable role in the attempt to lower the unacceptable global burden of NAFLD and ASCVD.

### Author contribution

All authors contributed to this letter. KL did the literature search, and drafted the letter. EL and JAD reviewed and edited the manuscript. All authors approved the final version of the manuscript.

### Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

### References

1. Janssen A, Grobbee DE and Dendale P. Non-alcoholic fatty liver disease, a new and growing risk indicator for cardiovascular disease. *Eur J Prev Cardiol*. Epub ahead of print 4 December 2019. DOI: 10.1177/2047487319891783.
2. Shapiro MD, Maron DJ, Morris PB, et al. Preventive cardiology as a subspecialty of cardiovascular medicine: JACC Council Perspectives. *J Am Coll Cardiol* 2019; 74: 1926–1942.
3. Stahl EP, Dhindsa DS, Lee SK, et al. Nonalcoholic fatty liver disease and the heart: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2019; 73: 948–963.
4. Lechner K, von Schacky C, McKenzie AL, et al. Lifestyle factors and high-risk atherosclerosis: Pathways and mechanisms beyond traditional risk factors. *Eur J Prev Cardiol*. Epub ahead of print 13 August 2019. DOI: 10.1177/2047487319869400.
5. Lechner K, Halle M, Scherr J, et al. Exercise recommendations in athletes with coronary artery calcification. *Eur J Prev Cardiol*, Epub ahead of print 11 October 2019. DOI: 10.1177/2047487319881700.
6. Gepner Y, Shelef I, Komy O, et al. The beneficial effects of Mediterranean diet over low-fat diet may be mediated by decreasing hepatic fat content. *J Hepatol* 2019; 71: 379–388.