



Ragweed plants grown under elevated CO₂ levels produce pollen which elicit stronger allergic lung inflammation

Denise Rauer¹ | Stefanie Gilles¹ | Maria Wimmer^{2,3} | Ulrike Frank⁴ |
Constanze Mueller⁵ | Stephanie Musiol^{2,3} | Behnam Vafadari¹ | Lorenz Aglas⁶ |
Fatima Ferreira⁶ | Philippe Schmitt-Kopplin⁵ | Jörg Durner⁴ | Jana Barbro Winkler⁷ |
Dieter Ernst⁴ | Heidrun Behrendt² | Carsten B. Schmidt-Weber^{2,3} |
Claudia Traidl-Hoffmann^{1,8,9} | Francesca Alessandrini^{2,3}

¹Chair and Institute of Environmental Medicine, UNIKA-T, Technical University of Munich and Helmholtz Zentrum München, Augsburg, Germany

²Center of Allergy & Environment (ZAUM), Technical University of Munich (TUM) and Helmholtz Zentrum München, Munich, Germany

³Members of the German Center of Lung Research (DZL), Munich, Germany

⁴Institute of Biochemical Plant Pathology (BIOP), Helmholtz Zentrum München, Neuherberg, Germany

⁵BGC, Research Unit Analytical BioGeoChemistry, Helmholtz Zentrum München, Neuherberg, Germany

⁶Department of Biosciences, University of Salzburg, Salzburg, Austria

⁷Research Unit Environmental Simulation, Institute of Biochemical Plant Pathology, Helmholtz Zentrum München, Neuherberg, Germany

⁸Outpatient Clinic for Environmental Medicine, University Clinic Augsburg, Augsburg, Germany

⁹Christine-Kühne Center for Allergy Research and Education (CK-Care), Davos, Switzerland

Correspondence

Francesca Alessandrini, Center of Allergy & Environment (ZAUM), Technical University

Abstract

Background: Common ragweed has been spreading as a neophyte in Europe. Elevated CO₂ levels, a hallmark of global climate change, have been shown to increase ragweed pollen production, but their effects on pollen allergenicity remain to be elucidated.

Methods: Ragweed was grown in climate-controlled chambers under normal (380 ppm, control) or elevated (700 ppm, based on RCP4.5 scenario) CO₂ levels. Aqueous pollen extracts (RWE) from control- or CO₂-pollen were administered in vivo in a mouse model for allergic disease (daily for 3–11 days, n = 5) and employed in human in vitro systems of nasal epithelial cells (HNECs), monocyte-derived dendritic cells (DCs), and HNEC-DC co-cultures. Additionally, adjuvant factors and metabolites in control- and CO₂-RWE were investigated using ELISA and untargeted metabolomics.

Results: In vivo, CO₂-RWE induced stronger allergic lung inflammation compared to control-RWE, as indicated by lung inflammatory cell infiltrate and mediators, mucus hypersecretion, and serum total IgE. In vitro, HNECs stimulated with RWE increased indistinctively the production of pro-inflammatory cytokines (IL-8, IL-1β, and IL-6). In contrast, supernatants from CO₂-RWE-stimulated HNECs, compared to control-RWE-stimulated HNECs, significantly increased TNF and decreased IL-10 production in DCs. Comparable results were obtained by stimulating DCs directly with RWEs. The metabolome analysis revealed differential expression of secondary plant metabolites in control- vs CO₂-RWE. Mixes of these metabolites elicited similar responses in DCs as compared to respective RWEs.

Abbreviations: 9-OTrE, 9-oxo-10E,12Z,15Z-octadecatrienoic acid; ADO, adenosine; BAL, bronchoalveolar lavage; Cat, catalposide; DC, dendritic cell; HNEC, human nasal epithelial cell; ICI, inflammatory cell infiltration; IL, interleukin; ILC2, type 2 innate lymphoid cell; i.n., intranasal; IPCC, intergovernmental panel on climate change; LPS, lipopolysaccharide; Lumi, lumichrome; Mal, Malvidin; MoDC, monocyte-derived dendritic cell; PALMs, pollen-associated lipid mediators; pC4OG, p-Coumaryl alcohol 4-O-glucoside; Pel, pelargonidin; Q3OS, quercetin-3-O-sophoroside; RCP, representative concentration pathway; RWE, aqueous ragweed pollen extract; Th2, T helper type 2 cell; Treg, regulatory T cell.

Denise Rauer, Stefanie Gilles, Claudia Traidl-Hoffmann, and Francesca Alessandrini equal contribution.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Allergy* published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd

of Munich and Helmholtz Zentrum Munich, German Research Center for Environmental Health, Ingolstaedter Landstr. 1, D-85764 Neuherberg, Germany.
Email: franci@helmholtz-muenchen.de


Funding information

Christine-Kühne Center for Allergy Research and Education (CK-Care), HGF-HICAM Initiative.

Conclusion: Our results indicate that elevated ambient CO₂ levels elicit a stronger RWE-induced allergic response in vivo and in vitro and that RWE increased allergenicity depends on the interplay of multiple metabolites.

KEYWORDS

allergic lung inflammation, carbon dioxide, climate change, pollen metabolome, ragweed



GRAPHICAL ABSTRACT

Pollen from ragweed grown under elevated CO₂ levels (700 ppm, based on RCP4.5 scenario) elicit a stronger allergic inflammatory response in vitro and in vivo by: Enhancing pro-inflammatory cytokine release in DCs stimulated with RWE or RWE-conditioned HNEC supernatants and increasing lung inflammatory infiltrate and serum total IgE. Increased allergenicity of CO₂-RWE depends on the interplay of multiple metabolites.

1 | INTRODUCTION

Common ragweed (*Ambrosia artemisiifolia* L.) is native to North America, where 26% of the population is sensitized to its pollen,^{1,2} causing hay fever, asthma, and allergic rhinitis. In recent decades, this invasive neophyte has been spreading in Europe.^{3,4} In 2016, around 33 million Europeans were sensitized to ragweed and these numbers are estimated to more than double by 2041-2060.⁵ Because weed pollen are highly allergenic, even low exposure induces strong allergic reactions.⁶

This is important for understanding future health burdens, which will be heavily increased by climate change.⁷ As a result of rising temperatures and favorable precipitation, we will experience a more widespread distribution of ragweed across Europe, expanding from Central toward Northern and Eastern European countries.⁸⁻¹⁰ Rising temperatures lead to earlier pollen seasons of anemophilous plants in the Northern hemisphere, thereby increasing the abundance of airborne allergenic pollen.^{11,12} Additionally, rising atmospheric CO₂ levels are driving forces of climate change, which resulted in higher ragweed biomass and pollen production

in an experimental Intergovernmental Panel on Climate Change (IPCC) scenario.^{13,14} Likewise, elevated CO₂ levels combined with drought stress increased the amount of ragweed allergens (Amb a 1, Amb a 8 and Amb a 9) at the protein and transcriptional level.^{15,16}

The allergenic potential of pollen is also determined by pollen-associated lipid mediators (PALMs) and low molecular weight compounds.¹⁷ PALMs, such as phytoprostanes, shift dendritic cell-mediated T-cell polarization toward a Th2 response.¹⁸ Also, pollen-derived lipids of the linoleic acid pathway act as chemoattractants for granulocytes.¹⁹ Additionally, low molecular weight compounds and lipid mediators such as PGE₂ and LTB₄ enhance cutaneous reactions and nasal allergic inflammation to common allergens.²⁰

Research determining whether and to what extent rising CO₂ levels influence the potential of ragweed pollen to induce pulmonary allergic disease is lacking. We used a combined approach of an in vivo mouse allergy model, human in vitro tests, and untargeted metabolomics to investigate whether elevated ambient CO₂ levels representative of climate change scenarios lead to enhanced allergenicity of ragweed pollen.

2 | METHODS

2.1 | Growth of ragweed plants in climate chambers

In 2013, ragweed plants were cultivated as previously described.²¹ Plants were grown under ambient (380 ppm) or enriched (700 ppm, based on IPCC scenario RCP4.5)²² CO₂ levels for the whole vegetation period. Aqueous ragweed pollen extracts (control-RWE and CO₂-RWE) were prepared as previously described.²³ Here, concentrations of RWE correspond to the amount of pollen used for the extraction. For more information on plant cultivation and aqueous pollen extract preparation, see online supplement.

2.2 | Murine sensitization model

Experiments were conducted according to federal guidelines for the use and care of laboratory animals and approved by the Government of the District of Upper Bavaria and the Animal Care and Use Committee of the Helmholtz Zentrum München (Approval # 55.2-1-54-2532-156-12).

An adjuvant-free ragweed sensitization protocol was performed as previously described.²⁴ In short, female, 6- to 10-week-old BALB/c

mice received intranasal (i.n.) instillations of control-RWE (10 mg/mL, 10 µL/nostril), CO₂-RWE (10 mg/mL, 10 µL/nostril), or PBS (10 µL/nostril) on 3, 8, or 11 consecutive days. Mice were sacrificed 24 hours after the last instillation (Figure 1A). Blood samples were taken prior to the first instillation and at sacrifice. Measurements of airway hyperresponsiveness, performed after 11 RWE instillations and bronchoalveolar lavage (BAL), occurred as previously described.²⁴ Lung tissue was prepared for histology and FACS analysis.

2.3 | Blood and nasal cell donors

Isolation, culture, and stimulation of primary cells for this study were approved by the ethical committee of the Medical Faculty of the Technical University Munich (ethics statement code: 54/17 S) and the consultative commission of the Augsburg University Medical School (ethics statement code: 2016-7). Blood samples or human nasal epithelial cells (HNEC) from turbinoplasty surgery of healthy non-atopic donors were collected after written informed consent. Atopy status of blood or nasal cell donors was determined by measuring total serum IgE and allergen-specific IgE by serum ImmunoCAP (ThermoFisher, Massachusetts, USA). An overview of the donors, specifying gender, age, total IgE, and RAST classes for the measured aeroallergens is available in Table 1.




FIGURE 1 Pollen of ragweed plants grown under elevated CO₂ levels elicits stronger allergic inflammation *in vivo*. A, Experimental setup. B, BAL cell analysis. C, Airway hyperresponsiveness measured 24 hours after 11x intranasal exposures. n = 5 mice/group; ***P < .01, ***P < .001 vs PBS at same methacholine concentrations. D, Total IgE levels *in vivo* after 11 instillations and (E) *ex vivo*. *In vivo*: n = 5 mice/group; ***P ≤ .001. Representative data of two independent experiments; Mann-Whitney U test, except AHR: ANOVA with post hoc Bonferroni test. *Ex vivo*: n = 7 mice/group; Wilcoxon signed-rank test; *P < .05, dashed line represents unstimulated control

2.4 | Human nasal epithelial cells stimulations

HNEC isolation was performed as recently described.²⁵ For details, see online supplement.

Submerged monolayer cultures of second passage HNECs were seeded in 48-well plates at a density of 2×10^4 cells/well in complete Airway Epithelial Cell Growth Medium (PromoCell, Heidelberg, Germany) and incubated at 37°C, 5% CO₂ for five days. At 80% confluence, the medium was changed to Airway Epithelial Cell Growth Medium without hydrocortisone (PromoCell) and cells were stimulated with control-RWE or CO₂-RWE (0.3 to 2.5 mg/mL). After 24 hours, supernatants were collected and subjected to IL-8, IL-1β, TNF, CCL2, CCL22 (BDOptEIA, BDBioscience Pharmingen, San Diego, CA, USA), IL-33 (R&D Systems, Wiesbaden, Germany), and IL-6 (eBioscience, San Diego, CA, USA) ELISA.

2.5 | Human monocyte-derived dendritic cells stimulations

Dendritic cells (DCs) were isolated from PBMCs as previously described.²⁶ For details, see online supplement. A total of 10⁵ day 5 immature DCs were stimulated with control- or CO₂-RWE (2.5 mg/mL), single pollen-derived compounds (3×10^{-7} M, Table 1), or corresponding compound mixes (3×10^{-7} M, Table 1). For DC stimulation with RWE-conditioned HNEC supernatants, HNEC supernatants of all donors were pooled and supernatants from cells stimulated with the two lowest (0.3 and 0.6 mg/mL) or highest (1.25 and 2.5 mg/mL) concentrations were combined resulting in 0.5 mg/mL (low) and 1.8 mg/mL (high) RWE stimulus concentrations, respectively. Unstimulated DCs correspond to DCs incubated with medium-stimulated HNEC. After 24 hours, supernatants were analyzed by ELISA for IL-10, IL-1β, TNF (BDOptEIA), CCL17 (R&D Systems), and IL-6 (eBioscience) secretion and DC maturation markers were analyzed by flow cytometry. For details, see online supplement.

2.6 | Untargeted metabolome analysis

The metabolome of control-RWE and CO₂-RWE was analyzed using ultra high-resolution mass spectroscopy (ICR-FT/MS) as previously described.²¹ For details, see online supplement.

2.7 | Statistical analysis

In vivo and in vitro data are shown as boxplots indicating minimum, 25% percentile, median, 75% percentile, and maximum, or as mean ± SD. Statistical significance of the in vivo data was determined by Mann-Whitney U test or by two-way ANOVA with post hoc Bonferroni test for lung function analysis. In vitro data were normalized to unstimulated controls, mean ± SD of raw values is available in Tables S2 and S3. Wilcoxon signed-rank test was used to compare two treatment groups of non-normally distributed data. Repeated measures one-way ANOVA with Sidak's post hoc test or Friedman using Dunn's correction was applied for multiple comparisons. Statistical analysis and graph design were performed using GraphPad Prism version 8.4.1. Spider plots for cytokine profiles were created in Excel (2013), using normalized data. Metabolomics data were analyzed using MetaboAnalyst 4.0.²⁷

3 | RESULTS

3.1 | Pollen of ragweed plants grown under elevated CO₂ levels elicit stronger allergic inflammation in vivo

The impact of elevated CO₂ exposure during plant growth on the allergenic potential of ragweed pollen was analyzed in an adjuvant-free mouse model of allergic lung inflammation.²⁴ To assess the kinetics of the allergic response on lung cell infiltration, mice were i.n. instilled on 3, 8, or 11 consecutive days with either PBS, control-RWE, or CO₂-RWE (Figure 1A). Increasing numbers of RWE instillations showed the typical shift from an early, neutrophil-based, to a later, eosinophil-based lung inflammation (Figure 1B). Contrary to control-RWE, in the CO₂-RWE-treated group both neutrophil and eosinophil numbers were significantly elevated after 3 (neutrophils; $P < .05$) and 11 (eosinophils; $P < .05$) i.n. instillations (Figure 1B). Total serum IgE measured 24 hours after 11× i.n. instillations was significantly increased in the CO₂-RWE-treated group compared to the other groups ($P < .001$, Figure 1D), including control-RWE. Ex vivo IgE production of mouse splenic B cells was also elevated upon CO₂-RWE stimulation compared to control-RWE ($P < .05$, Figure 1E). Eleven i.n. instillations of RWE significantly increased airway resistance in both treatment groups compared to PBS control ($P < .01$ for CO₂-RWE and $P < .001$ for RWE, Figure 1C), but no difference was detected between control-RWE and CO₂-RWE.

Gender	No. Donors	Age	Total IgE kU/ml (mean)	Aeroallergens (RAST class; HDM/Cat/Dog/Oat/Grasses/Rye/Penicillium/Cladosporium/Aspergillus/Alternaria/Botrytis/Alder/Birch/Hazel/Ash/Mugwort/Buckhorn)
Female	23	20 - 61	37.00	0/0/0/0/0/0/0/0/0/0/0/0/0/0/0/0/0
Male	13	32 - 67	68.78	0/0/0/0/0/0/0/0/0/0/0/0/0/0/0/0/0

TABLE 1 Overview of cell donors for this study. A total IgE of <100 kU/mL and/or RAST class 0 for common airborne allergens was considered non-atopic

Flow cytometric analysis of lung tissue retrieved 24 hours after the last i.n. instillation revealed a significant increase in eosinophils after 8x and 11x instillations in the CO₂-RWE group, confirming the BAL data (*P* < .05, Figure 2A, top). Furthermore, 8x instillations increased type 2 innate lymphoid cells (ILC2s) in lung tissues of mice treated with control-RWE and CO₂-RWE compared to 3x (*P* < .05) whereby the increase in CO₂-RWE was higher compared to control-RWE, but not significantly. 8x instillations increased Treg numbers in lung tissues of mice treated with control-RWE (*P* < .001 vs 3x and *P* < .05 vs PBS control, Figure 2A, middle). An increased percentage of CD11b⁺DCs in lung tissue was detected in the CO₂-RWE group after 11x instillations compared to the other groups, although significantly only vs PBS (*P* < .05, Figure 2A, bottom).

ILC2s were significantly increased in cervical lymph nodes of mice treated with CO₂-RWE vs control-RWE, although at a later time point compared to lung tissue (11x, *P* < .01 vs PBS, *P* < .05 vs control-RWE, Figure S2). Tregs in the same lymph nodes showed

no significant differences between the treatment groups, whereas higher percentage of DCs was detected after 8x instillations in the CO₂-RWE vs control-RWE group (*P* < .05, Figure S2).

Histopathological analysis of H&E- and PAS-stained lungs after 11x instillations revealed increased perivascular and peribronchiolar inflammatory cell infiltration (ICI) and mucus hypersecretion in control-RWE and CO₂-RWE mice, with CO₂-RWE scoring highest (mucus hypersecretion: *P* < .01 and ICI: *P* < .001 for PBS vs. CO₂-RWE and *P* < .05 for PBS vs. control-RWE in both parameters, Figure 2E,F). Analysis of Th1/Th2 and pro-inflammatory cytokines, as well as chemokines, revealed significant increases of IL-17A (*P* < .001) and IL-17F (*P* < .01) after 11x instillations and CCL22 (*P* < .05) after 3x instillations in CO₂-RWE vs PBS. In control-RWE, only IL-17A after 11x instillations and CCL17 after 3x instillations were significantly increased to PBS (*P* < .05; Figure 3, Figure S2). All other mediators were higher in CO₂-RWE, but did not reach statistical significance, apart from chemokines regulating neutrophil recruitment (CCL3, CCL4,




FIGURE 2 Pollen of ragweed plants grown under elevated CO₂ levels elicit stronger allergic inflammation in vivo. A, Flow cytometric analysis of lung tissue. B-D, Representative PAS-staining of lung sections from mice instilled 11x with pollen extract (B: PBS, C: control-RWE, D: CO₂-RWE). Arrows: inflammatory infiltrate; arrowheads: mucus hypersecretion; scale bar: 100 μm. E and F, Histological scores after 11x instillations. n = 5 mice/group; Mann-Whitney U test; **P* < .05; ***P* < .01; and ****P* < .001 vs PBS, same number of instillations (if applicable). #*P* < .05; ###*P* < .001 vs same experimental group, 3x instillations

and CXCL1), which were slightly higher in control-RWE (Figure 3 and Figure S3).

3.2 | RWEs induce pro-inflammatory responses in human nasal epithelial cells

To analyze the allergenic potential of the different RWEs in a human in vitro system, we stimulated nasal epithelial cells as first port of entry for pollen into the body. Control- and CO₂-RWEs significantly increased IL-8 (control-RWE: $P < .01$ and $P < .001$; CO₂-RWE: $P < .01$), IL-1 β ($P < .01$, $P < .001$) and IL-6 (control-RWE: $P < .01$; CO₂-RWE: $P < .05$) secretion compared to the unstimulated control (Figure S4B,D,E). CCL2 and CCL22 secretion were unchanged (Figure S4A,C). Only TNF release was differentially regulated by CO₂-RWE and control-RWE, being increased by low CO₂-RWE and high control-RWE concentrations ($P < .001$, $P < .01$ vs control-RWE and $P < .01$ vs CO₂-RWE, Figure S4F). IL-33 could not be detected in the supernatants.

3.3 | RWEs induce pro-inflammatory responses in human dendritic cells stimulated with RWE-conditioned epithelial cell supernatants

Because epithelial cells are important modulators of immune responses in the lung,²⁸ we investigated the effect of HNEC supernatants after RWE stimulation downstream of the nasal epithelium. Immature DCs were stimulated with the above characterized HNEC supernatants subsequently pooled in RWE-low and RWE-high, and the cytokine/chemokine profile and maturation markers were analyzed. IL-6 and CCL17 secretion was significantly higher than the baseline across all treatments (Figure 4C, IL-6: $P < .001$; Figure 4E, CCL17: $P < .001$ RWE-high, $P < .0001$ RWE-low, $P < .05$ unstimulated HNEC supernatants). CO₂-RWE-treated HNEC supernatants increased IL-6 ($P < .05$) and CCL17 ($P < .01$) secretion compared to unstimulated HNEC supernatants. IL-10 (Figure 4A) was increased by unstimulated- and CO₂-RWE-treated HNEC supernatants ($P < .01$) and strongly increased by high control-RWE-treated HNEC supernatants ($P < .0001$). TNF (Figure 4B) secretion was higher upon stimulation with CO₂-RWE-treated HNEC




FIGURE 3 Inflammatory mediators in BAL fluid. All mediators were measured 24 hours after 3x, 8x, or 11x i.n. instillations with pollen extract. $n = 5$ mice/group; Mann-Whitney U test; * $P < .05$; ** $P < .01$; and *** $P < .001$ vs PBS, same number of instillations

supernatants than control-RWE ($P < .01$). IL-1 β was only increased by supernatants from unstimulated or CO₂-RWE-stimulated HNECs ($P < .05$, Figure 4D). Overall, the cytokine profile induced by CO₂-RWE-treated HNEC supernatants was strongly pro-inflammatory (Figure 4F). CD80 and CD86 were increased by supernatants from CO₂-RWE-stimulated HNECs ($P < .05$, $P < .0001$, Figure S5B,D), whereas no difference in CD40 and HLA-DR was shown (Figure S5A,E). CD83 (Figure S5C) was elevated by unstimulated HNEC supernatants ($P < .01$) and reduced by CO₂-RWE-stimulated HNEC supernatants compared to unstimulated ($P < .01$).

3.4 | CO₂-RWE induces a more pro-inflammatory response profile in human dendritic cells

Lastly, we analyzed the direct effect of RWEs on dendritic cell cytokine/chemokine secretion and surface marker expression. IL-10 was significantly less secreted by DCs stimulated with CO₂-RWE than control-RWE ($P < .05$, Figure 5A). In contrast, CO₂-RWE significantly increased TNF levels ($P < .05$, Figure 5B). No differences were detected for IL-1 β , IL-6, and CCL17/TARC (Figure 5C-E). Similar to the above described co-culture experiments, CO₂-RWE induced a pro-inflammatory cytokine profile (Figure 5F). Both RWEs induced maturation profiles distinct from the unstimulated control, but similar between the treatments (Figure S6, bottom). Expression of CD86 was increased by both RWEs ($P < .05$), while CD80 was only higher in control-RWE-treated DCs ($P < .05$), (Figure S6, top).

3.5 | Extract analysis reveals candidate substances for enhanced allergenic potential of RWE

Pollen-derived substances act as immune modulators or have pro-inflammatory properties.^{18-20,24} As such, LTB₄, PGE₂, adenosine, and LPS were slightly, although non-significantly, higher in CO₂-RWE (Figure 6A). The content of the major allergen Amb a 1 did not differ between control- and CO₂-RWE (Figure 6A). To gain insight into secondary metabolites present in the RWEs, we used untargeted mass spectroscopy. The metabolite profile of the extracts was distinctly different as revealed by principal component analysis (PCA) (Figure 6C). We observed six candidate substances present only in control-RWE and 13 candidate substances present only in CO₂-RWE (Figure 6D) and chose the ones commercially available or their analogues to stimulate moDCs (Table 2).

3.6 | Pooled, but not single substances are responsible for the cytokine profiles of dendritic cells induced by control- and CO₂-RWE

We used the compounds either separately or in two combinations as present in control- or CO₂-RWE (Table 2) to stimulate DCs and compared the resulting cytokine response to whole RWEs.

Pelargonidin and malvidin enhanced IL-10 secretion ($P < .0001$ and $P < .01$ vs unstimulated), whereas pC4OG decreased IL-10 secretion ($P < .05$ vs. unstimulated) (Figure S7A). Malvidin ($P < .001$) and 9-OTrE ($P < .05$) increased IL-1 β secretion (Figure S7D), and lumicrome decreased IL-6 secretion ($P < .05$ vs. unstimulated, Figure S7C). Compared to a relatively low response to single substances, DCs stimulated with a compound pool mimicking CO₂-RWE secreted less IL-10 ($P < .05$, Figure 6E) and more IL-1 β ($P < .01$, Figure 6H) than with the control-RWE compound mix. TNF and IL-6 secretion did not differ between the two compound mixes.

4 | DISCUSSION

Climate change poses a considerable threat to global health in the foreseeable future.²⁹ Elevated CO₂ levels are part of the driving forces behind our changing climate.³⁰ CO₂ naturally contributes to plant growth, and doubling ambient CO₂ levels have led to increased pollen production of ragweed plants,^{13,14} raising their impact on allergic patients.³¹⁻³³

Here, we investigated whether doubling ambient CO₂ levels to 700 ppm, a still rather conservative IPCC scenario, could also affect the allergenic potential of pollen.

We observed that pollen extracts from plants grown under 700 ppm CO₂ induced a stronger allergic phenotype in a mouse model, characterized by higher serum IgE levels, enhanced lung inflammatory cell recruitment, and mucus hypersecretion, key hallmarks of allergic inflammation.^{34,35} Moreover, we observed moderately increased inflammatory mediators in BAL fluid. In lung and cervical lymph nodes, numbers of dendritic and ILC2 cells, which play a critical role in mounting Th2 responses via IL-33/ST2 signaling under acute and chronic ragweed allergen exposure,³⁶ were increased. Airway hyperresponsiveness was increased by both RWEs compared to PBS control, but no difference was detected between them probably because of the overall moderately increased cytokine response in this study.

To translate our mouse-based results to humans, we used different *in vitro* models to simulate the pollen passage through different immune checkpoints. As a first barrier, the nasal epithelium plays a key role in the allergic sensitization to airborne allergens, responding to pollen stimulation with inflammasome-related cytokines IL-18 and IL-1 β .²⁵ RWEs also activate the inflammasome in keratinocytes by IL-1 β secretion and caspase-1 activation.³⁷ In our study, RWEs induced IL-1 β together with pro-inflammatory cytokines in HNECs, irrespectively of the plant growth conditions. In the absence of IL-12, IL-1 family cytokines have been shown to promote Th2^{38,39} and, in the presence of TGF- β , Th9 differentiation⁴⁰ as well as proliferation of Th2 clones.^{41,42} IL-1 has also been shown to be required for allergen-specific Th2 cell activation and airway inflammation in a mouse model of asthma.⁴³ Indeed, secretion of IL-1 β in our RWE-stimulated HNECs potentially contributes to the Th2 promoting effect downstream of the nasal epithelium.




FIGURE 4 Co-culture of moDCs with supernatants of CO₂-RWE-stimulated HNECs elicits pro-inflammatory cytokine profile. A-E, IL-10, TNF, IL-6, IL-1β, CCL17/TARC secretion, and (F) cytokine profile of moDCs after 24-h stimulation with RWE-conditioned HNEC supernatants (corresponding to 0.5 and 1.8 mg/mL RWE). Dashed line indicates baseline cytokine production of moDCs. n = 35 independent experiments using cells from different donors; A, C, D-E RM one-way ANOVA with Sidak's correction for multiple comparisons, B Friedman's test with Dunn's correction; **P < .01; ***P < .001; ****P < .0001 vs baseline unless indicated otherwise

It is important to note that we used submerged HNEC monolayer cultures instead of air-liquid interface. Although we did not measure tight junctions in our cultures, a characteristic of differentiated epithelia, they have been detected in confluent monolayer cultures of non-atopic donors, similarly to air-liquid interface.^{25,44}

Contrarily to the results obtained by stimulating HNECs with RWEs directly, we report stronger effects of plant treatments upon activation of DCs as downstream effector cells with HNEC supernatants. DCs incubated with supernatants from CO₂-RWE-stimulated HNECs produced more pro-inflammatory cytokines, especially Th2-cell attractant CCL17 and pro-inflammatory IL-6 and TNF, compared to DCs stimulated with control-RWE-treated HNEC supernatants.


Direct stimulation of DCs with pollen extracts clearly demonstrates that CO₂-RWE, which induced allergic airway inflammation in vivo more potently, induced less IL-10 in human DCs in vitro compared to control-RWE. IL-10 is the hallmark cytokine for DC-induced Treg differentiation.⁴⁵ This cytokine was reduced in vitro by CO₂-RWE and by the CO₂ compound mix, consistent with reduced pulmonary Treg numbers upon sensitization with CO₂-RWE in vivo. Our

results are in line with a recent study indicating IL-10 signaling in DCs as essential for efficient tolerance induction.⁴⁶

TNF is another critical factor in allergic sensitization,^{47,48} acting as an adjuvant in house-dust mite allergic sensitization⁴⁹ and exacerbating allergic asthma.⁵⁰ CO₂-RWE induced TNF consistently in our in vitro experiments, but unfortunately we could not detect this cytokine in BAL fluid in vivo. IL-6 secretion, which was upregulated in DCs stimulated with CO₂-RWE-conditioned HNEC supernatants, is also implicated in facilitating Th2 polarization and simultaneous Th1 inhibition by activating NFAC and upregulating SOCS-1 expression in naïve CD4⁺ T cells.⁵¹

Expression of CD80 and CD86 on antigen-presenting cells is important for Th2 differentiation.⁵² Both markers were increased by CO₂-RWE-stimulated HNEC supernatants or by both RWEs by direct DCs stimulations. The role of CD83 on DCs is controversial,⁵³ but seems to be important for CD4⁺ T-cell activation.⁵⁴ CD83 was downregulated by CO₂-RWE-stimulated HNEC supernatants compared to unstimulated. Combined with the expression of CD80/CD86, our findings emphasize the importance of the mode of DC stimulation, either by RWE directly or indirectly via HNEC supernatants.

FIGURE 5 Pollen of ragweed plants grown under elevated CO₂ levels induce pro-inflammatory cytokine profile in moDCs. A-E, IL-10, TNF, IL-6, IL-1 β , and CCL17/TARC were measured in cell culture supernatants after 24-h stimulation with 2.5 mg/mL control- or CO₂-RWE, and the results were summarized in a profile (F). Dashed line indicates unstimulated control. n = 24 independent experiments using cells from different donors; A, RM one-way ANOVA with Sidak's test for multiple comparisons. B-E, Friedman's test with Dunn's correction for multiple comparisons; **P* < .05 and ***P* < .01 comparison between treatment groups



In addition to activating epithelial-DC cross-talk, RWE acts directly on B cells, increasing IgE secretion under Th2-mimicking conditions.⁵⁵ We demonstrate that CO₂-RWE increased the IgE response ex vivo as well as in vivo compared to control-RWE. Thus, RWEs appear to act on several levels of the immune response contributing to the clinical phenotype of ragweed allergy, that is, DC-mediated sensitization and B cell-mediated IgE production, which are both enhanced under exposure to CO₂-RWE.

To identify one or more substances responsible for the observed CO₂-RWE-induced increased allergic response, we first analyzed PALMs, known pollen-derived immune modulators.^{18,20} Pollen-derived adenosine appears to be protective during allergic sensitization by inducing regulatory responses in dendritic-primed T cells in vitro,²⁶ whereas it mediates exacerbation of allergic lung inflammation in vivo.²⁴ Slightly elevated PALMS and adenosine in CO₂-RWE can only partly explain the increased inflammatory response following CO₂-RWE exposure. Therefore, we broaden the analysis investigating the pollen metabolome. Here, we found a plethora of secondary plant metabolites differentially regulated by growth conditions. Metabolites which were exclusively present in CO₂-RWE (malvidin, pelargonidin, catalposide, and 9-oxo-OTrE) or in control-RWE (lumichrome, Q3OS and *p*-Coumaryl-alcohol-4-O-glucoside), exhibiting mostly anti-inflammatory/tolerogenic characteristics⁵⁶⁻⁶³ were employed for in vitro stimulations of DCs. Pelargonidin and malvidin alone were anti-inflammatory, while the opposite was seen for *p*-Coumaryl-alcohol-4-O-glucoside, and the other substances had almost no effect. We showed synergistic effects of the compound mixes, which induced a cytokine profile comparable to whole pollen extracts. Indeed, substances with known anti-inflammatory properties exhibited

pro-inflammatory properties when applied as a mix. Metabolomic screening was performed in a non-targeted, semi-quantitative manner, providing a global overview of the pollen metabolome without delivering absolute quantities of the significantly modulated compounds. The substances were annotated by their exact mass and elemental composition and chosen according to their immunological properties and commercial availability in case of multi-annotation. Nevertheless, we can conclude that more than a single adjuvant substance in the allergen matrix is needed to transmit an integrated signal via DCs to downstream effectors of the adaptive immune response, that is, T and B cells.

In summary, we showed that CO₂-RWE elicits a stronger allergic response compared to control-RWE and that allergenicity cannot be confined to a single factor, but rather stems from the interplay of different mediators. Given that IPCC reports predict a rise in atmospheric CO₂ from currently around 400 ppm to a range of 730-1020 ppm expected by the year 2100,³⁰ it should be noted that the impact of most pessimistic IPCC scenarios (eg, 1000 ppm CO₂) might further enhance not only pollen biomass, but also pollen allergenicity, which will most probably contribute to an increase of allergic responses to ragweed in the population. Together with our previous research on effects of climate change scenarios on pollen,^{16,64} we demonstrate that climate change affects plants and pollen allergenicity, emphasizing the importance of viewing climate change as an existential threat to our health.

ACKNOWLEDGMENTS

The authors wish to thank the animal caretakers of the Helmholtz Center Munich and Johanna Grosch, Benjamin Schnautz, Selina Eisenbart, and Stephanie Lukas for technical assistance. We




FIGURE 6 Metabolome analysis of RWEs reveals differentially expressed clusters of substances. A, PALMs, LPS, adenosine, and Amb a 1 measured in extracts of single plants (n = 10). B, Heatmap of substances present in RWE are clustered using Euclidean distance measure and Ward's linkage-clustering algorithm. C, Principal component analysis (PCA). D, Univariate volcano plot analysis of all metabolites. n = 3 control-RWEs and n = 4 CO₂-RWEs for metabolome analysis. E-H, Cytokines measured in DC supernatants 24 hours after stimulation with compound mixes (concentration 3×10^{-7} M, Table 1). n = 27 independent experiments using cells from different donors; Wilcoxon signed-rank test; *P < .05 and **P < .01 comparison between treatment groups

TABLE 2 Putative substances identified in CO₂-RWE and control-RWE, their compound class, and corresponding compound mix





Compound	Compound class	Compound Mix	Company
Pelargonidin (Pel)	Anthocyanidins	CO ₂ -Mix	Sigma-Aldrich (Taufkirchen, Germany)
Malvidin (Mal)	Anthocyanidins		
Catalposide (Cat)	Terpenoids		
9-Oxo-OTrE (9-OTrE)	α-Linolenic acid metabolites		
p-Coumaryl alcohol 4-O-glucoside (pC4OG)	Phenylpropanoids	Control mix	
Lumichrome (Lumi)	Riboflavins		
Quercetin-3-O-sophoroside (Q3OS)	Flavones and flavonols		F. Ferreira and L. Aglas, University of Salzburg, Austria

gratefully acknowledge Hans Lang for his excellent support in CO₂ fumigation in the phytotron chambers.

CONFLICTS OF INTEREST

Ms Rauer, Dr Gilles, Dr Wimmer, Dr Frank, Dr Mueller, Dr Musiol, Dr Vafadari, Dr Aglas, Prof. Dr Ferreira, Prof. Dr Schmitt-Kopplin, Prof. Dr Durner, Dr Winkler, Dr Ernst, Prof. Dr Behrendt, Prof. Dr Schmidt-Weber, Prof. Dr Traidl-Hoffmann, and Prof. Dr Alessandrini have nothing to disclose.

ORCID

Stefanie Gilles  <https://orcid.org/0000-0002-5159-2558>
 Stephanie Musiol  <https://orcid.org/0000-0001-8356-4343>
 Fatima Ferreira  <https://orcid.org/0000-0003-0989-2335>
 Carsten B. Schmidt-Weber  <https://orcid.org/0000-0002-3203-8084>
 Claudia Traidl-Hoffmann  <https://orcid.org/0000-0001-5085-5179>
 Francesca Alessandrini  <https://orcid.org/0000-0002-9854-8968>

REFERENCES

- Arbes SJ Jr, Gergen PJ, Elliott L, Zeldin DC. Prevalences of positive skin test responses to 10 common allergens in the US population: results from the third National Health and Nutrition Examination Survey. *J Allergy Clin Immunol.* 2005;116(2):377-383.
- Gergen PJ, Arbes SJ Jr, Calatroni A, Mitchell HE, Zeldin DC. Total IgE levels and asthma prevalence in the US population: results from the National Health and Nutrition Examination Survey 2005–2006. *J Allergy Clin Immunol.* 2009;124(3):447-453.
- Buters J, Alberterst B, Nawrath S, et al. *Ambrosia artemisiifolia* (ragweed) in Germany—current presence, allergological relevance and containment procedures. *Allergo J Int.* 2015;24:108-120.
- Chen KW, Marusciac L, Tamas PT, Valenta R, Panaitescu C. Ragweed pollen allergy: burden, characteristics, and management of an imported allergen source in Europe. *Int Arch Allergy Immunol.* 2018;176(3–4):163-180.
- Lake IR, Jones NR, Agnew M, et al. climate change and future pollen allergy in Europe. *Environ Health Perspect.* 2017;125(3):385-391.
- DellaValle CT, Triche EW, Leaderer BP, Bell ML. Effects of ambient pollen concentrations on frequency and severity of asthma symptoms among asthmatic children. *Epidemiology.* 2012;23(1):55-63.
- Heuson C, Traidl-Hoffmann C. The significance of climate and environment protection for health under special consideration of skin barrier damages and allergic sequelae. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitschutz.* 2018;61(6):684-696.
- Storkey J, Stratonovitch P, Chapman DS, Vidotto F, Semenov MA. A process-based approach to predicting the effect of climate change on the distribution of an invasive allergenic plant in Europe. *PLoS ONE* 2014;9(2):e88156.
- Rasmussen K, Thyrring J, Muscarella R, Borchsenius F. Climate-change-induced range shifts of three allergenic ragweeds (*Ambrosia* L.) in Europe and their potential impact on human health. *PeerJ.* 2017;5:e3104.
- Cunze S, Leiblein MC, Tackenberg O. Range expansion of *Ambrosia artemisiifolia* in Europe is promoted by climate change. *ISRN Ecol.* 2013;2013:1-9.
- Ziska LH, Makra L, Harry SK, et al. Temperature-related changes in airborne allergenic pollen abundance and seasonality across the northern hemisphere: a retrospective data analysis. *Lancet Planet Health.* 2019;3(3):e124-e131.
- Ziska L, Knowlton K, Rogers C, et al. Recent warming by latitude associated with increased length of ragweed pollen season in central North America. *Proc Natl Acad Sci U S A.* 2011;108(10):4248-4251.
- Wayne P, Foster S, Connolly J, Bazzaz F, Epstein P. Production of allergenic pollen by ragweed (*Ambrosia artemisiifolia* L.) is increased in CO₂-enriched atmospheres. *Ann Allergy Asthma Immunol.* 2002;88(3):279-282.
- Rogers CA, Wayne PM, Macklin EA, et al. Interaction of the onset of spring and elevated atmospheric CO₂ on ragweed (*Ambrosia artemisiifolia* L.) pollen production. *Environ Health Perspect.* 2006;114(6):865-869.
- Singer BD, Ziska LH, Frenz DA, Gebhard DE, Straka JG. Increasing Amb a 1 content in common ragweed (*Ambrosia artemisiifolia*) pollen as a function of rising atmospheric CO₂ concentration. *Funct Plant Biol* 2005;32(7):667-670.

16. El Kelish A, Zhao F, Heller W, et al. Ragweed (*Ambrosia artemisiifolia*) pollen allergenicity: SuperSAGE transcriptomic analysis upon elevated CO₂ and drought stress. *BMC Plant Biol.* 2014;14:176-191.
17. Gilles S, Akdis C, Lauener R, et al. The role of environmental factors in allergy: a critical reappraisal. *Exp Dermatol.* 2018;27(11):1193-1200.
18. Traidl-Hoffmann C, Mariani V, Hochrein H, et al. Pollen-associated phytoprostanes inhibit dendritic cell interleukin-12 production and augment T helper type 2 cell polarization. *J Exp Med.* 2005;201(4):627-636.
19. Traidl-Hoffmann C, Kasche A, Jakob T, et al. Lipid mediators from pollen act as chemoattractants and activators of polymorphonuclear granulocytes. *J Allergy Clin Immunol.* 2002;109(5):831-838.
20. Gilles-Stein S, Beck I, Chaker A, et al. Pollen derived low molecular compounds enhance the human allergen specific immune response in vivo. *Clin Exp Allergy.* 2016;46(10):1355-1365.
21. Kanter U, Heller W, Durner J, et al. Molecular and immunological characterization of ragweed (*Ambrosia artemisiifolia* L.) pollen after exposure of the plants to elevated ozone over a whole growing season. *PLoS ONE.* 2013;8(4):e61518.
22. van Vuuren DP, Edmonds J, Kainuma M, et al. The representative concentration pathways: an overview. *Clim Change.* 2011;109(1):5-31.
23. Buters JTM, Kasche A, Weichenmeier I, et al. Year-to-year variation in release of Bet v 1 allergen from birch pollen: evidence for geographical differences between West and South Germany. *Int Arch Allergy Immunol.* 2008;145(2):122-130.
24. Wimmer M, Alessandrini F, Gilles S, et al. Pollen-derived adenosine is a necessary cofactor for ragweed allergy. *Allergy.* 2015;70(8):944-954.
25. Bergougnan C, Dittlein DC, Hümmer E, et al. Physical and immunological barrier of human primary nasal epithelial cells from non-allergic and allergic donors. *World Allergy Organ J.* 2020;13(3):100109.
26. Gilles S, Fekete A, Zhang X, et al. Pollen metabolome analysis reveals adenosine as a major regulator of dendritic cell-primed T(H) cell responses. *J Allergy Clin Immunol.* 2011;127(2):454-461 e451-459.
27. Chong J, Wishart DS, Xia J. Using MetaboAnalyst 4.0 for comprehensive and integrative metabolomics data analysis. *Curr Protoc Bioinformatics.* 2019;68(1):e86.
28. Weitnauer M, Mijosek V, Dalpke AH. Control of local immunity by airway epithelial cells. *Mucosal Immunol.* 2016;9(2):287-298.
29. Watts N, Amann M, Arnell N, et al. The 2019 report of The Lancet Countdown on health and climate change: ensuring that the health of a child born today is not defined by a changing climate. *Lancet.* 2019;394(10211):1836-1878.
30. Meehl GA, Stocker TF, Collins WD, et al. Global climate projections. In: Solomon S, Qin D, Manning M, et al., eds. *Climate Change 2007: The Physical Science Basis Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge, UK: Cambridge University Press; 2007.
31. Jones NR, Agnew M, Banic I, et al. Ragweed pollen and allergic symptoms in children: results from a three-year longitudinal study. *Sci Total Environ.* 2019;683:240-248.
32. Ariano R, Berra D, Chiodini E, et al. Ragweed allergy: Pollen count and sensitization and allergy prevalence in two Italian allergy centers. *Allergy Rhinol.* 2015;6(3):177-183.
33. Schmidt CW. Pollen overload: seasonal allergies in a changing climate. *Environ Health Perspect.* 2016;124(4):A70-A75.
34. Alessandrini F, Schulz H, Takenaka S, et al. Effects of ultrafine carbon particle inhalation on allergic inflammation of the lung. *J Allergy Clin Immunol.* 2006;117(4):824-830.
35. Yu QL, Chen Z. Establishment of different experimental asthma models in mice. *Exp Ther Med.* 2018;15(3):2492-2498.
36. Akasaki S, Matsushita K, Kato Y, et al. Murine allergic rhinitis and nasal Th2 activation are mediated via TSLP- and IL-33-signaling pathways. *Int Immunol.* 2016;28(2):65-76.
37. Dittlein DC, Gilles-Stein S, Hiller J, et al. Pollen and UV-B radiation strongly affect the inflammasome response in human primary keratinocytes. *Exp Dermatol.* 2016;25(12):991-993.
38. Caucheteux SM, Hu-Li J, Guo L, et al. IL-1 β enhances inflammatory TH2 differentiation. *J Allergy Clin Immunol.* 2016;138(3):898-901. e894.
39. Xu D, Trajkovic V, Hunter D, et al. IL-18 induces the differentiation of Th1 or Th2 cells depending upon cytokine milieu and genetic background. *Eur J Immunol.* 2000;30(11):3147-3156.
40. Uyttenhove C, Brombacher F, Van Snick J. TGF-beta interactions with IL-1 family members trigger IL-4-independent IL-9 production by mouse CD4(+) T cells. *Eur J Immunol.* 2010;40(8):2230-2235.
41. Lichtman AH, Chin J, Schmidt JA, Abbas AK. Role of interleukin 1 in the activation of T lymphocytes. *Proc Natl Acad Sci U S A.* 1988;85(24):9699-9703.
42. Taylor-Robinson AW, Phillips RS. Expression of the IL-1 receptor discriminates Th2 from Th1 cloned CD4+ T cells specific for *Plasmodium chabaudi*. *Immunology.* 1994;81(2):216-221.
43. Nakae S, Komiyama Y, Yokoyama H, et al. IL-1 is required for allergen-specific Th2 cell activation and the development of airway hypersensitivity response. *Int Immunol.* 2003;15(4):483-490.
44. Blume C, Swindle EJ, Gilles S, Traidl-Hoffmann C, Davies DE. Low molecular weight components of pollen alter bronchial epithelial barrier functions. *Tissue Barriers.* 2015;3(3):e1062316.
45. Akdis CA, Akdis M. Mechanisms of immune tolerance to allergens: role of IL-10 and Tregs. *J Clin Invest.* 2014;124(11):4678-4680.
46. Dolch A, Kunz S, Dorn B, et al. IL-10 signaling in dendritic cells is required for tolerance induction in a murine model of allergic airway inflammation. *Eur J Immunol.* 2019;49(2):302-312.
47. Bachus H, Kaur K, Papillion AM, et al. Impaired tumor-necrosis-factor-alpha-driven dendritic cell activation limits lipopolysaccharide-induced protection from allergic inflammation in infants. *Immunity.* 2019;50(1):225-240.e224.
48. Choi J-P, Kim Y-S, Kim OY, et al. TNF-alpha is a key mediator in the development of Th2 cell response to inhaled allergens induced by a viral PAMP double-stranded RNA. *Allergy.* 2012;67(9):1138-1148.
49. Lambert AL, Selgrade MK, Winsett DW, Gilmour MI. TNF-alpha enhanced allergic sensitization to house dust mite in brown Norway rats. *Exp Lung Res.* 2001;27(7):617-635.
50. Kips JC. Cytokines in asthma. *Eur Respir J Suppl.* 2001;34:24s-33s.
51. Diehl S, Rincon M. The two faces of IL-6 on Th1/Th2 differentiation. *Mol Immunol.* 2002;39(9):531-536.
52. Li J-G, Du Y-M, Yan Z-D, et al. CD80 and CD86 knockdown in dendritic cells regulates Th1/Th2 cytokine production in asthmatic mice. *Exp Ther Med.* 2016;11(3):878-884.
53. Prazma CM, Tedder TF. Dendritic cell CD83: a therapeutic target or innocent bystander? *Immunol Lett.* 2008;115(1):1-8.
54. Aerts-Toegaert C, Heirman C, Tuyaerts S, et al. CD83 expression on dendritic cells and T cells: correlation with effective immune responses. *Eur J Immunol.* 2007;37(3):686-695.
55. Oeder S, Alessandrini F, Wirz OF, et al. Pollen-derived nonallergenic substances enhance Th2-induced IgE production in B cells. *Allergy.* 2015;70(11):1450-1460.
56. Bognar E, Sarszegi Z, Szabo A, et al. Antioxidant and anti-inflammatory effects in RAW264.7 macrophages of malvidin, a major red wine polyphenol. *PLoS ONE.* 2013;8(6):e65355.
57. Hamalainen M, Nieminen R, Vuorela P, Heinonen M, Moilanen E. Anti-inflammatory effects of flavonoids: genistein, kaempferol, quercetin, and daidzein inhibit STAT-1 and NF-kappaB activations, whereas flavone, isorhamnetin, naringenin, and pelargonidin inhibit only NF-kappaB activation along with their inhibitory effect

- on iNOS expression and NO production in activated macrophages. *Mediators Inflamm.* 2007;2007:45673.
58. An SJ, Pae HO, Oh GS, et al. Inhibition of TNF-alpha, IL-1beta, and IL-6 productions and NF-kappa B activation in lipopolysaccharide-activated RAW 264.7 macrophages by catalposide, an iridoid glycoside isolated from *Catalpa ovata* G. Don (Bignoniaceae). *Int Immunopharmacol.* 2002;2(8):1173-1181.
59. Prost I, Dhondt S, Rothe G, et al. Evaluation of the antimicrobial activities of plant oxylipins supports their involvement in defense against pathogens. *Plant Physiol* 2005;139(4):1902-1913.
60. Schramm M, Wiegmann K, Schramm S, et al. Riboflavin (vitamin B2) deficiency impairs NADPH oxidase 2 (Nox2) priming and defense against *Listeria monocytogenes*. *Eur J Immunol.* 2014;44(3):728-741.
61. Jansen F, Gillessen B, Mueller F, Commandeur U, Fischer R, Kreuzaler F. Metabolic engineering for p-coumaroyl alcohol production in *Escherichia coli* by introducing an artificial phenylpropanoid pathway. *Biotechnol Appl Biochem.* 2014;61(6):646-654.
62. Tsuji R, Ikado K, Fujiwara D. Modulation of innate immunity by lignin-carbohydrate, a novel TLR4 ligand, results in augmentation of mucosal IgA and systemic IgG production. *Int J Mol Sci.* 2017;19(1):64-78.
63. Seutter von Loetzen C, Hoffmann T, Hartl M, et al. Secret of the major birch pollen allergen Bet v 1: identification of the physiological ligand. *Biochem J.* 2014;457(3):379-390.
64. Beck I, Jochner S, Gilles S, et al. High environmental ozone levels lead to enhanced allergenicity of birch pollen. *PLoS ONE.* 2013;8(11):e80147.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Rauer D, Gilles S, Wimmer M, et al. Ragweed plants grown under elevated CO₂ levels produce pollen which elicit stronger allergic lung inflammation. *Allergy.* 2020;00:1-13. <https://doi.org/10.1111/all.14618>