

Strict convexity of the free energy for non-convex gradient models at moderate β

Codina Cotar ^{*†‡}, Jean-Dominique Deuschel ^{*§}, and Stefan Müller ^{*¶}

February 2, 2008

Abstract

We consider a gradient interface model on the lattice with interaction potential which is a non-convex perturbation of a convex potential. We show using a one-step multiple scale analysis the strict convexity of the surface tension at high temperature. This is an extension of Funaki and Spohn's result [10], where the strict convexity of potential was crucial in their proof.

AMS 2000 Subject Classification. 60K35, 82B24, 35J15

Key words and phrases. effective non-convex gradient interface models, surface tension, strict convexity, Helffer-Sjöstrand representation

1 Introduction

We consider an an effective model with gradient interaction. The model describes a phase separation in \mathbb{R}^{d+1} , eg. between the liquid and vapor phase. For simplicity we consider a discrete basis $\Lambda_M \subset \mathbb{Z}^d$, and continuous height variables

$$x \in \Lambda_M \longrightarrow \phi(x) \in \mathbb{R}.$$

This model ignores overhangs like in Ising models, but gives a good approximation in the vicinity of the phase separation. The distribution of the interface is given in terms of its Gibbs distribution with nearest neighbor interactions of gradient type, that is, the interaction between two neighboring sites x, y depends only on the discrete gradient, $\nabla\phi(x, y) = \phi(y) - \phi(x)$. More precisely, the Hamiltonian is of the form

$$H_M(\phi) = \sum_{x, y \in \Lambda_{M+1}, |x-y|=1} V(\phi(y) - \phi(x)) \quad (1.1)$$

*Corresponding Author

†Supported by the DFG-Forschergruppe 718 'Analysis and stochastics in complex physical systems'

‡TU Berlin - Fakultät II, Institut für Mathematik, Strasse des 17. Juni 136, D-10623 Berlin, Germany. E-mail: cotar@math.tu-berlin.de

§TU Berlin - Fakultät II Institut für Mathematik Strasse des 17. Juni 136 D-10623 Berlin, Germany. E-mail: deuschel@math.tu-berlin.de

¶Max Planck Institute for Mathematics in the Sciences Inselstrasse 22-26 D-04103 Leipzig, Germany. E-mail: sm@mis.mpg.de

where $V \in C^2(\mathbb{R})$ is a function with quadratic growth at infinity:

$$V(\eta) \geq A|\eta|^2 - B, \quad \eta \in \mathbb{R} \quad (1.2)$$

for some $A > 0, B \in \mathbb{R}$.

For a given boundary condition $\psi \in \mathbb{R}^{\partial\Lambda_M}$, where $\partial\Lambda_M = \Lambda_{M+1} \setminus \Lambda_M$, the (finite) Gibbs distribution on $\mathbb{R}^{\Lambda_{M+1}}$ at inverse temperature $\beta > 0$ is given by

$$\mu_{V_M, \psi}^\beta(d\phi) \equiv \frac{1}{Z_{M, \psi}^\beta} \exp(-\beta H_M(\phi)) \prod_{x \in \Lambda_M} d\phi(x) \prod_{x \in \partial\Lambda_M} \delta_{\psi(x)}(d\phi(x)).$$

Here $Z_{M, \psi}^\beta$ is a normalizing constant given by

$$Z_{M, \psi}^\beta = \int_{\mathbb{R}^{\Lambda_{M+1}}} \exp(-\beta H_M(\phi)) \prod_{x \in \Lambda_M} d\phi(x) \prod_{x \in \partial\Lambda_M} \delta_{\psi(x)}(d\phi(x)).$$

One is particularly interested in tilted boundary conditions

$$\psi_u(x) = \langle x, u \rangle = \sum_{i=1}^d x_i u_i$$

for some given 'tilt' $u \in \mathbb{R}^d$. This corresponds to an interface in \mathbb{R}^{d+1} which stays normal to the vector $n_u = (u, -1) \in \mathbb{R}^{d+1}$.

An object of basic relevance in this context is the surface tension or free energy defined by the limit

$$\sigma(u) = \lim_{M \rightarrow \infty} -\frac{1}{\beta} \log Z_{M, \psi_u}^\beta. \quad (1.3)$$

The existence of the above limit follows from a standard sub-additivity argument. In fact the surface tension can also be defined in terms of the partition function on the torus, see below and [10]. In case of *strictly* convex potential V with

$$c_1 \leq V'' \leq c_2 \quad (1.4)$$

where $0 < c_1 \leq c_2 < \infty$, Funaki and Spohn showed in [10] that σ is strictly convex.

The simplest strictly convex potential is the quadratic one with $V(\eta) = |\eta|^2$, which corresponds to a Gaussian model, also called gradient free field or harmonic crystal. Models with non quadratic potentials V are sometimes called anharmonic crystals.

The strict convexity of the surface tension σ plays a crucial role in the derivation of the hydrodynamical limit of the Landau-Ginsburg model in [10].

Under the condition (1.4), a large deviation principle for the rescaled profile with rate function given in terms of the integrated surface tension has been derived in [7]. Here also the strict convexity of σ is very important. Both papers [10] and [7] use very explicitly the condition (1.4) in their proof. In particular they rely on the Brascamp Lieb inequality and on the random walk representation of Helffer and Sjöstrand, which requires a strictly convex potential V .

The objective of our work is to prove strict convexity of σ also for some non convex potential. One cannot expect strict convexity for any non convex V , see

below. Our result is perturbative at high temperature (small β), and shows strict convexity of $\sigma(u)$ at every $u \in \mathbb{R}$ for potentials V of the form

$$V(\eta) = V_0(\eta) + g_0(\eta)$$

where V_0 satisfies (1.4) and $g_0 \in C^2(\mathbb{R})$ has a negative bounded second derivative such that $\sqrt{\beta} \cdot \|g_0''\|_{L^1(\mathbb{R})}$ is small enough.

Our proof is based on the scale decomposition of the free field as the sum of two independent free fields ϕ_1 and ϕ_2 , where we choose the variance of ϕ_1 small enough to match the non-convexity of g . This particular type of scale decomposition was used earlier by Haru Pinson in [11], who also suggested to us the use of this approach. The partition function Z_{N,ψ_u}^β can be then expressed in terms of a double integral, with respect to both ϕ_1 and ϕ_2 . We fix ϕ_2 and perform first the integration with respect to ϕ_1 . This yields a new induced Hamiltonian, which is a function of the remaining variable ϕ_2 . The main point is that our choice of the variance of ϕ_1 and smallness of β allow us to show convexity in ϕ_2 of the induced Hamiltonian. Of course this Hamiltonian is no longer of the simple form (1.1), in particular we lose the locality of the interaction. However an extension of the technique introduced in [7] shows strict convexity of σ . The idea behind the proof is that one can gain convexity via integration. This procedure is called "one step decomposition", since we perform only one integration. Of course this procedure could be iterated which would allow to lower the temperature. However for general non convex g we do not expect that this procedure works at low temperature for every tilt u .

At low temperature an approach in the spirit of [5], [4] looks more promising [1].

Finally note that, due to the gradient interaction, the Hamiltonian has a continuous symmetry. In particular this implies that no infinite Gibbs state exists for the lower lattice dimensions, $d = 1, 2$ where the field "delocalizes" as $M \rightarrow \infty$, c.f. [8]. On the other hand, it is very natural in this setting to consider the *gradient* Gibbs distributions, that is the image of $\mu_{V_M,\psi}$ under the gradient operation $\phi \in \mathbb{R}^{\mathbb{Z}^d} \rightarrow \nabla\phi$. It is easy to verify that this distribution depends only on $\nabla\psi$, the gradient of the boundary condition, in fact one can also introduce gradient Gibbs distributions in terms of conditional distributions satisfying DLR conditions, c.f. [10]. Using the quadratic bound (1.1), one can easily see that the corresponding measures are tight. In particular for each tilt $u \in \mathbb{R}^d$ one can construct a translation invariant gradient Gibbs state $\tilde{\mu}_u$ on \mathbb{Z}^d with mean u :

$$\mathbb{E}_{\tilde{\mu}_u}[\phi(y) - \phi(x)] = \langle y - x, u \rangle .$$

Under (1.4), Funaki and Spohn proved the existence and unicity of extremal, ie. ergodic, gradient Gibbs state, for each tilt $u \in \mathbb{R}$. In the case of non convex V , unicity of the ergodic states can be violated, even at $u = 0$ tilt, c.f. [2]. However in this situation, the surface tension is not strictly convex at $u = 0$.

2 Main result and outline of the proof

We study the convexity properties of the free energy (as a function of the tilt u) for non-convex gradient models on a lattice. Using the results of [10], we

work on the torus, instead of the box Λ_M , see Remark 2.4 below. Thus, let $\mathbb{T}_M^d = (\mathbb{Z}/M\mathbb{Z})^d = \mathbb{Z}^d \bmod (M)$ be the lattice torus in \mathbb{Z}^d , let $u \in \mathbb{R}^d$ and let $\beta > 0$. For a function $\phi : \mathbb{T}_M^d \rightarrow \mathbb{R}$, we consider the discrete derivative

$$\nabla_i \phi(x) = \phi(x + e_i) - \phi(x) \quad (2.5)$$

and the Hamiltonian

$$H(u, \phi) = \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d [V_0^i(\nabla_i \phi(x) + u_i) + g_0^i(\nabla_i \phi(x) + u_i)], \quad (2.6)$$

where V_0^i is convex and g_0^i is non-convex (see (2.11) below). We consider the partition function

$$Z_M^\beta(u) = \int_X e^{-\beta H(u, \phi)} m_M(d\phi), \quad (2.7)$$

where

$$X = \{\phi : \mathbb{T}_M^d \rightarrow \mathbb{R} : \phi(0) = 0\} \quad (2.8)$$

and

$$m_M(d\phi) = \prod_{x \in \mathbb{T}_M^d \setminus \{0\}} d\phi(x) \delta_0(d\phi(0)), \quad (2.9)$$

and the free energy

$$f_M^\beta(u) = -\frac{1}{\beta} \log Z_M^\beta(u). \quad (2.10)$$

We will prove

Theorem 2.1 *Suppose that V_0^i and g_0^i are C^2 functions on \mathbb{R} and that there exist constants C_0, C_1, C_2 and*

$$0 < C_1 \leq (V_0^i)'' \leq C_2, \quad -C_0 \leq (g_0^i)'' \leq 0. \quad (2.11)$$

Set

$$\bar{C} = \max\left(\frac{C_0}{C_1}, \frac{C_2}{C_1} - 1, 1\right). \quad (2.12)$$

If $(g_0^i)'' \in L^1(\mathbb{R})$ and for $i \in \{1, 2, \dots, d\}$

$$\frac{4}{\pi} (12d\bar{C})^{1/2} \sqrt{\beta C_1} \frac{1}{C_1} \|(g_0^i)''\|_{L^1(\mathbb{R})} \leq \frac{1}{2}, \quad (2.13)$$

then

$$(D^2 f_M^\beta)(u) \geq \frac{C_1}{2} |\mathbb{T}_M^d| \text{Id}, \quad \forall u \in \mathbb{R}^d, \quad (2.14)$$

where $|\mathbb{T}_M^d| = M^d$ denotes the number of points in \mathbb{T}_M^d . In other words, the free energy per particle is uniformly convex, uniformly in M .

Remark 2.2 The main point is that the convexity estimate (2.14) holds uniformly in the size M of the torus. Indeed a direct calculation of $D^2 f_M^1$ yields at u

$$D^2 f_M^1(u) = \langle D_u^2 H(u, \cdot) \rangle_H - \text{var}_H D_u H(u, \cdot), \quad (2.15)$$

where

$$\langle f \rangle_H = \frac{\int_X f(\phi) e^{-H(u, \phi)} m_M(d\phi)}{\int_X e^{-H(u, \phi)} m_M(d\phi)} \quad (2.16)$$

and

$$\text{var}_H f = \langle (f - \langle f \rangle_H)^2 \rangle_H. \quad (2.17)$$

Now one might expect that a condition like (2.13) implies that $\langle (D_u^2 H(u, \cdot)) \rangle_H \geq cC_1 |\mathbb{T}_M^d| \text{Id}$ (see Lemma 4.1 below). The problem is that naively the variance term scales like $|\mathbb{T}_M^d|^2$ since $D_u H$ is a sum of $d|\mathbb{T}_M^d|$ terms. To get a better estimate, one has to show that in a suitable sense, the terms

$$\text{cov}_H (D_u(V_0 + g_0)(u + \nabla_i \phi(x)), D_u(V_0 + g_0)(u + \nabla_j \phi(y))) \quad (2.18)$$

decay if $|x - y|$ is large. If H is not convex such a decay of correlations is, presently, only proved for the class of potentials studied in [6]. As discussed above, the Helffer-Sjöstrand estimates do not apply directly. The main idea is to rewrite $Z_M^\beta(u)$ as an iterated integral in such a way that each integration involves a convex hamiltonian to which the Helffer-Sjöstrand theory can be applied (see (2.42) below).

Remark 2.3 Instead of $\|g_0''\|_{L^1(\mathbb{R})}$ one can also use bounds on lower order derivatives. More precisely, condition (2.13) can, for example be replaced by

$$\frac{50}{\sqrt{2\pi}} d\bar{C}(\beta C_1)^{3/4} \frac{1}{C_1} \|g_0'\|_{L^2(\mathbb{R})} \leq \frac{1}{2} \quad (2.19)$$

(see Remark 4.2 below). In view of the estimate

$$\int_{\mathbb{R}} (g_0')^2(s) ds = \int_{\mathbb{R}} g_0(s) g_0''(s) ds \leq C_0 \|g_0\|_{L^1(\mathbb{R})}, \quad (2.20)$$

we can see that (2.13) can be replaced by

$$cd^2 \bar{C}^3 (\beta C_1)^{3/2} \frac{1}{C_1} \|g_0\|_{L^1(\mathbb{R})} \leq \frac{1}{4} \quad (2.21)$$

with $c = \frac{2500}{2\pi}$.

Remark 2.4 Note that the surface tensions defined in (1.3) and (2.10) coincide (see, for example, [10]). Because of this, we will work from now on with the definition of the surface tension on a torus, as it is easier to use.

Outline of the proof for Theorem 2.1

Step 1: Scaling argument

A simple scaling argument shows that it suffices to prove the result for

$$\beta = 1, C_1 = 1. \quad (2.22)$$

Indeed, suppose that the result is true for $\beta = 1$ and $C_1 = 1$. Given β , V_0^i and g_0^i which satisfy (2.11) and (2.13), we define

$$\tilde{V}_0^i(s) = \beta V_0^i\left(\frac{s}{\sqrt{\beta C_1}}\right), \quad \tilde{g}_0^i(s) = \beta g_0^i\left(\frac{s}{\sqrt{\beta C_1}}\right). \quad (2.23)$$

Then

$$\begin{aligned} 1 \leq (\tilde{V}_0^i)'' \leq \frac{C_2}{C_1}, \quad -\frac{C_0}{C_1} \leq (\tilde{g}_0^i)'' \leq 0, \\ \|(\tilde{g}_0^i)''\|_{L^1(\mathbb{R})} = \sqrt{\beta C_1} \frac{1}{C_1} \| (g_0^i)'' \|_{L^1(\mathbb{R})}. \end{aligned} \quad (2.24)$$

Hence \tilde{V}_0^i , \tilde{g}_0^i satisfy the assumptions of Theorem 2.1 with $\beta = 1$ and $C_1 = 1$. Thus

$$D^2 f_M^1(\cdot, \tilde{V}_0^i, \tilde{g}_0^i) \geq \frac{1}{2} |\mathbb{T}_M^d| \text{Id}. \quad (2.25)$$

On the other hand, the change of variables

$$\tilde{\phi}(x) = \sqrt{\beta C_1} \phi(x), \quad \tilde{u} = \sqrt{\beta C_1} u \quad (2.26)$$

yields

$$\tilde{V}_0^i(\tilde{u}_i + \nabla_i \tilde{\phi}(x)) = V_0^i(u + \nabla_i \phi(x)) \quad (2.27)$$

and thus

$$Z_M^\beta(u, V_0^i, g_0^i) = (\beta C_1)^{-(|\mathbb{T}_M^d|-1)/2} Z_M^1(\tilde{u}, \tilde{V}_0^i, \tilde{g}_0^i). \quad (2.28)$$

Hence

$$f_M^\beta(u, V_0, g_0) = \text{const}(\beta, C_1) + \frac{1}{\beta} f_M^1(\sqrt{\beta C_1} u, \tilde{V}_0, \tilde{g}_0). \quad (2.29)$$

Thus (2.25) implies (2.13), as claimed.

Step 2: Separation of the Gaussian part

Next we separate the Gaussian part in the Hamiltonian. From now on, we will always assume that $\beta = 1$ and $C_1 = 1$. Set

$$V_1(s) = V_0(s) - \frac{1}{2} s^2, \quad g = V_1 + g_0. \quad (2.30)$$

Then

$$0 \leq V_1'' \leq C_2 - 1, \quad -C_0 \leq g'' \leq C_2 - 1 \quad (2.31)$$

and the Hamiltonian can be rewritten as

$$H(u, \phi) = \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d \frac{1}{2} (u_i + \nabla_i \phi(x))^2 + G(u, \phi), \quad (2.32)$$

where

$$G(u, \phi) = \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d g(u_i + \nabla_i \phi(x)). \quad (2.33)$$

Since for all functions ϕ on the torus and for all $i \in \{1, 2, \dots, d\}$

$$\sum_{x \in \mathbb{T}_M^d} \nabla_i \phi(x) = 0, \quad (2.34)$$

we get

$$H(u, \phi) = \frac{1}{2} |\mathbb{T}_M^d| |u|^2 + \frac{1}{2} \|\nabla \phi\|^2 + G(u, \phi), \quad (2.35)$$

where $\|\nabla \phi\|^2 = \sum_{x \in \mathbb{T}_M} \sum_{i=1}^d |\nabla_i \phi(x)|^2$. Let

$$Z_0 = \int_X e^{-\frac{1}{2} \|\nabla \phi\|^2} m_M(d\phi). \quad (2.36)$$

Then the measure

$$\mu = \frac{1}{Z_0} e^{-\frac{1}{2} \|\nabla \phi\|^2} m_M(d\phi) \quad (2.37)$$

is a Gaussian measure. Its covariance C is a positive definite symmetric operator on X (equipped with a standard scalar product $(\phi, \psi) = \sum_{x \in \mathbb{T}_M^d} \phi(x)\psi(x)$) such that

$$(C^{-1}\phi, \phi) = \|\nabla \phi\|^2, \quad \forall \phi \in X. \quad (2.38)$$

The partition function thus becomes (recall that we take $\beta = 1$)

$$Z_M(u) = Z_0 e^{-\frac{1}{2} |\mathbb{T}_M^d| |u|^2} \int_X e^{-G(u, \phi)} \mu(d\phi). \quad (2.39)$$

Step 3: Decomposition of μ and Helffer-Sjöstrand calculus

By standard Gaussian calculus, $\mu = \mu_1 * \mu_2$, where μ_1 and μ_2 are Gaussian with covariances

$$C_1 = \lambda C, \quad C_2 = (1 - \lambda)C, \quad \text{where } \lambda \in (0, 1). \quad (2.40)$$

More explicitly, for $i \in \{1, 2\}$

$$\mu_i(d\phi) = \frac{1}{Z_i} e^{-\frac{1}{2\lambda_i} \|\nabla \phi\|^2} m_M(d\phi), \quad \text{where } \lambda_1 = \lambda, \lambda_2 = 1 - \lambda. \quad (2.41)$$

Thus

$$Z_M(u) = Z_0 e^{-\frac{1}{2}|\mathbb{T}_M^d||u|^2} \int_X \int_X e^{-G(u, \psi + \theta)} \mu_1(d\theta) \mu_2(d\psi). \quad (2.42)$$

To write the free energy in a more compact form, we introduce the renormalization maps R_i . For $f \in C(\mathbb{R}^d \times X)$ we define $R_i f$ by

$$e^{-R_i f(u, a)} := \int_X e^{-f(u, a + b)} d\mu_i(b). \quad (2.43)$$

Taking the logarithm of (2.42), we get

$$f_M(u) = \text{const}(M) + \frac{1}{2}|\mathbb{T}_M^d||u|^2 + (R_2 R_1 G)(0, u). \quad (2.44)$$

The main point now is that the map

$$H_1(\theta) = G(u, \psi + \theta) + \frac{1}{2\lambda} \|\nabla \theta\|^2 \quad (2.45)$$

becomes uniformly convex for sufficiently small λ . This will allow us to use the Helffer-Sjöstrand representation to get a good lower bound for $D^2(R_1 G)$, which involves, roughly speaking, the expectation of $G_{i,x}(\theta) = g_0''(u_i + \nabla_i \psi(x) + \nabla_i \theta(x))$ with respect to e^{-H_1} (see (4.82)). This expectation can be controlled in terms of $\|g_0''\|_{L^1(\mathbb{R})}$ (see Lemma 4.1). Under the smallness condition (2.13) one then easily obtains the lower bound for $D^2(R_2 R_1 G)$ (see (4.85) and (4.87)).

3 Consequence of the Helffer-Sjöstrand representation

Let \mathbb{U} and X be finite-dimensional inner product spaces, let C be a positive definite symmetric operator on X and let μ_C be the Gaussian measure with covariance C on X , i.e

$$\mu_C(db) = \frac{1}{Z_C} e^{-\frac{1}{2}(C^{-1}b, b)} db, \quad (3.46)$$

where db is the $\dim X$ dimensional Hausdorff measure on X (i.e $db = \prod db_i$ if the b_i are the coordinates with respect to an orthonormal basis). For a continuous function $f \in C(\mathbb{U} \times X)$ we define $R_C f$ by

$$e^{-R_C f(u, a)} = \int_X e^{-f(u, a + b)} d\mu_C(db). \quad (3.47)$$

In the situation we will consider, $b \rightarrow f(u, a + b) + \frac{1}{2}(C^{-1}b, b)$ will be convex and hence bounded from below so that the right hand side of the above identity is strictly positive.

For $f \in C^2(\mathbb{U} \times X)$ we write $D^2 f(u, a)$ for the Hessian at (u, a) , viewed as an operator from $\mathbb{U} \times X$ to itself. The restriction of the Hessian to X is denoted by $D_X^2 f := P_X D^2 f P_X$, where P_X is the orthogonal projection $\mathbb{U} \times X \rightarrow X$. On the level of quadratic forms we thus have

$$(D_X^2 f(u, a)(\dot{u}, \dot{a}), (\dot{u}, \dot{a})) = (D^2 f(u, a)(0, \dot{a}), (0, \dot{a})). \quad (3.48)$$

From the Helffer-Sjöstrand representation of the variance (see, e.g., [9] (2.6.15)) and the duality relation

$$\frac{1}{2} (A^{-1}a, a) = \sup_{b \in D(A^{\frac{1}{2}})} \left((a, b) - \frac{1}{2} (Ab, b) \right), \quad (3.49)$$

which holds for any positive definite self-adjoint operator A on a Hilbert space Y_0 , one immediately obtains the following estimate:

Lemma 3.1 *Suppose that $H \in C^2(X)$, $\sup_X |D^2H| < \infty$ and there exists a $\delta > 0$ such that*

$$D^2H(a) \geq \delta \text{Id}, \quad \forall a \in X. \quad (3.50)$$

Set

$$Y_0 = \{K \in L^2_{loc}(X) : \langle |DK|^2 \rangle_H < \infty\}, \quad (3.51)$$

$$Y = \{K \in Y_0 : \langle \|D^2K\|_{HS}^2 \rangle_H < \infty\}, \quad (3.52)$$

where the derivatives are understood in the weak sense and

$$\|DK^2\|_{HS}^2 := \sum_{x, y \in \mathbb{T}_M^d \setminus \{0\}} \left(\frac{\partial^2}{\partial \phi(x) \partial \phi(y)} K \right)^2 \quad (3.53)$$

denotes the Hilbert-Schmidt norm. Then for all $G \in Y$ we have

$$\text{var}_H G = \sup_{K \in Y} \langle 2(DG, DK) - (DK, D^2HDK) - \|D^2K\|_{HS}^2 \rangle_H \quad (3.54)$$

Therefore

$$\text{var}_H G \leq \sup_{K \in Y} \langle 2(DG, DK) - (DK, D^2HDK) \rangle_H. \quad (3.55)$$

We will use (3.55) from Lemma 3.1 in the proof of the lemma below.

Lemma 3.2 *Suppose that $f \in C^2(\mathbb{U} \times X)$ and $\sup_{\mathbb{U} \times X} |D^2f| < \infty$. Suppose moreover that there exists a $\delta > 0$ such that*

$$D^2f(u, a) + C^{-1} \geq \delta \text{Id}, \quad \forall (u, a) \in \mathbb{U} \times X. \quad (3.56)$$

Then $Rf \in C^2(\mathbb{U} \times X)$ and for all $u, \dot{u} \in \mathbb{U}$, $a, \dot{a} \in X$

$$\begin{aligned} & ((D^2Rf)(u, a)(\dot{u}, \dot{a}), (\dot{u}, \dot{a})) \\ & \geq \inf_{K \in Y} \langle (D^2f(u, a + \cdot)(\dot{u}, \dot{a} - DK(\cdot), (\dot{u}, \dot{a} - DK(\cdot)))) \rangle_{H, a} \\ & + \langle (C^{-1}DK(\cdot), DK(\cdot)) \rangle_{H, u, a} \end{aligned} \quad (3.57)$$

where

$$H_{u, a}(b) = f(u, a + b) + \frac{1}{2}(C^{-1}b, b), \quad (3.58)$$

$$\langle g \rangle_{H, u, a} = \frac{\int g(b) e^{-H_{u, a}(b)} db}{\int e^{-H_{u, a}(b)} db}. \quad (3.59)$$

Proof

We have

$$e^{-Rf(u,a)} = \int_X e^{-[f(u,a+b) + \frac{1}{2}(C^{-1}b,b)]} db. \quad (3.60)$$

It follows from (3.56) that

$$f(u, a + b) + (C^{-1}(a + b), (a + b)) \geq \frac{1}{2}\delta|a + b|^2 - c \quad (3.61)$$

and standard estimates yield

$$f(u, a + b) + (C^{-1}b, b) \geq \frac{1}{4}\delta|b|^2 - c(1 + |a|^2). \quad (3.62)$$

Hence, by the dominated convergence theorem, the right-hand side of (3.60) is a C^2 function in (u, a) and the same applies to Rf since the right-hand side of (3.60) does not vanish.

To prove the estimate (3.57) for D^2Rf , we may assume without loss of generality that $a = 0, u = 0$ (otherwise we can consider the shifted function $f(\cdot - u, \cdot - a)$). Set

$$h(t) := Rf(t\dot{u}, t\dot{a}). \quad (3.63)$$

Then

$$h''(0) = (D^2(Rf))(0, 0)(\dot{u}, \dot{a}), (\dot{u}, \dot{a}). \quad (3.64)$$

Now

$$h(t) = -\log \int_X e^{-f(t\dot{u}, t\dot{a}+b)} \mu_C(db), \quad (3.65)$$

$$h'(t) = \frac{\int_X e^{-f(t\dot{u}, t\dot{a}+b)} Df(t\dot{u}, t\dot{a} + b)(\dot{u}, \dot{a}) \mu_C(db)}{\int_X e^{-f(t\dot{u}, t\dot{a}+b)} \mu_C(db)} \quad (3.66)$$

and

$$h''(0) = \langle (D^2f(0, \cdot)(\dot{u}, \dot{a}), (\dot{u}, \dot{a})) \rangle_H - \text{var}_H Df(0, \cdot)(\dot{u}, \dot{a}), \quad (3.67)$$

where

$$H(b) = f(0, b) + \frac{1}{2}(C^{-1}b, b). \quad (3.68)$$

By assumption,

$$D^2H(b) \geq \delta \text{Id}, \quad (3.69)$$

i.e. H is uniformly convex. \square

Hence by (3.55) from Lemma 3.1

$$-\text{var}_{Hg} \geq \inf_{K \in \mathcal{Y}} \langle -2(Dg, DK) + (DK, D^2HDK) \rangle_H. \quad (3.70)$$

Apply this with

$$g(b) = Df(0, b)(\dot{u}, \dot{a}) \quad (3.71)$$

and write

$$D^2H = D_X^2f + C^{-1}. \quad (3.72)$$

Then

$$\begin{aligned} & -2(Dg, DK) + (DK, D^2HDK) \\ &= -2D^2f(0, \cdot)((\dot{u}, \dot{a}), (0, DK)) + D^2f(0, \cdot)((0, DK), (0, DK)) \\ & \quad + (C^{-1}DK, DK). \end{aligned} \quad (3.73)$$

Together with (3.70) and (3.67) this yields (3.57). \square

4 Proof of Theorem 2.1

By (2.44)

$$f_M(u) = \text{const}(M) + \frac{1}{2}|\mathbb{T}_M^d||u|^2 + (R_2R_1G)(0, u), \quad (4.74)$$

where

$$G(u, \phi) = \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d g^i(u_i + \nabla_i \phi). \quad (4.75)$$

We first estimate D^2R_1G from below. By (2.31)

$$(g^i)'' \geq -C_0 \geq -\bar{C} \quad (4.76)$$

(recall that we always assume $C_1 = 1$). By (2.12), we have $\bar{C} \geq 1$. If we take

$$\lambda = \frac{1}{2\bar{C}} \quad (4.77)$$

then

$$H_{u, \psi}(\theta) := G(u, \psi + \theta) + \frac{1}{\lambda} \|\nabla \theta\|^2 \quad (4.78)$$

is uniformly convex, i.e.

$$D^2H_{u, \psi}(\theta)(\dot{\theta}, \dot{\theta}) \geq \bar{C} \|\nabla \dot{\theta}\|^2 \geq \delta_M \bar{C} \|\dot{\theta}\|^2, \quad (4.79)$$

with $\delta_M > 0$. Here we used the discrete Poincare inequality

$$\|\nabla \eta\|^2 \geq \delta_M \|\eta\|^2 \quad \text{for } \eta \in X \quad (4.80)$$

which follows from a simple compactness argument since \mathbb{T}_M^d is a finite set. Hence, by Lemma 3.2, we have

$$\begin{aligned}
& (D^2 R_1(G)(u, \psi)(\bar{u}, \bar{\psi}), (\bar{u}, \bar{\psi})) \\
& \geq \inf_{K \in Y} \left\{ \left\langle \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d (g^i)''(u_i + \nabla_i \psi(x) + \nabla_i \cdot (x)) \left(u_i + \nabla_i \psi(x) - \nabla_i \frac{\partial K}{\partial \phi(x)}(\cdot) \right)^2 \right. \right. \\
& \quad \left. \left. + \frac{1}{\lambda} \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d \left| \nabla_i \frac{\partial K}{\partial \phi(x)}(\cdot) \right|^2 \right\rangle_{H_{u, \psi}} \right\}, \tag{4.81}
\end{aligned}$$

where Y is defined by (3.52). Now $(g^i)'' = (V^i)'' + g_0'' \geq g_0''$ (see (2.30) and (2.31)) and together with the estimate $(a - b)^2 \leq 2a^2 + 2b^2$ and the assumption $-C_0 \leq g_0'' \leq 0$, this yields

$$\begin{aligned}
& (D^2 R_1(G)(u, \psi), (\dot{u}, \dot{\psi}), (\dot{u}, \dot{\psi})) \\
& \geq 2 \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d \left\langle (g_0^i)''(u_i + \nabla_i \psi(x) + \nabla_i \cdot (x)) (u_i + \nabla_i \psi(x))^2 \right\rangle_{H_{u, \psi}} \\
& \quad + \left\langle \left(\frac{1}{\lambda} - 2C_0 \right) \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d \left| \nabla_i \frac{\partial K}{\partial \phi(x)}(\cdot) \right|^2 \right\rangle_{H_{u, \psi}}, \tag{4.82}
\end{aligned}$$

where $\frac{1}{\lambda} - 2C_0 \geq 0$. We will now use the following result, which will be proven at the end of this section.

Lemma 4.1 *For $h \in L^1(\mathbb{R}) \cap C^0(\mathbb{R})$, $\psi \in X$, $x \in \mathbb{T}_M^d$ and $i \in \{1, 2, \dots, d\}$ consider $F \in C(X)$ given by*

$$F(\theta) = h(u_i + \nabla_i \psi(x) + \nabla_i \theta(x)). \tag{4.83}$$

Then

$$\left| \langle F \rangle_{H_{u, \psi}} \right| \leq \frac{2}{\pi} (12d\bar{C})^{1/2} \|h\|_{L^1(\mathbb{R})}. \tag{4.84}$$

Together with (4.82), the smallness condition (2.13) and the relation $\sum_{x \in \mathbb{T}_M^d} \nabla_i \psi(x) = 0$, this lemma yields

$$\begin{aligned}
& D^2 R_1 G(u, \psi)(\dot{u}, \dot{\psi})(\dot{u}, \dot{\psi}) \\
& \geq -\frac{1}{2} \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d \left| \dot{u}_i + \nabla_i \dot{\psi}(x) \right|^2 = -\frac{1}{2} |\mathbb{T}_M^d| |\dot{u}|^2 - \frac{1}{2} \|\nabla \dot{\psi}\|^2. \tag{4.85}
\end{aligned}$$

Thus

$$H_2(\psi) := (R_1 G)(u, \psi) + \frac{1}{2(1 - \lambda)} \|\nabla \psi\|^2 \tag{4.86}$$

is uniformly convex and another application of Lemma 3.2 gives

$$\begin{aligned}
& (D^2(R_2R_1G)(u, 0)(\dot{u}, 0), (\dot{u}, 0)) \\
& \geq \inf_K \left\langle D^2(R_1G)(u, \cdot)(\dot{u}, -DK)(\dot{u}, -DK) + \frac{1}{1-\lambda} \|\nabla DK\|^2 \right\rangle_{H_2} \\
& \geq -\frac{1}{2} |\mathbb{T}_M^d| |\dot{u}|^2 + \inf_K \left\{ \left(\frac{1}{1-\lambda} - \frac{1}{2} \right) \langle \|\nabla DK\|^2 \rangle_{H_2} \right\} \\
& \geq -\frac{1}{2} |\mathbb{T}_m^d| |\dot{u}|^2, \tag{4.87}
\end{aligned}$$

where in the last inequality we used that fact that $\frac{1}{1-\lambda} - \frac{1}{2} \geq 0$. In view of (4.74), this finishes the proof of Theorem 2.1.

Proof of Lemma 4.1

Note that u and ψ are fixed. Since the function $\tilde{h}(s) = h(u_i + \nabla_i \psi(x) + s)$ has the same L^1 norm as h , it suffices to prove the estimate for the function $F \in C(X)$ given by

$$F(\theta) = h(\nabla_i \theta(x)). \tag{4.88}$$

Moreover, we write H instead of $H_{u,\psi}$. Let

$$\hat{h}(k) = \int_{\mathbb{R}} e^{-iks} h(s) ds \tag{4.89}$$

denote the Fourier transform of h . Then

$$\|\hat{h}\|_{L^\infty(\mathbb{R})} \leq \|h\|_{L^1(\mathbb{R})} \tag{4.90}$$

and

$$h(s) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{iks} \hat{h}(s) dk. \tag{4.91}$$

Set

$$A(k) = \langle F_k \rangle_H, \quad \text{where } F_k(\theta) = e^{ik\nabla_i \theta(x)}. \tag{4.92}$$

Then

$$\langle F \rangle_H = \frac{1}{2\pi} \int_{\mathbb{R}} A(k) h(k) dk \tag{4.93}$$

and, in view of (4.90), it suffices to show that

$$\int_{\mathbb{R}} |A(k)| dk \leq 4(12d\bar{C})^{1/2}. \tag{4.94}$$

First note that $|F_k| = 1$. Hence

$$|A(k)| \leq 1, \quad \forall k \in \mathbb{R}. \tag{4.95}$$

To get decay of $A(k)$ for large k we use integration by parts. First note that for $G_i \in C^1(X)$, with $\sup_{a \in X} e^{-\delta|a|} (|G_i|(a) + |DG_i|(a)) < \infty$ for all $\delta > 0$, we have

$$\left\langle \frac{\partial G_1}{\partial \phi(x)} G_2 \right\rangle_H = \left\langle -G_1 \frac{\partial G_2}{\partial \phi(x)} \right\rangle_H + \left\langle \frac{\partial H}{\partial \phi(x)} G_1 G_2 \right\rangle_H. \quad (4.96)$$

Assume first that $x \in \mathbb{T}_M^d \setminus \{0\}$. Then

$$F_k(\theta) = -\frac{1}{k^2} \frac{\partial^2 F_k}{\partial \theta^2(x)}(\theta) \quad (4.97)$$

and thus

$$\begin{aligned} -k^2 A(k) &= \left\langle \frac{\partial^2 F_k}{\partial \theta^2(x)} \cdot 1 \right\rangle_H = \left\langle \frac{\partial F_k}{\partial \theta(x)} \frac{\partial H}{\partial \theta(x)} \right\rangle_H \\ &= -\left\langle F_k \frac{\partial^2 H}{\partial \theta^2(x)} \right\rangle_H + \left\langle F_k \left(\frac{\partial H}{\partial \theta(x)} \right)^2 \right\rangle_H. \end{aligned} \quad (4.98)$$

Since $|F_k| = 1$, this yields

$$|A(k)| \leq \frac{1}{k^2} \left\langle \left| \frac{\partial^2 H}{\partial \theta^2(x)} \right| \right\rangle_H + \frac{1}{k^2} \left\langle \left(\frac{\partial H}{\partial \theta(x)} \right)^2 \right\rangle_H. \quad (4.99)$$

Application of (4.96) with $G_2 = 1, G_1 = \frac{\partial H}{\partial \theta(x)}$ gives

$$\left\langle \frac{\partial^2 H}{\partial \theta^2(x)} \right\rangle_H = \left\langle \left(\frac{\partial H}{\partial \theta(x)} \right)^2 \right\rangle_H. \quad (4.100)$$

Thus

$$|A(k)| \leq \frac{2}{k^2} \left\langle \left| \frac{\partial^2 H}{\partial \theta^2(x)} \right| \right\rangle_H. \quad (4.101)$$

Now recall that

$$H(\theta) = \sum_{x \in \mathbb{T}_M^d} \sum_{i=1}^d g^i (u_i + \nabla_i \psi(x) + \nabla_i \theta(x)) + \frac{1}{2\lambda} |\nabla_i \theta(x)|^2. \quad (4.102)$$

Since $\lambda^{-1} = 2\bar{C}$, it follows that

$$\left| \frac{\partial^2 H}{\partial \theta^2(x)} \right| \leq 2d \left(\sup_{\mathbb{R}} |(g^i)''| + \frac{1}{\lambda} \right) \leq 6d\bar{C}. \quad (4.103)$$

Hence

$$|A(k)| \leq \frac{12d\bar{C}}{k^2}. \quad (4.104)$$

Using (4.104) for $|k| \geq (12d\bar{C})^{1/2}$ and (4.95) for $|k| \leq (12d\bar{C})^{1/2}$, we get (4.94).

Finally, if $x = 0$ we note that

$$F_k(\theta) = -\frac{1}{k^2} \frac{\partial^2}{\partial \theta^2(e_i)} F_k(\theta) \quad (4.105)$$

and we proceed as before. \square

Remark 4.2 The proof shows that for $h = g''$ we can also use norms involving only lower derivatives of g . In particular, we have

$$\begin{aligned}
|\langle g'' \rangle_H| &\leq \frac{1}{2\pi} \int_{\mathbb{R}} |\hat{g}''(k)| |A(k)| dk \\
&\leq \frac{1}{2\pi} \|\hat{g}'(k)\|_{L^2(\mathbb{R})} \left(\int_{\mathbb{R}} k^2 |A(k)|^2 dk \right)^{1/2} \\
&\leq \frac{1}{\sqrt{2\pi}} \|g'\|_{L^2(\mathbb{R})} \left(2 \left(\frac{1}{3} + (12d\bar{C})^2 \right) \right)^{1/2}, \quad (4.106)
\end{aligned}$$

where we used (4.95) for $|k| \leq 1$ and (4.104) for $|k| \geq 1$.

Remark 4.3 Note that our proofs can be very easily adapted to any decomposition of $\mu = \mu_1 * \mu_2$, where μ_1 and μ_2 are Gaussian with covariances C_1 and C_2 , such that $H_{u,\psi}(\theta) := G(u, \psi + \theta) + \frac{1}{2}(C^{-1}\theta, \theta)$ is uniformly convex.

Remark 4.4 The procedure for the one-step decomposition can be iterated and the proofs can be adapted to the multi-scale decomposition; iterating the method would lower the temperature and weaken the conditions on the perturbation function g . However, our iteration procedure would not allow us to get results involving the low temperature case.

Examples

- (a) $V(s) = s^2 + a - \log(s^2 + a)$, where $0 < a < 1$. Then $C_1 = C_2 = 2$, $C_0 = \frac{2}{a}$, $\|(g_0^i)''\|_{L^1(\mathbb{R})} = 2\sqrt{\frac{1}{a}}$ and $\beta \leq \frac{a^2 \pi^2}{6 \times 16^2 d}$.




Figure 1: Example (a)

- (b) Let $0 < \delta < 1$ and

$$V(s) = \begin{cases} \frac{x^2}{2} - \frac{4}{\delta^4} x^3 (\delta - x)^3 & \text{if } 0 \leq x \leq \delta \\ \frac{x^2}{2} & \text{otherwise.} \end{cases}$$

Then $C_1 = C_2 = 1$, $\bar{C} = \frac{6}{5}$, $\|(g_0^i)''\|_{L^1(\mathbb{R})} \leq \frac{3}{10\sqrt{5}}\delta^5$ and $\beta \leq \left(\frac{5\sqrt{5d\pi}}{2\delta}\right)^2$.

Note that if $\delta \ll 1$, the surface tension is convex for very large values of β .




Figure 2: Example (b)

(c) Let $p \in (0, 1)$ and $0 < k_2 < k_1$. Let

$$V(s) = -\log\left(pe^{-k_1 \frac{s^2}{2}} + (1-p)e^{-k_2 \frac{s^2}{2}}\right).$$

Then

$$V_0''(s) = \frac{pk_1 e^{-k_1 \frac{s^2}{2}} + (1-p)k_2 e^{-k_2 \frac{s^2}{2}}}{pe^{-k_1 \frac{s^2}{2}} + (1-p)e^{-k_2 \frac{s^2}{2}}}$$

and

$$g_0''(s) = -\frac{p(1-p)(k_1 - k_2)^2 s^2}{p^2 e^{-(k_1 - k_2) \frac{s^2}{2}} + 2p(1-p) + (1-p)^2 e^{(k_1 - k_2) \frac{s^2}{2}}}.$$

We have

$$k_2 \leq V_0''(s) \leq pk_1 + (1-p)k_2 \quad \text{and} \quad -\frac{p(k_1 - k_2)}{1-p} \leq g_0''(s) \leq 0,$$

where the lower bound inequality for $g_0''(s)$ follows from the fact that $g_0''(s)$ attains its minimum for $s \geq \sqrt{\frac{2}{k_1 - k_2}}$. Then

$$\|g_0''(s)\|_{L^1(\mathbb{R})} \leq \frac{2p}{1-p} \sqrt{(k_1 - k_2)\pi} \quad \text{and} \quad \beta \leq \left(\frac{1-p}{16p}\right)^2 \frac{\pi k_2}{12d\bar{C}(k_1 - k_2)}.$$

Note that example c) is the one used in [2] to prove that unicity of ergodic states can be violated for non-convex V for large enough β .

Acknowledgment

Codina Cotar thanks David Brydges and Haru Pinson for invaluable advice and suggestions during the writing of the manuscript.

References

- [1] Adams, S., Kotecky, R., Müller, S., in preparation.
- [2] Biskup, M., Kotecky, R., *Phase coexistence of gradient Gibbs states*, Probab. Theory Relat. Fields, 139, 1-39, (2007).
- [3] Biskup, M., Spohn, M., *Scaling limit for a class of gradient fields with non-convex potentials*, preprint.
- [4] Brydges, D., *Park City lecture notes*, 2007, in preparation.
- [5] Brydges, D., Yau, S.T., *Grad ϕ perturbations of massless Gaussian fields*, Comm. Math. Phys. 129, no. 2, 351-392, (1990).
- [6] Cotar, C., Deuschel, J.D., *Decay of covariances, uniqueness of ergodic component and scaling limit for $\nabla\phi$ systems with non-convex potential*, preprint.
- [7] Deuschel, J.D., Giacomin, G., Ioffe, D., *Large deviations and concentration properties for $\nabla\phi$ interface models*, Prob. Theory Relat. Fields 117, 49-111, (2000).
- [8] Fröhlich, J. Pfister, *On the absence of spontaneous symmetry breaking and of crystalline ordering in two-dimensional systems*. Comm. Math. Phys. 81, 277-298 (1981).
- [9] Helffer, B., *Semiclassical analysis, Witten Laplacians and statistical mechanics*, World Scientific, (2002).
- [10] Funaki, T., Spohn, H., *Motion by Mean Curvature from the Ginzburg-Landau $\nabla\phi$ Interface Model*, Commun. Math. Phys. 185, 1-36, (1997).
- [11] Pinson, H., *Towards a Nonperturbative Renormalization Group Analysis*, preprint.