

RESEARCH ARTICLE

Cryopreservation of Endothelial Cells in Various Cryoprotective Agents and Media – Vitrification versus Slow Freezing Methods

Achim von Bomhard^{1*}, Alexander Elsässer², Lucas Maximilian Ritschl¹, Silke Schwarz², Nicole Rotter²

1 Technical University of Munich, Department of Maxillofacial Surgery, Langerstr. 3, 81675 Munich, Germany, **2** University Medical Center Ulm, Department of Otorhinolaryngology, Frauensteige 12, 89075 Ulm, Germany

* achimvonbomhard@gmail.com



Abstract

Vitrification of endothelial cells (MHECT-5) has not previously been compared with controlled slow freezing methods under standardized conditions. To identify the best cryopreservation technique, we evaluated vitrification and standardized controlled-rate $-1^{\circ}\text{C}/\text{minute}$ cell freezing in a -80°C freezer and tested four cryoprotective agents (CPA), namely dimethyl sulfoxide (DMSO), ethylene glycol (EG), propylene glycol (PG), and glycerol (GLY), and two media, namely Dulbecco's modified Eagle medium Ham's F-12 (DMEM) and K^{+} -modified TiProtec (K^{+}TiP), which is a high-potassium-containing medium. Numbers of viable cells in proliferation were evaluated by the CellTiter 96[®] A_{QUEOUS} One Solution Cell Proliferation Assay (Promega Corporation, Mannheim, Germany). To detect the exact frozen cell number per cryo vial, DNA content was measured by using Hoechst 33258 dye prior to analysis. Thus, results could be evaluated unconstrained by absolute cell number. Thawed cells were cultured in 25 cm^2 cell culture flasks to confluence and examined daily by phase contrast imaging. With regard to cell recovery immediately after thawing, DMSO was the most suitable CPA combined with K^{+}TiP in vitrification ($99 \pm 0.5\%$) and with DMEM in slow freezing ($92 \pm 1.6\%$). The most viable cells in proliferation after three days of culture were obtained in cells vitrified by using GLY with K^{+}TiP ($308 \pm 34\%$) and PG with DMEM in slow freezing ($280 \pm 27\%$).

OPEN ACCESS

Citation: von Bomhard A, Elsässer A, Ritschl LM, Schwarz S, Rotter N (2016) Cryopreservation of Endothelial Cells in Various Cryoprotective Agents and Media – Vitrification versus Slow Freezing Methods. PLoS ONE 11(2): e0149660. doi:10.1371/journal.pone.0149660

Editor: Boris Rubinsky, University of California at Berkeley, UNITED STATES

Received: November 3, 2015

Accepted: February 3, 2016

Published: February 18, 2016

Copyright: © 2016 von Bomhard et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The authors have no support or funding to report.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

The endothelium is the monolayer of endothelial cells lining the lumen of all blood vessels. Disorders in the endothelium predispose the vessel wall to vasoconstriction, leukocyte adherence, platelet activation, mitogenesis, pro-oxidation, thrombosis, impaired coagulation, vascular inflammation, and atherosclerosis [1]. To be able to study all these vascular disorders in vitro, the cryopreservation, storage, and shipment of endothelial cells is extremely important. Relevant cell culture models are also crucial to the study of the pathobiology of the lung microvascular endothelium and for an understanding of lung metastasis [2] and disorders such as acute

respiratory distress syndrome [3, 4]. Large numbers of quality-controlled cells are required for in vitro research and could, potentially, be used for cell therapy in clinical application [5].

Cryopreservation with controlled slow cooling rate was first described by Polge, Smith and Parkes in 1949 using the protective properties of glycerol (GLY) [6]. Dimethyl sulfoxide (DMSO) was proposed as a cryoprotectant in 1960 by Lovelock & Bishop and was rapidly shown to have far more widespread applicability than glycerol, particularly for the preservation of cells in tissue culture [7]. The literature of the 1950s and 1960s is dominated by reports of efforts to achieve or improve the cryopreservation of a variety of cell types through empirical variations of freezing rates, thawing rates, cryoprotectant concentrations, and associated solutes [8]. Nevertheless, a recovery of viability of more than 80–90% is still rarely achieved.

Slow cooling avoids intracellular ice buildup, which can cause the rupture of the cell membrane at temperatures between 0°C and –70°C [9]. However, the formation of extracellular ice can still result in the dehydration of the cells. To prevent this, an ideal cooling rate should be chosen, and a cryoprotective agent (CPA) added [10]. CPAs are divided into intracellular agents, which penetrate inside the cells and prevent ice crystal formation and membrane rupture, and extracellular compounds that do not penetrate the cell membrane and act by reducing the hyperosmotic effect present during the freezing procedure [10]. DMSO, ethylene glycol (EG), propylene glycol (PG), and GLY are intracellular CPAs. Among the extracellular compounds are sucrose, trehalose, dextrose, and polyvinylpyrrolidone [11]. DMSO is the most commonly used CPA. It provides a high rate of postfreezing cell survival but presents chemical cytotoxicity at room temperature and can damage the cells osmotically [12]. This cytotoxicity forces the experimenter to work rapidly. Moreover, different workers require different amounts of time to thaw their cells completely and to wash them free of CPA. Therefore, less cytotoxic CPAs are needed for the standardization of such procedures.

Over the last few years, another fast freezing method has become increasingly popular: vitrification. This promising and easy process avoids damage to the cells caused by their dehydration during slow freezing. To prevent intracellular ice buildup, a higher concentration of CPA is necessary, but its cytotoxic effect can be reduced by immediate freezing in liquid nitrogen (LN₂). Vitrification has not previously been compared with controlled slow freezing methods under standardized conditions for endothelial cells. Many reports have shown that the toxicity to cells not only depends on the chemical properties of CPAs, but also on the cell types to be frozen. The ways in which the various CPAs and their combinations affect a given cell type cannot be predicted, and hence, the selection of cryoprotectants for each type of cell is still largely dependent on an empirical and experimental approach [13].

To identify the best cryopreservation technique for endothelial cells, four CPAs, namely DMSO, EG, PG, and GLY, and two different types of medium were tested in two different cryopreservation protocols (slow freezing vs. vitrification).

Materials and Methods

All reagents, cell culture media, and supplements were purchased from Sigma Aldrich (Munich, Germany), Invitrogen (Darmstadt, Germany), and Biochrom (Berlin, Germany), unless indicated otherwise. Cell culture material was obtained from Greiner Bio One (Frickenhäusen, Germany).

Derivation and culture of endothelial cells

Murine heart endothelial cells (MHEC5-T, Leibniz Institute DSMZ—German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) were used in this study. Cell

culture was performed as recommended by DSMZ at 37°C and 5% CO₂. Cells were seeded at 0.1 x10⁵ cells/cm² in monolayer culture.

Cryopreservation solution (CPS), vitrification solution (VS), and cell culture media

DMEM Ham's F-12 + 2mM L-glutamine (Gln+) and K⁺TiP were tested as carrier solutions. K⁺TiP was prepared by supplementing 500mL TiProtec (Dr. Franz Köhler Chemie GmbH, Bensheim, Germany) with 0.871g K₂SO₄ and 3.87g 56.3% w/w potassium lactate and was diluted with 50mL distilled water. [Table 1](#) shows the final composition of K⁺TiP.

Prior to freezing or vitrification, the cryopreservation solution (CPS) and vitrification solution (VS) were prepared containing foetal bovine serum (FBS), sucrose (in VS), and one of the CPAs.

CPS was used for the controlled-rate -1°C/minute freezing. It contained one of the two tested carrier solutions, 40% FBS, and 3M of one of the following CPAs: DMSO, EG, PG, or GLY. For freezing, CPS was added 1:1 to cells in precooled carrier solution; therefore, the final concentration of FBS was 20% and of each CPA was 1.5M.

VS was used for vitrification. It contained one of the two tested carrier solutions, 40% FBS, 1M sucrose, and 6M of one of the following CPAs: DMSO, EG, PG, or GLY. For vitrification, the VS was added 1:1 to cells in precooled carrier solution; therefore, the final concentration of FBS was 20%, that of sucrose was 0.5M, and that of each CPAs was 3M. DMSO, GLY, and EG are reported to lead reliably to vitrification with tolerable toxicity at a concentration of 3M [14]. PG was also set to 3M in order to compare its efficacy.

The cell culture media and warming solution was DMEM Ham's F-12 Gln+ supplemented with 10% FBS. [Table 2](#) shows the features of the four cryoprotective agents used.

Table 1. Composition of K⁺TiP.

Cl-	94	mM
H ₂ PO ₄ -	1	mM
α-ketoglutarate	2	mM
aspartate	5	mM
Na+	21	mM
K+	131	mM
Mg ²⁺	7	mM
Ca ²⁺	0.02	mM
N-acetylcysteine	27	mM
glycine	9	mM
alanine	5	mM
tryptophan	2	mM
saccharose	27	mM
glucose	9	mM
Desferal	0.075	mM
LK 614	0.015	mM
osmolarity	338	mosm/l
pH	6.93	

K⁺TiP was prepared by supplementing 500 mL TiProtec (Dr. Franz Köhler Chemie GmbH, Bensheim, Germany) with 0.871g K₂SO₄ and 3.87g 56.3% w/w potassium lactate and was diluted with 50mL distilled water prior to use. Table 1 shows the final composition of K⁺TiP.

doi:10.1371/journal.pone.0149660.t001

Table 2. Cryoprotective agents.

	dimethyl sulfoxide (DMSO)	ethylene glycol (EG)	glycerol (GLY)	propylene glycol (PG)	
molar mass (M)	78.13	62.07	92.09	76.09	g/mol
density	1.19	1.11	1.26	1.04	g/cm ³
purity	0.999	0.999	0.999	0.999	%
molarity	15.21579419	17.86515225	13.66858508	13.65435668	mol/L

Table 2 shows the features of the four cryoprotective agents used.

doi:10.1371/journal.pone.0149660.t002

Cryopreservation and thawing

MHEC5-Ts were frozen at a final concentration of 10^6 cells/mL by two different methods. At 80% confluence in Monolayer culture, the cells were trypsinized and counted, and 1×10^6 cells were transferred in 500 μ L precooled 4°C DMEM Ham’s F-12 Gln+ or 500 μ L precooled 4°C K⁺TiP into cryo vials. In the controlled-rate -1°C/minute freezing protocol, 500 μ L CPS was added, the cap was immediately tightened, and the vials were placed in a CoolCell (BioCision, Mill Valley, USA) in a -80°C freezer, left over night, and transferred into liquid nitrogen (LN₂) the next day. Alternatively, for vitrification, 500 μ L VS was added, the cap was tightened, and vials were directly plunged into LN₂. The whole process was performed within less than 15 seconds per vial. The samples were stored for 3 and 6 months.

Thawing was carried out quickly in a water bath at 37°C. To accelerate the warming process and to dilute the concentration of CPA, 1mL 37°C warming solution was added. Immediately after thawing, all samples were washed with 40mL DMEM Ham’s F-12 Gln+ to remove residual CPA, pelleted by centrifugation at 200x g for 5min, and resuspended in 2mL culture media. After repetitive gentle vortexing, 1mL of this cell suspension was used to determine the exact cell number by using Hoechst 33258. Aliquots of 500 μ L were used to determine cell viability at day 0, 1, 2, and 3. The residual 500 μ L of each sample was plated in a 25cm² cell culture flask.

Ice formation during vitrification and thawing was evaluated by visual inspection for the presence of a milky appearance as described by Sheffen et al. [15].

Determination of cell viability, number, recovery, and cell doubling time

The CellTiter 96® A_{QUEOUS} One Solution Cell Proliferation Assay (COS, Promega Corporation, Mannheim, Germany) was used to determine viable cells in proliferation at one hour after thawing and at day one, two, and three. COS reagent contains a novel tetrazolium compound [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium, inner salt; MTS] and an electron-coupling reagent (phenazine ethosulfate; PES). The MTS tetrazolium compound (Owen’s reagent) is bioreduced by cells into a colored formazan product that is soluble in tissue culture medium. This conversion is presumed to be accomplished by NADPH or NADH produced by dehydrogenase enzymes in metabolically active cells [16]. Aliquots of 20 μ L of the COS reagent were pipetted directly into transparent 96-well cell culture plates containing endothelial cells in 100 μ L culture media and incubated for 1.5 hours. The absorbance was then recorded at 490nm with a 96-well plate reader (Infinite 200 PRO series, Tecan Group Ltd., Männedorf, Switzerland).

To detect the exact cell number per cryo vial, DNA content was measured according to Kim et al. by using Hoechst 33258 dye [17]. Cells were digested by using 0.5mg/mL proteinase K (AppliChem, Darmstadt, Germany) in 30mM Tris-HCl pH 8.0 at 56°C in a thermomixer comfort (Eppendorf, Hamburg, Germany) at 1000rpm. The digest was centrifuged at 2500x g for 5min. Aliquots of 100 μ L of each supernatant were diluted 1:5 in TNE-buffer (10mM Tris,

1mM Na₂EDTA, 0.1mM NaCl, pH 7.4), and 500 μ L Hoechst 33258 dye 0.1 μ g/mL in TNE-buffer was added. After 20min, fluorescence was measured in triplicate in a black 96-well plate with a 96-well plate reader (Infinite® 200 PRO series, Tecan Group Ltd., Männedorf, Switzerland). The excitation wavelength was 356nm with a bandpass of 9nm, and the emission wavelength was 465nm with a bandpass of 20nm. Blanks contained 0.5mg/mL proteinase K in 30mM Tris-HCl pH 8.0 diluted 1:5 with TNE. For the determination of cell number, a calibration curve with MHEC5-Ts was performed. Samples were analyzed on day 0 (each n = 6).

Recovered MHEC5-Ts are able to attach to the culture flask bottom after thawing. Therefore, the cell recovery rate (cells attached to the flask / cells seeded x100%) was calculated by counting all dead cells in the supernatant after 24 hours by using a Neubauer chamber (Carl Roth, Karlsruhe, Germany) and trypan blue.

MHEC-5Ts doubling time could be analyzed by comparing COS-assay values immediately after thawing and on day 3 by using a standard curve.

The 25cm² cell culture flasks were analyzed microscopically and photographed in six specific areas under phase contrast on a microscope (Axio Observer A1 inverted stand, Carl Zeiss GmbH, Jena, Germany) every day until confluence was reached. Cell numbers on day one till four were counted in these photographs with adobe illustrator CS5 (Adobe Systems, Delaware, USA).

Statistical analysis

The two-tailed Mann-Whitney test was performed by using GraphPad Prism (version 6.0b for Mac OS X, GraphPad Software, La Jolla, USA). Values were considered significant when $p < 0.05$ (*), medium significant when $p < 0.01$ (**), highly significant when $p < 0.001$ (***), and of the highest significance when $p < 0.0001$ (****).

Results

Standardized controlled-rate -1°C/minute cell freezing in a -80°C freezer

On thawing, cell recovery of MHEC-5Ts that had been frozen utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was 91.6 \pm 1.6% with DMSO, 83.6 \pm 3.8% with PG, 69.6 \pm 1.3% with EG, and 53.0 \pm 13.1% with GLY as the CPA. On thawing, cell recovery of MHEC-5Ts that had been frozen utilizing K⁺TiP as a cryopreservation medium was 81.7 \pm 5.2% with DMSO, 76.3 \pm 8.3% with EG, 64.7 \pm 5.6% with PG, and 41.3 \pm 4.4% with GLY as the CPA. [Fig 1](#) illustrates these findings.

The numbers of viable cells in proliferation were determined colorimetrically. The highest absorbance at 1.5 hours after thawing relative to the cell count following the use of DMEM Ham's F-12 Gln+ as the cryopreservation medium was 0.62 \pm 0.06 x10⁻⁷ absorbance units (au) (59 \pm 5.9% relative to unfrozen cells) with DMSO, and then 0.52 \pm 0.03 x10⁻⁷ au (48.8 \pm 3%) with PG, 0.44 \pm 0.02 x10⁻⁷ au (41.6 \pm 2.1%) with EG, and 0.18 \pm 0.04 x10⁻⁷ au (17.2 \pm 3.9%) with GLY as the CPA. The highest absorbance after thawing following the use of K⁺TiP as the cryopreservation medium was 0.65 \pm 0.01 x10⁻⁷ au (62.4 \pm 1.2%) with DMSO, and then 0.54 \pm 0.03 x10⁻⁷ au (50.8 \pm 2.7%) with PG, 0.43 \pm 0.09 x10⁻⁷ au (41 \pm 8.4%) with EG, and 0.14 \pm 0.03 x10⁻⁷ au (13.6 \pm 2.6%) with GLY as the CPA.

At 1 day after thawing, the highest absorbance utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was 1.03 \pm 0.14 x10⁻⁷ au (97.8 \pm 12.8%) with DMSO, and then 0.83 \pm 0.11 x10⁻⁷ au (78.2 \pm 10.1%) with PG, 0.7 \pm 0.02 x10⁻⁷ au (66.3 \pm 1.8%) with EG, and 0.14 \pm 0.02 x10⁻⁷ au (13.1 \pm 2.3%) with GLY as the CPA. The highest absorbance after thawing utilizing K⁺TiP as cryopreservation medium was 0.77 \pm 0.09 x10⁻⁷ au (74.5 \pm 8.4%) with DMSO,




Fig 1. Cell recovery by using slow freezing methods. Figure 1 illustrates cell recovery of slowly frozen MHEC-5Ts immediately after thawing. DMEM = DMEM Ham's F-12 Gln+, 20% FBS, and 1.5M of one of the CPAs; K⁺TiP = K⁺TiP, 0.5M sucrose, 20% FBS, and 1.5M of one of the CPAs; ** = medium significant ($p < 0.01$); **** = highest significance ($p < 0.0001$).

doi:10.1371/journal.pone.0149660.g001

and then $0.61 \pm 0.03 \times 10^{-7}$ au ($58.2 \pm 2.5\%$) with PG, $0.61 \pm 0.15 \times 10^{-7}$ au ($57.6 \pm 14.3\%$) with EG, and $0.15 \pm 0.03 \times 10^{-7}$ au ($13.7 \pm 2.6\%$) with GLY as the CPA.

Two days after thawing, the highest absorbance utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was $2.21 \pm 0.21 \times 10^{-7}$ au ($209.5 \pm 20.3\%$) with DMSO, and then $2.03 \pm 0.24 \times 10^{-7}$ au ($192 \pm 22.5\%$) with PG, $1.68 \pm 0.03 \times 10^{-7}$ au ($159.3 \pm 2.5\%$) with EG, and $0.17 \pm 0.03 \times 10^{-7}$ au ($15.7 \pm 2.9\%$) with GLY as the CPA. The highest absorbance after thawing utilizing K⁺TiP as cryopreservation medium was $1.94 \pm 0.18 \times 10^{-7}$ au ($183.3 \pm 16.8\%$) with DMSO, and then $1.56 \pm 0.07 \times 10^{-7}$ au ($147.9 \pm 6.7\%$) with PG, $1.5 \pm 0.37 \times 10^{-7}$ au ($141.8 \pm 35.4\%$) with EG, and $0.2 \pm 0.04 \times 10^{-7}$ au ($19.1 \pm 3.5\%$) with GLY as the CPA.

Three days after thawing, the highest absorbance utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was $2.95 \pm 0.28 \times 10^{-7}$ au ($279.6 \pm 26.9\%$) with PG, and then $2.5 \pm 0.1 \times 10^{-7}$ au ($238.1 \pm 9.5\%$) with EG, $2.31 \pm 0.12 \times 10^{-7}$ au ($218.6 \pm 11.4\%$) with DMSO, and $0.29 \pm 0.05 \times 10^{-7}$ au ($27.7 \pm 4.5\%$) with GLY as the CPA. The highest absorbance after thawing utilizing K⁺TiP as the cryopreservation medium was $2.66 \pm 0.15 \times 10^{-7}$ au ($251.7 \pm 14.1\%$) with DMSO, and then $2.64 \pm 0.28 \times 10^{-7}$ au ($249.8 \pm 26.3\%$) with PG, $2.41 \pm 0.43 \times 10^{-7}$ au ($228.6 \pm 41\%$) with EG, and $0.48 \pm 0.15 \times 10^{-7}$ au ($45.4 \pm 14.5\%$) with GLY as the CPA. [Fig 2](#) illustrates these findings.

The fastest MHEC-5T doubling-time following the use of DMEM Ham's F-12 Gln+ as the cryopreservation medium was 25 ± 0.6 h with EG, and then 25 ± 1 h with PG, 39 ± 0.7 h with DMSO, and 88 ± 3.4 h with GLY as the CPA. Doubling-time utilizing K⁺TiP as the cryopreservation media was 26 ± 1.1 h with EG, 29 ± 1.2 h with PG, 36 ± 0.8 h with DMSO, and 45 ± 5.8 h with GLY as the CPA. [Fig 3](#) illustrates these findings.




Fig 2. Viable cells in proliferation after use of slow freezing methods. The CellTiter 96[®] A_{QUEOUS} One Solution Cell Proliferation Assay (COS, Promega Corporation, Mannheim, Germany) was used to determine viable cells in proliferation at one hour after thawing and at day one, two, and three. Figure 2 illustrates the absorption of viable cells in proliferation after the use of slow freezing methods. DMEM = DMEM Ham's F-12 Gln+, 20% FBS, and 1.5M of one of the CPAs; K⁺TiP = K⁺TiP, 0.5M sucrose, 20% FBS, and 1.5M of one of the CPAs; DMSO = dimethyl sulfoxide; EG = ethylene glycol; PG = propylene glycol; GLY = glycerin; * = significant (p < 0.05); ** = medium significant (p < 0.01); au = absorbance units.

doi:10.1371/journal.pone.0149660.g002

All frozen cells could be cultured until confluence in the 25cm² cell culture flasks after thawing. Confluence was reached on day 4 by cells previously frozen utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and DMSO as the CPA and utilizing K⁺TiP as the cryopreservation medium and DMSO or EG as the CPA. On day 5, confluence was reached by cells previously frozen utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and EG or PG as the CPA and utilizing K⁺TiP as the cryopreservation medium and PG as the CPA. On day 6, confluence was reached by cells previously frozen by utilizing K⁺TiP as the cryopreservation medium and GLY as the CPA. On day 8, confluence was reached by cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and GLY as the CPA.

Vitrification

After thawing, cell recovery of MHEC-5Ts frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was 98.6 ± 1.1% with DMSO, 57.4 ± 17.6% with EG, 56.9 ± 8.6% with GLY, and 52.0 ± 1.0% with PG as the CPA. After thawing, cell recovery of MHEC-5Ts frozen by utilizing K⁺TiP as the cryopreservation medium was 99.7 ± 0.5% with DMSO,




Fig 3. Cell doubling-time after the use of slow freezing methods. Figure 3 illustrates MHEC-5T doubling-time after the use of slow freezing methods. DMEM = DMEM Ham's F-12 Gln+, 20% FBS, and 1.5M of one of the CPAs; K+TiP = K+TiP. 0.5M sucrose, 20% FBS, and 1.5M of one of the CPAs; **** = highest significance ($p < 0.0001$).

doi:10.1371/journal.pone.0149660.g003

93.8 ± 3.6% with GLY, 75.9 ± 13.1% with EG, and 61.4 ± 1.0% with PG as the CPA. Fig 4 illustrates these findings.

The numbers of viable cells in proliferation were determined colorimetrically. The highest absorbance relative to cell count 1.5 hours after the thawing of cells frozen by utilizing DMEM Ham's F-12 Gln+ as cryopreservation media was $0.85 \pm 0.05 \times 10^{-7}$ au (80.7 ± 4.9% relative to unfrozen cells) with DMSO, and then $0.16 \pm 0.04 \times 10^{-7}$ au (14.7 ± 3.3%) with GLY, $0.11 \pm 0.01 \times 10^{-7}$ au (10 ± 0.6%) with PG, and $0.1 \pm 0.01 \times 10^{-7}$ au (9.2 ± 1.2%) with EG as the CPA. The highest absorbance after thawing following the use of K+TiP as the cryopreservation medium was $0.98 \pm 0.06 \times 10^{-7}$ au (92.6 ± 5.4%) with DMSO, and then $0.41 \pm 0.05 \times 10^{-7}$ au (38.7 ± 4.8%) with GLY, $0.19 \pm 0.05 \times 10^{-7}$ au (17.5 ± 5%) with PG, and $0.08 \pm 0.01 \times 10^{-7}$ au (7.2 ± 1%) with EG as the CPA.

At 1 day after thawing, the highest absorbance of cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation media was $1.8 \pm 0.09 \times 10^{-7}$ au (170.8 ± 8.7% relative to unfrozen cells) with DMSO, and then $0.19 \pm 0.05 \times 10^{-7}$ au (17.9 ± 5.1%) with GLY, $0.09 \pm 0.01 \times 10^{-7}$ au (8.6 ± 0.8%) with EG, and $0.08 \pm 0.01 \times 10^{-7}$ au (8 ± 1%) with PG as the CPA. The highest absorbance after thawing cells previously frozen by utilizing K+TiP as the cryopreservation medium was $1.93 \pm 0.03 \times 10^{-7}$ au (182.7 ± 3.1%) with DMSO, and then $0.74 \pm 0.13 \times 10^{-7}$ au (70.4 ± 12.6%) with GLY, $0.18 \pm 0.06 \times 10^{-7}$ au (17 ± 6.1%) with PG, and $0.06 \pm 0.01 \times 10^{-7}$ au (6 ± 0.8%) with EG as the CPA.

Two days after thawing, the highest absorbance of cells frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was $2.9 \pm 0.2 \times 10^{-7}$ au (274.3 ± 18.6% relative to unfrozen cells) with DMSO, and then $0.38 \pm 0.14 \times 10^{-7}$ au (35.7 ± 13.6%) with GLY, $0.14 \pm 0.01 \times 10^{-7}$ au (12.9 ± 0.5%) with PG, and $0.13 \pm 0.03 \times 10^{-7}$ au (12.3 ± 2.5%) with EG as the CPA. The




Fig 4. Cell recovery by using vitrification. Figure 4 illustrates cell recovery of vitrified MHEC-5Ts immediately after thawing. DMEM = DMEM Ham's F-12 Gln+, 20% FBS, and 3M of one of the CPAs; K⁺TiP = K⁺TiP. 0.5M sucrose, 20% FBS, and 3M of one of the CPAs; * = significant ($p < 0.05$); **** = highest significance ($p < 0.0001$).

doi:10.1371/journal.pone.0149660.g004

highest absorbance after the thawing of cells frozen by utilizing K⁺TiP as the cryopreservation medium was $2.83 \pm 0.16 \times 10^{-7}$ au ($268.2 \pm 15\%$) with DMSO, and then $2.28 \pm 0.26 \times 10^{-7}$ au ($216.1 \pm 25\%$) with GLY, $0.24 \pm 0.05 \times 10^{-7}$ au ($22.5 \pm 4.9\%$) with PG, and $0.1 \pm 0.03 \times 10^{-7}$ au ($9.2 \pm 2.8\%$) with EG as the CPA.

Three days after thawing, the highest absorbance of cells frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was $2.93 \pm 0.25 \times 10^{-7}$ au ($277.6 \pm 23.2\%$ relative to unfrozen control at day 0) with DMSO, and then $0.98 \pm 0.46 \times 10^{-7}$ au ($92.7 \pm 44\%$) with GLY, $0.27 \pm 0.11 \times 10^{-7}$ au ($26.1 \pm 10\%$) with EG, and $0.12 \pm 0.01 \times 10^{-7}$ au ($11.4 \pm 1.3\%$) with PG as the CPA. The highest absorbance after the thawing of cells frozen by utilizing K⁺TiP as the cryopreservation medium was $3.25 \pm 0.36 \times 10^{-7}$ au ($308.2 \pm 34.4\%$) with GLY, and then $2.65 \pm 0.16 \times 10^{-7}$ au ($250.6 \pm 14.9\%$) with DMSO, $0.28 \pm 0.09 \times 10^{-7}$ au ($26.6 \pm 8.1\%$) with PG, and $0.15 \pm 0.07 \times 10^{-7}$ au ($14 \pm 7\%$) with EG as the CPA. Fig 5 illustrates these findings.

The fastest MHEC-5T doubling-time of cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium was 26 ± 7.3 h with GLY, and then 42 ± 2.4 h with DMSO, 56 ± 19.5 h with EG, and 128 ± 6.9 h with PG as the CPA. The doubling-time of cells frozen by utilizing K⁺TiP as the cryopreservation medium was 18 ± 4.2 h with GLY, 53 ± 2.6 h with DMSO, 87 ± 28.7 h with EG, and 96 ± 12.2 h with PG as the CPA. Fig 6 illustrates these findings.

All frozen cells could be cultured until confluence in the 25 cm² cell culture flasks after thawing. Confluence was reached as early as day 3 by cells previously frozen by utilizing K⁺TiP as the cryopreservation medium and DMSO as the CPA. On day 4, confluence was reached by




Fig 5. Viable cells in proliferation after vitrification. The CellTiter 96[®] A_{QUEOUS} One Solution Cell Proliferation Assay (COS, Promega Corporation, Mannheim, Germany) was used to determine viable cells in proliferation at one hour after thawing and at day one, two, and three. Figure 5 illustrates the absorption of viable cells in proliferation after vitrification. DMEM = DMEM Ham's F-12 Gln+, 20% FBS, and 3M of one of the CPAs; K⁺TiP = K⁺TiP. 0.5M sucrose, 20% FBS, and 3M of one of the CPAs; DMSO = dimethyl sulfoxide; EG = ethylene glycol; PG = propylene glycol; GLY = glycerin; * = significant ($p < 0.05$); ** = medium significant ($p < 0.01$); au = absorbance units.

doi:10.1371/journal.pone.0149660.g005

cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and DMSO as the CPA and by utilizing K⁺TiP as the cryopreservation medium and GLY as the CPA. On day 6, confluence was reached by cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and GLY as the CPA. On day 8, confluence was reached by cells previously frozen by utilizing K⁺TiP as the cryopreservation medium and EG as the CPA. On day 9, confluence was reached by cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and EG as the CPA. On day 10, confluence was reached by cells previously frozen by utilizing K⁺TiP as the cryopreservation medium and PG as the CPA. Only on day 18 was confluence reached by cells previously frozen by utilizing DMEM Ham's F-12 Gln+ as the cryopreservation medium and PG as the CPA. Fig 7 illustrates these findings.

Discussion

Our aim was to compare the promising but sparsely examined technique of vitrification with the standard slow freezing technique. To optimize the media used in vitrification, we




Fig 6. Cell doubling-time after vitrification. Figure 6 illustrates MHEC-5T doubling-time after vitrification. DMEM = DMEM Ham's F-12 Gln+, 20% FBS, and 3M of one of the CPAs; K⁺TiP = K⁺TiP. 0.5M sucrose, 20% FBS, and 3M of one of the CPAs; * = significant (p < 0.05); *** = highly significant (p < 0.001); **** = highest significance (p < 0.0001).

doi:10.1371/journal.pone.0149660.g006

introduced a modified organ preservation solution that showed excellent results in the prevention of iron-dependent cold-induced injury during the cold storage of vessels. K⁺TiP contains 131 mM potassium in order to terminate the membrane resting potential during freezing. With a terminated resting potential, the O₂ consumption of the cells decreases [18]. Wille et al. have reported significantly lower damage to endothelial cells by using potassium as the main cation, instead of sodium [19]. High extracellular potassium is responsible for the prevention of the K⁺-outflow of the cell [20, 21], which leads to its apoptosis [22]. The pH of K⁺TiP was adjusted to 7.0, as a mild acidosis is protective against multiple mechanisms of cellular damage [23].

TiProtec (Dr. Franz Köhler Chemie GmbH, Bensheim, Germany) organ storage solution contains several ingredients that counteract the damage mechanism of cold storage. TiProtec was used as the basis for K⁺TiP. It contains the following additions:

Several studies have shown that the cold-induced injury of several cultured cell types is mediated via the iron-dependent formation of reactive oxygen species (ROS). Chelatable “redox-active” iron plays the major role in the development of cold-induced apoptosis [24, 25]. To prevent cold-induced injury, the hydroxamic acid derivative LK 614 is therefore included in TiProtec. The aromatic ring system gives the molecule a sufficiently high lipophilicity to permeate membranes. Glucose is added as a substrate for the glycolytically highly active endothelial cells [26]. To prevent the formation of non-specific pores during hypoxia followed by sodium influx, TiProtec contains the amino acid glycine and the structurally related alanine

The cryopreservation process involving vitrification is fast and reliable. The precooled DMEM Ham's F-12 Gln+ is added to the centrifuged cells in a cryo vial, followed by the prepared and precooled VS. The mixture is pipetted up and down two to three times to ensure good mixing, and then the cryo vial is tightly capped and directly plunged into LN₂. This process can be conducted within less than 15 seconds per vial and can be performed by various researchers within the same time span. For slow freezing, the mixing process is the same, but the first prepared vial needs to rest in the CoolCell or on ice until the last one is ready, and everything can be stored in -20°C.

Cell survival is normally evaluated by using live/dead staining with membrane-permeant molecules that are cleaved by esterases in live cells to yield cytoplasmic fluorescence and the membrane-impermeant dye, propidium iodide. Because propidium iodide is cell-toxic, this method is highly time-dependent, not very sensitive, and difficult to reproduce. Therefore, we developed and used a new time-independent and cell-count-adjusted method to evaluate cell survival and the number of viable cells in proliferation after thawing.

Because of the large variability of the amount of cells that can be plated into a particular well or cryo vial, we decided first to minimize this error by repetitive gentle vortex of the cell suspension prior to the plating of each well; to this end, we used a multistep pipette (Eppendorf repeater stream combitip advanced, Hamburg, Germany). Our results indicated, that in spite of great care being taken, with repetitive gentle vortexing and the use of a multistep pipette, the number of cells per cryo vial varied between 279,000 and 590,000 (median 461,000, standard deviation 67,000, $n = 32$). To detect the exact cell number per cryo vial, the DNA content was measured according to Kim et al. by using Hoechst 33258 dye. This assay is stable for at least 2 h and sensitive to as little as 6 ng of DNA or equivalently less than 1,000 cells [17]. We finally evaluated not the simple results achieved from CellTiter 96® A_{QUEOUS} One Solution Cell Proliferation Assay, but those normalized by building a quotient of fluorescence intensity and cell number per cryo vial. The result is a ratio of measured absorbance that is directly proportional to the number of living cells in culture compared with the total cell number.

In slow freezing, we used a CoolCell, because it precisely controls air flow and heat removal to ensure standardized cell freezing runs and no variability in the cryopreservation period. The elimination of alcohol from the cryopreservation process means no maintenance, no ongoing expense, no waste, and no variability [30].

After use of the standardized controlled-rate -1°C/minute protocol, the cell recovery rates after thawing could be ranked in the following order with respect to CPA GLY < EG ≈ PG < DMSO. The use of DMEM Ham's F-12 Gln+ as the cryopreservation medium showed a statistical benefit compared with K⁺TiP with DMSO or PG as the CPA. Vitrification gave the following order with regard to cell recovery rates after thawing: PG < EG << GLY ≈ DMSO. In this case, the use of K⁺TiP as the cryopreservation medium showed a statistical benefit compared with DMEM Ham's F-12 Gln+ with DMSO and particularly PG and GLY as CPA. Finally, the best cryopreservation method was vitrification with VS containing K⁺TiP, 40% FBS, 0.5 M sucrose, and 6M CPA. With regard to cell recovery immediately after thawing, DMSO was the most suitable CPA (99 ± 0.5%). The most viable cells in proliferation after three days of culture could be achieved by using GLY (308 ± 34%). However, against expectation, K⁺TiP had no benefit in the slow freezing methods. We conclude that, only in vitrification, is the termination of membrane resting potential beneficial. In slow cooling, the termination of the membrane resting potential is harmful to cells, and therefore, GLY as a low-toxic CPA, shows good long-term results when used in slow freezing technology with K⁺TiP as the cryopreservation medium. The preservation of surface markers is possibly better with low-toxic CPAs. To clarify this, further studies with fluorescence-activated cell sorting analysis are necessary. Additional investigation is also needed to determine whether this effect is influenced by

the potassium concentration or another modification made compared with DMEM Ham's F-12 Gln+. The various methods of assessment of recovery and the various cryopreservation media used by other groups make direct quantitative comparisons of the various results difficult or even impossible.

Supporting Information

S1 Data Set. Minimal data set. The minimal data set includes the following files: results_freezing.xls, results_vitrific.xls and results_cells_25cm2_counted.xls. (ZIP)

Acknowledgments

The authors wish to thank Monika Jerg, Katja Urlbauer, and Gabriela Cudek for their excellent technical support.

Author Contributions

Conceived and designed the experiments: AvB NR. Performed the experiments: AvB AE. Analyzed the data: AvB AE LR. Contributed reagents/materials/analysis tools: AvB AE SS. Wrote the paper: AvB.

References

1. Verma S, Anderson TJ. Fundamentals of endothelial function for the clinical cardiologist. *Circulation*. 2002; 105(5):546–9. PMID: [11827916](#).
2. Lehr HA, Bittinger F, Kirkpatrick CJ. Microcirculatory dysfunction in sepsis: a pathogenetic basis for therapy? *The Journal of pathology*. 2000; 190(3):373–86. doi: [10.1002/\(SICI\)1096-9896\(200002\)190:3<373::AID-PATH593>3.0.CO;2-3](#) PMID: [10685071](#).
3. Kirkpatrick CJ, Bittinger F, Klein CL, Hauptmann S, Klosterhalfen B. The role of the microcirculation in multiple organ dysfunction syndrome (MODS): a review and perspective. *Virchows Archiv: an international journal of pathology*. 1996; 427(5):461–76. PMID: [8624575](#).
4. Krump-Konvalinkova V, Bittinger F, Unger RE, Peters K, Lehr HA, Kirkpatrick CJ. Generation of human pulmonary microvascular endothelial cell lines. *Laboratory investigation; a journal of technical methods and pathology*. 2001; 81(12):1717–27. PMID: [11742042](#).
5. Katkov II, Kan NG, Cimadamore F, Nelson B, Snyder EY, Tersikh AV. DMSO-Free Programmed Cryopreservation of Fully Dissociated and Adherent Human Induced Pluripotent Stem Cells. *Stem cells international*. 2011; 2011:981606. doi: [10.4061/2011/981606](#) PMID: [21716669](#); PubMed Central PMCID: PMC3118298.
6. Polge C, Smith AU, Parkes AS. Revival of spermatozoa after vitrification and dehydration at low temperatures. *Nature*. 1949; 164(4172):666. PMID: [18143360](#).
7. Lovelock JE, Bishop MW. Prevention of freezing damage to living cells by dimethyl sulphoxide. *Nature*. 1959; 183(4672):1394–5. PMID: [13657132](#).
8. Meryman HT. Freezing injury and its prevention in living cells. *Annual review of biophysics and bioengineering*. 1974; 3(0):341–63. doi: [10.1146/annurev.bb.03.060174.002013](#) PMID: [4609398](#).
9. Moon JH, Lee JR, Jee BC, Suh CS, Kim SH, Lim HJ, et al. Successful vitrification of human amnion-derived mesenchymal stem cells. *Human reproduction*. 2008; 23(8):1760–70. doi: [10.1093/humrep/den202](#) PMID: [18541648](#).
10. Janz Fde L, Debes Ade A, Cavaglieri Rde C, Duarte SA, Romao CM, Moron AF, et al. Evaluation of distinct freezing methods and cryoprotectants for human amniotic fluid stem cells cryopreservation. *Journal of biomedicine & biotechnology*. 2012; 2012:649353. doi: [10.1155/2012/649353](#) PMID: [22665987](#); PubMed Central PMCID: PMC3361720.
11. Davis JM, Rowley SD, Braine HG, Piantadosi S, Santos GW. Clinical toxicity of cryopreserved bone marrow graft infusion. *Blood*. 1990; 75(3):781–6. PMID: [2297578](#).
12. Syme R, Bewick M, Stewart D, Porter K, Chadderton T, Gluck S. The role of depletion of dimethyl sulphoxide before autografting: on hematologic recovery, side effects, and toxicity. *Biology of blood and*

- marrow transplantation: journal of the American Society for Blood and Marrow Transplantation. 2004; 10(2):135–41. doi: [10.1016/j.bbmt.2003.09.016](https://doi.org/10.1016/j.bbmt.2003.09.016) PMID: [14750079](https://pubmed.ncbi.nlm.nih.gov/14750079/).
13. Fan WX, Ma XH, Ge D, Liu TQ, Cui ZF. Cryoprotectants for the vitrification of corneal endothelial cells. *Cryobiology*. 2009; 58(1):28–36. doi: [10.1016/j.cryobiol.2008.10.124](https://doi.org/10.1016/j.cryobiol.2008.10.124) PMID: [18976647](https://pubmed.ncbi.nlm.nih.gov/18976647/).
 14. Kawakami Y, Ishihara M, Saito T, Fujimoto T, Adachi S, Arai K, et al. Cryopreservation of green fluorescent protein (GFP)-labeled primordial germ cells with GFP fused to the 3' untranslated region of the nanos gene by vitrification of Japanese eel (*Anguilla japonica*) somite stage embryos. *Journal of animal science*. 2012; 90(12):4256–65. doi: [10.2527/jas.2011-4884](https://doi.org/10.2527/jas.2011-4884) PMID: [22829617](https://pubmed.ncbi.nlm.nih.gov/22829617/).
 15. Sheffen B, Van Der Zwalm P., and Massip A. A simple and efficient procedure for preservation of mouse embryos by vitrification. *Cryo Letters*. 1986; 7:260–9.
 16. Berridge MV, Tan AS. Characterization of the cellular reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT): subcellular localization, substrate dependence, and involvement of mitochondrial electron transport in MTT reduction. *Archives of biochemistry and biophysics*. 1993; 303(2):474–82. doi: [10.1006/abbi.1993.1311](https://doi.org/10.1006/abbi.1993.1311) PMID: [8390225](https://pubmed.ncbi.nlm.nih.gov/8390225/).
 17. Kim YJ, Sah RL, Doong JY, Grodzinsky AJ. Fluorometric assay of DNA in cartilage explants using Hoechst 33258. *Analytical biochemistry*. 1988; 174(1):168–76. PMID: [2464289](https://pubmed.ncbi.nlm.nih.gov/2464289/).
 18. Preusse CJ, Gebhard MM, Bretschneider HJ. Myocardial "equilibration processes" and myocardial energy turnover during initiation of artificial cardiac arrest with cardioplegic solution—reasons for a sufficiently long cardioplegic perfusion. *The Thoracic and cardiovascular surgeon*. 1981; 29(2):71–6. doi: [10.1055/s-2007-1023446](https://doi.org/10.1055/s-2007-1023446) PMID: [6166077](https://pubmed.ncbi.nlm.nih.gov/6166077/).
 19. Wille T, de Groot H, Rauen U. Improvement of the cold storage of blood vessels with a vascular preservation solution. Study in porcine aortic segments. *Journal of vascular surgery*. 2008; 47(2):422–31. doi: [10.1016/j.jvs.2007.09.048](https://doi.org/10.1016/j.jvs.2007.09.048) PMID: [18164170](https://pubmed.ncbi.nlm.nih.gov/18164170/).
 20. Collins GM, Bravo-Shugarman M, Terasaki PI. Kidney preservation for transportation. Initial perfusion and 30 hours' ice storage. *Lancet*. 1969; 2(7632):1219–22. PMID: [4187813](https://pubmed.ncbi.nlm.nih.gov/4187813/).
 21. D'Alessandro AM, Southard JH, Love RB, Belzer FO. Organ preservation. *The Surgical clinics of North America*. 1994; 74(5):1083–95. PMID: [7940062](https://pubmed.ncbi.nlm.nih.gov/7940062/).
 22. Lang F, Foller M, Lang KS, Lang PA, Ritter M, Gulbins E, et al. Ion channels in cell proliferation and apoptotic cell death. *The Journal of membrane biology*. 2005; 205(3):147–57. doi: [10.1007/s00232-005-0780-5](https://doi.org/10.1007/s00232-005-0780-5) PMID: [16362503](https://pubmed.ncbi.nlm.nih.gov/16362503/).
 23. Nishimura Y, Romer LH, Lemasters JJ. Mitochondrial dysfunction and cytoskeletal disruption during chemical hypoxia to cultured rat hepatic sinusoidal endothelial cells: the pH paradox and cytoprotection by glucose, acidotic pH, and glycine. *Hepatology*. 1998; 27(4):1039–49. doi: [10.1002/hep.510270420](https://doi.org/10.1002/hep.510270420) PMID: [9537444](https://pubmed.ncbi.nlm.nih.gov/9537444/).
 24. Rauen U, Kerkweg U, de Groot H. Iron-dependent vs. iron-independent cold-induced injury to cultured rat hepatocytes: a comparative study in physiological media and organ preservation solutions. *Cryobiology*. 2007; 54(1):77–86. doi: [10.1016/j.cryobiol.2006.11.008](https://doi.org/10.1016/j.cryobiol.2006.11.008) PMID: [17289012](https://pubmed.ncbi.nlm.nih.gov/17289012/).
 25. Rauen U, Polzar B, Stephan H, Mannherz HG, de Groot H. Cold-induced apoptosis in cultured hepatocytes and liver endothelial cells: mediation by reactive oxygen species. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology*. 1999; 13(1):155–68. PMID: [9872940](https://pubmed.ncbi.nlm.nih.gov/9872940/).
 26. Mertens S, Noll T, Spahr R, Krutzfeldt A, Piper HM. Energetic response of coronary endothelial cells to hypoxia. *The American journal of physiology*. 1990; 258(3 Pt 2):H689–94. PMID: [2316683](https://pubmed.ncbi.nlm.nih.gov/2316683/).
 27. Rauen U, de Groot H. New insights into the cellular and molecular mechanisms of cold storage injury. *Journal of investigative medicine: the official publication of the American Federation for Clinical Research*. 2004; 52(5):299–309. PMID: [15551652](https://pubmed.ncbi.nlm.nih.gov/15551652/).
 28. Wu S, Wohlschlaeger J, de Groot H, Rauen U. Evaluation of a modified HTK solution containing the new iron chelator LK 614 in an isolated rat liver perfusion model. *Journal of investigative surgery: the official journal of the Academy of Surgical Research*. 2009; 22(5):340–7. PMID: [19842888](https://pubmed.ncbi.nlm.nih.gov/19842888/).
 29. Sutton RL. Critical cooling rates for aqueous cryoprotectants in the presence of sugars and polysaccharides. *Cryobiology*. 1992; 29(5):585–98. PMID: [1424715](https://pubmed.ncbi.nlm.nih.gov/1424715/).
 30. Yokoyama WM, Thompson ML, Ehrhardt RO. Cryopreservation and thawing of cells. *Current protocols in immunology* / edited by Coligan John E [et al]. 2012; Appendix 3:3G. doi: [10.1002/0471142735.ima03gs99](https://doi.org/10.1002/0471142735.ima03gs99) PMID: [23129152](https://pubmed.ncbi.nlm.nih.gov/23129152/).