


# Genome-wide association study identified novel candidate loci affecting wood formation in Norway spruce

John Baisou<sup>1,\*</sup> , Amaryllis Vidalis<sup>2</sup>, Linghua Zhou<sup>1</sup>, Zhi-Qiang Chen<sup>1</sup>, Zitong Li<sup>3</sup>, Mikko J. Sillanpää<sup>4</sup>, Carolina Bernhardsson<sup>1,5</sup>, Douglas Scofield<sup>6</sup>, Nils Forsberg<sup>1</sup>, Thomas Grahn<sup>7</sup>, Lars Olsson<sup>7</sup>, Bo Karlsson<sup>8</sup>, Harry Wu<sup>1</sup>, Pär K. Ingvarsson<sup>5,9</sup>, Sven-Olof Lundqvist<sup>7,10</sup>, Totte Niittylä<sup>1,†</sup> and M Rosario García-Gil<sup>1,\*,†</sup>

<sup>1</sup>Department of Forest Genetics and Plant Physiology, Umeå Plant Science Centre, Swedish University of Agricultural Science, Parallellvägen 21, Umeå 907 36, Sweden,

<sup>2</sup>Section of Population Epigenetics and Epigenomics, Centre of Life and Food Sciences Weihenstephan, Technische Universität München, Lichtenbergstr. 2a, München 85748, Germany,

<sup>3</sup>Ecological Genetics Research Unit, Department of Biosciences, University of Helsinki, P.O. Box 65, FI-00014, Helsinki, Finland,

<sup>4</sup>Department of Mathematical Sciences, Biocenter Oulu, University of Oulu, Pentti Kaiteran katu 1, Oulu, Finland,

<sup>5</sup>Department of Ecology and Environmental Science, Umeå University, Linnaeus väg 4-6, Umeå 907 36, Sweden,

<sup>6</sup>Uppsala Multidisciplinary Centre for Advanced Computational Science, Uppsala University, Lägerhyddsvägen 2, Uppsala 752 37, Sweden,

<sup>7</sup>RISE Bioeconomy, Drottning Kristinas väg 61, SE-114 86, Stockholm, Sweden,

<sup>8</sup>Skogforsk, Ekebo 2250, SE-268 90, Svalöv, Sweden,

<sup>9</sup>Department of Ecology and Genetics: Evolutionary Biology, Uppsala University, Kåbovägen 4, Uppsala 752 36, Sweden, and

<sup>10</sup>IIC, Rosenlundsgatan 48B, SE-118 63, Stockholm, Sweden

Received 1 March 2019; revised 16 April 2019; accepted 20 May 2019; published online 5 June 2019.

\*For correspondence (e-mails m.rosario.garcia@slu.se; john.baisou@slu.se).

†Double last authorship.

## SUMMARY

Norway spruce is a boreal forest tree species of significant ecological and economic importance. Hence there is a strong imperative to dissect the genetics underlying important wood quality traits in the species. We performed a functional genome-wide association study (GWAS) of 17 wood traits in Norway spruce using 178 101 single nucleotide polymorphisms (SNPs) generated from exome genotyping of 517 mother trees. The wood traits were defined using functional modelling of wood properties across annual growth rings. We applied a Least Absolute Shrinkage and Selection Operator (LASSO-based) association mapping method using a functional multilocus mapping approach that utilizes latent traits, with a stability selection probability method as the hypothesis testing approach to determine a significant quantitative trait locus. The analysis provided 52 significant SNPs from 39 candidate genes, including genes previously implicated in wood formation and tree growth in spruce and other species. Our study represents a multilocus GWAS for complex wood traits in Norway spruce. The results advance our understanding of the genetics influencing wood traits and identifies candidate genes for future functional studies.

**Keywords:** candidate genes, functional trait mapping, genome-wide association mapping, Norway spruce, sequence capture, single nucleotide polymorphisms.

## INTRODUCTION

Norway spruce (*Picea abies* (L.) Karst.) is a dominant boreal species of significant economic and ecological importance (Hannrup *et al.*, 2004). Long-term Norway spruce breeding programmes for improvement of growth and survival were initiated in the 1940s and recently, wood quality

has become one of the priority traits (Bertaud and Holmbom, 2004; Hannrup *et al.*, 2004). Norway spruce breeding in Sweden completes one cycle in about 20 years and such long generation time makes improvements in growth and wood quality very slow. Among wood quality traits, wood

density is considered a key indicator of stability, strength and stiffness of sawn timber (Hauksson *et al.*, 2001). Several studies of wood quality observed that fast growth conflicts with high quality wood, as shown by the negative genetic correlation between wood volume growth and density in Norway spruce (Olesen, 1977; Dutilleul *et al.*, 1998; Chen *et al.*, 2014). However in several conifers such as Scots pine (*P. sylvestris* L.) and red pine the relationship has been inconsistent (Larocque and Marshall, 1995; Peltoja *et al.*, 2009). To combine fast growth and desirable wood properties through breeding, and to shorten the breeding cycle, it is therefore imperative to design effective early selection methods and breeding strategies. In an effort to design optimal breeding and selection strategies it is essential to identify alleles that are responsible for generating favourable or unfavourable genetic correlations (Hallingbäck *et al.*, 2014).

One of the early studies in conifers identified quantitative trait locus (QTLs) for wood density variation in loblolly pine using linkage analyses based on segregating family pedigrees (Groover *et al.*, 1994). However, marker-aided selection (MAS) based on results from QTL linkage analyses were never implemented in practical tree breeding due to the so-called Beavis effect (e.g. inflated estimates of allelic effects and underestimation of QTL number for economically important traits) (Beavis, 1998), inconsistent associations among different families and the low transferability of markers (Strauss *et al.*, 1992). Association mapping (AM), also called linkage disequilibrium (LD) mapping, is a powerful alternative QTL detection method that was introduced to tree genetics using a candidate gene approach (Thumma *et al.*, 2010). AM overcomes the limited resolution of family-based QTL linkage mapping by relying on historical recombination in the mapping population (Neale and Savolainen, 2004; Thavamanikumar *et al.*, 2013; Huang and Han, 2014). However, the genome-wide levels of LD in Norway spruce has been revealed to be complex and highly heterogeneous (Larsson *et al.*, 2013). Therefore, AM is also vulnerable to some confounding historical factors such as population admixture, selection pressures which include possible genetic drift. Therefore, population genetic structure, kinship and LD within the study population need to be carefully accounted for in the analysis to minimize false positives (Khan and Korban, 2012).

The availability of a draft genome sequence for Norway spruce (Nystedt *et al.*, 2013) has provided numerous possibilities for the development of genetic markers to conduct both AM at the genome-wide level (genome-wide association, GWAS) and genomic selection (GS). Several reduced representation-based approaches such as sequence capture and transcriptome sequencing (Hirsch *et al.*, 2014) have been developed for studying large genomes, such as the 20 Gb Norway spruce genome. These approaches

reduce the sequence space by decreasing the repetitive sequence content of the genome.

Several AM studies have been performed on traits in different tree species and have identified genetic loci linked to, for instance, wood properties in *Populus trichocarpa* Torr. & A. Gray ex. Hook (Porth *et al.*, 2013) and *Eucalyptus* (Resende *et al.*, 2017b), and adaptive traits in *Pinus contorta* Douglas (Parchman *et al.*, 2012). Some genes may impact the trait development at a particular developmental stage, whereas others may alter, or control, rates of change and transitions between consecutive stages (Xing *et al.*, 2012; Anderegg, 2015). Studies aimed at dissecting the genetic basis of such dynamics in wood properties can benefit from the application of mathematical functions that account for year-to-year variation across annual growth rings, cambial age and distance from pith (Li *et al.*, 2014). The development of mathematical methods for the analysis of these longitudinal traits has made it possible to map QTLs underlying the dynamics of developmental traits (Yang *et al.*, 2006; Li and Sillanpää, 2013; Camargo *et al.*, 2018), and to enhance our understanding of the genetic architecture of the growth trajectories of such dynamic traits (Ma *et al.*, 2002; Xing *et al.*, 2012). Such functional mapping analysis can be conducted using a multistage approach (Heuven and Janss, 2010). First, the phenotype trends of each individual are modelled using curve-fitting methods and the parameters describing the curve are then considered as latent traits. The latent traits are then used in an independent association analyses to search for genomic regions affecting the trait and to estimate genetic marker effects (Li and Sillanpää, 2013; Li *et al.*, 2014).

In this study, we applied a functional AM approach to identify genomic regions contributing to wood quality traits in Norway spruce. We applied spline models since traditional analyses that utilise a single point data across annual growth rings may confound the analyses by averaging across a full sample. Such averaging may obscure mechanisms acting at specific time points during wood formation and will make identification of underlying genes more difficult. This study has performed the analysis of number of cells per ring calculated from SilviScan data. Penalized LASSO regression (Tibshirani, 1996) and the stability selection probability method (Meinshausen and Bühlmann, 2010) were then used, to detect significant associations between latent traits derived from estimated breeding values (EBVs) and 178101 SNP markers covering the Norway spruce genome.

## RESULTS AND DISCUSSION

All 517 Norway spruce maternal trees in the study were considered for variant detection and an average of 1.5 million paired-end reads were sequenced per individual resulting in 178 101 SNPs. Most SNPs were missense (61%). Applying the probability of stability selection

(SSP) to the intercept, slope and two nodes ( $\beta_2$  and  $\beta_3$ ) we detected 52 significant QTLs in 17 individual traits whose phenotypic variance explained (PVE) ranged from 0.01 to 4.93% (Table 1). 14 of the significant markers

were consistent with overdominance ( $|d/a| > 1.25$ ), with the remaining being codominant (27) ( $|d/a| < 0.50$ ) and 10 exhibiting partial to full dominance ( $0.50 < |d/a| < 1.25$ ) (Table 2, Figure 3).

**Table 1** Phenotypes, latent traits, SNP, SNP feature, frequency and PVE

Phenotype	Latent Trait	QTL	SNP <sup>a</sup>	Allele	SNP Feature	Frequency	PVE (%)
WD	Intercept	167610	MA_10435406_13733	A/G	Downstream variant	0.71	4.64
	Slope	30469	MA_33109_11804	A/G	Upstream variant	0.72	4.50
	$\beta_2$	30469	MA_33109_11804	A/G	Upstream variant	0.551	4.15
	$\beta_3$	157442	MA_10432646_63090	G/A	Upstream variant	0.567	2.43
EWD	Intercept	167610	MA_10435406_13733	A/G	Downstream variant	0.545	3.38
	Slope	23798	MA_20321_44812	C/T	Upstream variant	0.53	0.69
		70955	MA_118446_4316	T/A	Upstream variant	0.644	0.40
TWD	Slope	131698	MA_10235390_3386	G/A	Stop gained	0.672	1.58
		160208	MA_10433411_3386	T/C	Intron variant	0.595	3.41
	$\beta_2$	89044	MA_212523_6278	T/C	Upstream variant	0.534	3.34
LWD	Slope	43797	MA_62987_13474	T/C	Missense variant	0.524	1.81
		165481	MA_10434805_21408	C/T	Intron variant	0.588	1.21
		171223	MA_10436058_4902	G/A	Intron variant	0.712	4.03
RW	Intercept	11535	MA_10694_9101	A/C	Synonymous variant	0.545	1.95
		112391	MA_879270_7373	C/T,A	Stop gained	0.532	1.45
		112394	MA_879384_3894	C/A	Splice region variant	0.692	2.56
	Slope	165481	MA_10434805_21408	C/T	Intron variant	0.521	2.66
		23808	MA_20322_28351	T/G	Synonymous variant	0.554	1.78
	$\beta_2$	165481	MA_10434805_21408	C/T	Intron variant	0.533	0.18
		23808	MA_20322_28351	T/G	Synonymous variant	0.55	1.20
TRW	Slope	165481	MA_10434805_21408	C/T	Intron variant	0.615	1.79
		111057	MA_817099_1105	T/A	Missense variant	0.685	1.12
	$\beta_2$	33110	MA_38472_13803	T/A	Upstream gene variant	0.657	3.23
		89295	MA_214776_1624	G/A	Upstream gene variant	0.688	4.51
$\beta_3$	111057	MA_817099_1105	T/A	Missense variant	0.672	1.20	
LRW	Intercept	143628	MA_10428744_29330	C/T	Downstream variant	0.668	0.5
		164772	MA_10434624_20686	C/A	Downstream variant	0.571	0.06
MOE	Slope	165481	MA_10434805_21408	C/T	Intron variant	0.602	1.00
NC	$\beta_2$	145839	MA_10429444_12692	G/C	Upstream variant	0.645	3.82
ENC	Slope	98508	MA_402880_2045	A/C	Upstream variant	0.667	0.03
		167610	MA_10435406_13733	A/G	Downstream variant	0.685	0.01
TNC	Intercept	95870	MA_346723_2241	T/C	Upstream variant	0.667	3.78
		126785	MA_9447489_687	A/C	Upstream gene variant	0.68	4.93
LNC	Intercept	143628	MA_10428744_29330	C/T	Downstream variant	0.66	3.14
		143628	MA_10428744_29330	C/T	Downstream variant	0.672	4.77
EP	Intercept	16868	MA_15729_40331	G/T	Intron variant	0.609	3.32
		91242	MA_246125_1213	G/A	Synonymous variant	0.594	3.41
TP	Intercept	101203	MA_462319_4322	A/C	Upstream gene variant	0.594	1.16
		132014	MA_10251995_2442	A/C	Upstream gene variant	0.601	3.22
LP	$\beta_2$	162397	MA_10434007_77578	C/T	Upstream gene variant	0.892	1.14
EP/LP	Intercept	51657	MA_80954_29644	G/A	Downstream variant	0.63	0.81
		60787	MA_98424_947	C/T	Intron variant	0.655	1.80
		123639	MA_8790100_1384	A/C	Upstream variant	0.628	0.75
	$\beta_2$	59480	MA_96191_7122	A/G	Synonymous	0.6	2.37
		117333	MA_1045136_4310	T/C	Missense variant	0.523	1.34
	Mass index (growth × density)	Intercept	166235	MA_10435002_4986	G/A	Intergenic variant	0.533
61096			MA_99004_17108	G/A	Synonymous variant	0.66	0.01
Slope		67181	MA_109804_10278	G/A	Missense variant	0.612	0.05
		1401	MA_1378_4718	C/A	Exon/stop gained	0.588	1.19
		138744	MA_10427214_13968	G/T	Missense variant	0.58	1.80
$\beta_2$		162397	MA_10434007_77578	C/T	Upstream variant	0.627	1.44
		21924	MA_19222_1789	A/G	Upstream variant	0.71	1.82

<sup>a</sup>SNP: The SNP name was composed of the contig (MA\_number) and SNP position on contig. For example, the first SNP MA\_1043540\_13733 was located on contig MA\_1043540 at position 13 733 bp; PVE is the phenotypic variance explained.

**Table 2** SNP modes of inheritance

Phenotype	QTL	SNP	Allele	2a <sup>a</sup>	d <sup>b</sup>	d/a
WD	167610	MA_10435406_13733	A/G	19.63	9.45	0.96
	30469	MA_33109_11804	A/G	5.19	6.22	2.40
	157442	MA_10432646_63090	G/A	4.49	2.80	1.25
EWD	167610	MA_10435406_13733	A/G	5.495	2.270	0.83
	23798	MA_20321_44812	C/T	4.814	5.905	2.45
	70955	MA_118446_4316	T/A	1.989	0.966	0.97
TWD	131698	MA_10235390_3386	G/A	3.849	-0.636	-0.33
	160208	MA_10433411_3386	T/C	2.313	3.908	3.38
	89044	MA_212523_6278	T/C	0.703	-0.962	-2.73
LWD	43797	MA_62987_13474	T/C	5.124	-1.086	-0.42
	165481	MA_10434805_21408	C/T	4.684	1.165	0.50
	171223	MA_10436058_4902	G/A	0.938	2.482	5.29
RW	11535	MA_10694_9101	A/C	0.111	0.049	0.88
	112391	MA_879270_7373	C/T,A	0.056	-0.027	-0.98
	112394	MA_879384_3894	C/A	0.194	-0.045	-0.45
	165481	MA_10434805_21408	C/T	0.158	0.039	0.49
	23808	MA_20322_28351	T/G	0.025	0.030	2.62
TRW	111057	MA_817099_1105	T/A	0.016	0.001	0.16
	33110	MA_38472_13803	T/A	0.029	-0.002	-0.19
	89295	MA_214776_1624	G/A	0.026	-0.001	-0.13
LRW	143628	MA_10428744_29330	C/T	0.006	-0.002	-0.67
	164772	MA_10434624_20686	C/A	0.002	0.003	2.90
MOE	165481	MA_10434805_21408	C/T	0.376	0.101	0.53
NC	145839	MA_10429444_12692	G/C	0.298	0.792	5.31
ENC	98508	MA_402880_2045	A/C	4.144	1.314	0.63
	167610	MA_10435406_13733	A/G	4.695	-3.033	-1.29
TNC	95870	MA_346723_2241	T/C	0.529	-0.187	-0.71
	126785	MA_9447489_687	A/C	0.083	-0.429	-10.21
LNC	143628	MA_10428744_29330	C/T	0.219	-0.057	-0.52
EP	16868	MA_15729_40331	G/T	0.542	0.149	0.55
	91242	MA_246125_1213	G/A	0.183	-0.129	-1.40
TP	101203	MA_462319_4322	A/C	0.469	-0.199	-0.85
	132014	MA_10251995_2442	A/C	0.339	-0.429	-1.63
LP	162397	MA_10434007_77578	C/T	0.127	-0.071	-1.11
EP/LP	51657	MA_80954_29644	G/A	0.081	0.062	1.49
	60787	MA_98424_947	C/T	0.254	-0.181	-1.43
	123639	MA_8790100_1384	A/C	0.032	-0.078	-4.81
	59480	MA_96191_7122	A/G	0.120	-0.117	-1.95
	117333	MA_1045136_4310	T/C	0.018	0.077	8.56
	166235	MA_10435002_4986	G/A	0.138	0.013	0.19
	61096	MA_99004_17108	G/A	0.006	-0.009	-3.16
	67181	MA_109804_10278	G/A	0.007	-0.012	-3.14
	1401	MA_1378_4718	C/A	0.003	0.004	2.67
	138744	MA_10427214_13968	G/T	0.002	0.017	17.00
MI	162397	MA_10434007_77578	C/T	0.025	-0.010	-0.79
	21924	MA_19222_1789	A/G	0.014	-0.008	-1.14

<sup>a</sup>Calculated as the difference between the phenotype means observed within each homozygous class ( $2a = |G_{BB} - G_{bb}|$ , where  $G_{ij}$  is the trait mean in the  $ij$ th genotype class).

<sup>b</sup>Calculated as the difference between the phenotypic mean observed within the heterozygous class and the average phenotypic mean across both homozygous classes [ $d = G_{Bb} - 0.5(G_{BB} + G_{bb})$ ], where  $G_{ij}$  is the trait mean in the  $ij$ th genotype class.

Previous work utilizing a functional mapping analysis in forest trees have used a limited number of molecular markers (Li *et al.*, 2014). Li *et al.* (2014) applied this analysis in a bi-parental Scots pine cross using 319 markers. Hence, our work represents an advance in that we have been able to apply this approach at the genome-wide scale (178 101 SNPs) on maternal trees, with a dynamic trait dataset

comprising 14 time points/annual growth rings (i.e. cambial age). Latent traits represent significant time points in the trait development allowing us to detect putative genes at these critical junctures in wood formation. Functional mapping has also been applied in ecological studies (Paine *et al.*, 2012) and crops more recently. The fitting of growth models to the data describing growth trajectories of wood

formation phenotypes allowed the identification of marker-trait associations. This enabled us to track phenotype development against the genetic contributions at key time points.

Wood properties have previously been indicated to have a complex genetic architecture, in which association studies that make use of historical recombination represent a method that presents a substantial increase in QTL detection power for such complex traits (Hall *et al.*, 2016). In our study, the number of QTLs detected reflected the complex nature of the traits under study, and our experimental design allowed the detection of the largest/most significant QTLs. A previous functional mapping study involving SNPs in conifers applied two levels of evaluating QTLs (Li *et al.*, 2014), for which they have suggestive and significant QTLs, with our study only reporting the significant QTLs (single level), hence the small number of QTLs in our study. The small number of significant QTLs might also be due to the complex nature of the approximately 20 Gbp spruce genome. The sequence capture method only covered a total of 2331.1 kbp of exonic sequence, 2470.9 kbp of intronic sequence, 40.7 kbp of UTR-like sequence and 9119 exon–intron boundaries (Vidalis *et al.*, 2018). Therefore, a large portion of the genome was not represented and this would be compounded by the rapid LD in spruce, which might affect the number of significant QTLs detected. However, the numbers of QTLs detected in our study are in line with some previous studies in conifers (González-Martínez *et al.*, 2007), and with the drought association study in Norway spruce (29 significant SNP) (Trujillo-Moya *et al.*, 2018).

The QTL detected in our study explain a small proportion of the genetic variation and this could be due to several factors. This is in line with previous studies examining genetic variation in complex traits in coniferous species using forward genetic approaches, such as QTL (Sewell *et al.*, 2000; Novaes *et al.*, 2009) and AM (Wegrzyn *et al.*, 2010; Du *et al.*, 2013, 2018; Porth *et al.*, 2013; McKown *et al.*, 2014; Lamara *et al.*, 2016). The large effective population size in forest tree populations closely resembles humans, therefore making the ‘missing heritability’ issue found in human AM experiments relevant to forest tree populations. First, one of the hypothesis attributed to this ‘missing heritability’ is the substantial amount of quantitative variation linked to the cumulative effect of rare alleles that cannot be detected in GWAS using small sample sizes. Therefore in our study increasing the sample size from 517 individuals might allow the inclusion of rare alleles, explaining some of the missing heritability (Hamblin *et al.*, 2011; De La Torre *et al.*, 2019). The detection of true low-frequency alleles associated with complex traits is challenging as it requires large and genetically diverse populations (Hall *et al.*, 2016). Variants with low minor allele frequencies are usually discarded due to the potential of genotyping errors. However, rare alleles play an important


role in both the genetic regulation of traits and explaining the ‘missing heritability’ in forest species (De La Torre *et al.*, 2019). Therefore, this could have also contributed to the small effect sizes detected in our study as we filtered SNPs with low minor allele frequencies (<0.05 MAF). Second, allelic heterogeneity in which multiple functional alleles exist and are associated with different phenotypes, especially for such complex traits as those linked with wood formation. The presence of allelic heterogeneity will require a large population size that will encompass the allelic variations to account for the missing heritability (Bergelson and Roux, 2010). Third, non-additive effects mainly epistatically derived variation between genes might go undiscovered (Storey *et al.*, 2005). Most GWAS models have been designed to only consider the additive effects of markers. Numerous studies have shown that non-additive effects constitute a large part of the genetic variation of complex traits, these studies considered intra-locus (dominance) and inter-locus (epistatic) effects (Huang *et al.*, 2012; Zhou *et al.*, 2012; Mackay, 2013; Yang *et al.*, 2014; Du *et al.*, 2015). Yang *et al.* (2014) showed in corn an increase in the proportion of heritability explained when a model considering dominance was utilized and therefore allowing a better overview of heterosis. In rice Zhou *et al.* (2012) demonstrated the accumulation of multiple effects, including dominance and overdominance, which might partially explain some of the genetic basis for heterosis. Du *et al.* (2015) identified additive, dominant and epistatic effects explaining nearly two-fold high heritability in *Populus tomentosa* for 10 growth and wood property traits utilizing pathway-based multiple gene associations.

Lastly, epigenetic variation is also likely to be one of the sources of the ‘missing heritability’. With the development of advanced sequencing platforms, sophisticated genotyping tools have been developed to unravel epigenetic variation (Johannes *et al.*, 2009). Therefore, the influence of each of these factors on heritability strongly depends on the population sampled and inclusion of sophisticated genotyping tools in the case of epigenetics. The incorporation of a combination of advanced statistical models such as regional heritability mapping (RHM) and the detection of structural variants, insertion/deletions (InDels) and copy number variants in GWAS studies from several tree species has resulted in higher heritabilities being detected (Resende *et al.*, 2017a; Gong *et al.*, 2018)

### Trait trajectories and functional mapping

EBVs were plotted as phenotype data versus 14 consecutive cambial ages (Figure 1). All phenotypes under investigation are represented with thin light blue (Black average) curves, to visualize the nature of variation and growth trajectories of the phenotype (Figure 1 and Supporting Information Figure S1). The dissection of dynamic traits in forest trees has been predominantly performed using





**Figure 1.** EBV trajectories of four wood quality traits over time: (a) wood density, (b) late wood density, (c) annual ring width and (d) late wood ring width. Individual trajectories for each trait are shown in light blue lines and the black line represents the mean trajectory for the phenotype. These trajectories were used to determine the four latent traits of each tree, using linear splines with two knots.

single data points representing the value of the trait at a given developmental stage. The major disadvantage of such an approach is that it overlooks many of the factors that define the process of formation and development for important traits such as density and ring width. We utilized splines that have the advantage of not making *a priori* assumption about the shape of the curve and allow for the trait growth trends to be unbiased. Splines also allow for the characterization of the dynamic traits in terms of a few parameters derived from the spline models (Al-Tamimi *et al.*, 2016). The fitting of growth trajectories is considered as optimal because it treats phenotypes measured over time as different traits and also takes into account the correlation generated by the ordered time points (Yang *et al.*, 2006). The growth trajectories of the traits over time were calculated from the fitted splines and time intervals were identified and selected based on the characteristic growth trajectory of each trait, resulting in associations *across* and *within* traits being identified (Table 1). Therefore, indicating the control by different sets of genes at different time points for our longitudinal traits (Table 2), just as in some age-specific QTLs found in other conifers and rice (Verhaegen *et al.*, 1997; Emebiri *et al.*, 1998; Wu *et al.*, 1999). This approach has the potential to be applied to genomic prediction and selection studies for predicting individuals that would have the highest impact through the formation and development of a trait of interest. With application of differentially penalized regression (DiPR), pooled significant association markers can be utilized in GS in order to increase prediction accuracies (Bentley *et al.*, 2014).

### Linkage disequilibrium

The zygotic LD (squared correlation coefficient  $r^2$ ) was determined through the pooling of all  $r^2$  values and plotting them against the physical distances between the same SNP pair (Figure 2a). This allowed us to estimate the genome-wide degree of LD in Norway spruce, with the average LD for linked SNPs being inferred from the trendline (curve) of the nonlinear regressions. The fitted curve indicates the LD is low in Norway spruce, rapidly decaying by over 50% (from 0.50 to 0.20) (Figure 2a). The average distance associated with the LD decline for  $r^2 = 0.1$  varied from 14 to 1500 bp (Figures 2c,d and S2). Neale and Savolainen (2004) reported an LD decayed to less than 0.20 within roughly 1500 bp based on 19 candidate genes in loblolly pine. As conifers are highly outcrossing a rapid LD decay is expected, however in spruces the LD displays diverse patterns among different genes or the same genes in different species. The LD decline in spruces was also noted to be roughly between a few base pairs and 2000 bp (Namroud *et al.*, 2010). These diverse heterogenous LD patterns were also observed when we analyzed the LD for individual contigs that had significant associations to our traits (Figure 2c,d, Figure S3). The general LD estimate of

all the SNP pairs indicated a fast LD decay (Figure 2a). This rapid decay could be due to the number of contigs analyzed in relation to the large Norway spruce genome, as well as the use of zygotic LD between genotypes. Lu *et al.* (2016), noted that the calculation of gametic LD from phased haplotypes indicated a slower LD decay than when using zygotic LD in loblolly pine. However, they also observed varying rates of LD decay between genes and across different genome regions (Lu *et al.*, 2016). Therefore, the generality of the LD patterns within the Norway spruce genome remains to be further analyzed because only a relatively small and highly specific portions of the genome was studied here.

### Population inference


To account for effects derived from population stratification we performed a principal component analysis (PCA). The top two explained a total of 5.3% of the variation. Population structure inference of clusters detected by PCA was performed by ADMIXTURE (Figure S3) and the best K value plotted from the cross-validation error term. Using the best K method, K = 2 better explained the genetic structure of the study population (Figure S3).

### Overall summary of genetic associations

Several associations were shared *within* each trait and *across* traits in the analysis. WD, Ring width (RW), Transitional ring width (TRW) and Latewood number of cells (LNC) had one (MA\_33109\_11804), two (MA\_10434805\_21408 and MA\_20322\_28351), one (MA\_817099\_1105) and one (MA\_10428744\_29330) QTLs shared by two or more latent traits, respectively. Common QTLs *within* RW were observed for slope,  $\beta_2$  and  $\beta_3$  latent traits, with moderate frequencies ranging from 0.521 to 0.615 and influenced their respective traits to modest degrees (PVE in ranges of 0.18–2.66%).

For QTLs common *across* the different latent traits, SNP MA\_10434805\_21408 was shared between latewood wood density (LWD), RW and Modulus of elasticity (MOE); this is not surprising because of the close correlation between MOE and wood density. Intron variant MA\_10434805\_21408 explained between 0.18 and 2.66% of the PVE observed in the respective traits. This SNP associated also had high frequencies of 0.602 and 0.615 in MOE and RW explaining PVE of 1.00 and 2.66%, respectively (Table 2). SNP MA\_10435406\_13733 was shared between WD, Earlywood wood density (EWD) and Earlywood number of cells (ENC), was associated with the intercept trait for WD and EWD and the slope latent trait in ENC (Table 1), with PVE ranging from 0.01 to 4.64%. The QTL had a high influence on the density related traits as it explained 4.64% (WD) and 3.38% (EWD) both exhibiting a partial dominant inheritance pattern (Table 2).

Numbers of cells (NC), ENC, TNC and LNC traits were associated with a total of three putative genes and three





**Figure 2.** (a) Decay of linkage disequilibrium (LD) across all the tagged genomic sequences, the majority being exonic regions. The squared correlation coefficient between loci ( $r^2$ ) is plotted against distance, in base pairs, separating loci. The fitted curve (red) is representative of the trend of decay from the 178 101 SNPs utilized in the association mapping (AM). (b) Decay of LD with distance in base pairs between sites from across 41 contigs with significant associations. (c) Decay of LD across contig MA\_96191 that has a significant association for ratio of percentage earlywood vs latewood on which two probes were captured. (d) Decay of LD on contig MA\_80033 indicating the variable LD in the genome.

protein domains. Of the three putative genes, two are associated with serine/kinase activity and one is involved in cysteine and methionine synthesis (Table S1). All the SNPs associated with these traits were either downstream or upstream of coding regions and may therefore act as modifiers of gene expression.

Wood percentage traits, early wood proportion (EP), LP, TP and the ratio of EP/LP had significant associations with 10 SNPs. Four of the six significant SNP variants for EP/LP are modifiers with the other two SNPs, being a synonymous (MA\_96191\_7122) and missense (MA\_1045136\_4310) variant. The synonymous SNP MA\_96191\_7122 was consistent with the codominant mode of inheritance (Table 2). The significant SNP MA\_15729\_40331, an intron variant, that is associated with EP, is located in the gene MA\_15729g0010, homologous to a DNA-3-methyladenine glycosylase II enzyme (Table 1).

WD, EWD, TWD and LWD had a total of 12 significant associations. A missense SNP, MA\_33109\_11804, was associated with WD and located within the gene homologous to an Arabidopsis senescence-associated gene 24 (Table S1). Of the three significant SNP associations for Transitional wood density (TWD), two, SNP MA\_10235390\_3386 (stop gained) and SNP MA\_212523\_89044 (upstream gene variant) were identified within genes. Two of the three significant SNPs identified for LWD were intron variants (MA\_10434805\_21408 and MA\_10436058\_4902) with the third being a missense variant (MA\_62987\_13474).

Trees showing a positive correlation between growth and density had seven QTL specific for this observed phenomenon (MI), explaining a PVE ranging between 0.05 and 1.82% (Table 1). The seven associated SNPs, were two upstream gene variants, two missense variants one intergenic variant, one stop gained variant and one synonymous nucleotide replacement (Table 1). The SNP MA\_1378g0010\_4718 encodes for a premature stop codon on gene MA\_1378g0010. Two SNPs associated with the slope latent trait for MI (MA\_1378\_4718 and MA\_10427214\_13968) have an overdominance inheritance pattern with the C and G alleles being dominant, respectively (Table 2; Figure 3).


#### Genetics associations for genes of known function in wood formation

**Intercept associations.** Our study identified several interesting genes linked to the significant QTLs from the intercept latent trait, which represents the mean from our spline model. This resulted in 17 significant associations with a

PVE ranging from 0.50 to 4.64% associated with the intercept latent trait. The modes of action determined by the non-additive effects of these significant SNP associations to the intercept latent trait were one for overdominance ( $|d/a| > 1.25$ ), codominant ( $|d/a| < 0.50$ ) 12 and four SNPs were partial to fully dominant ( $0.50 < |d/a| < 1.25$ ).

Ring width phenotypes RW, TRW, and LRW were linked with a total of three gene models associated with the intercept latent trait (Table S1). Of these putative genes associated with RW phenotypes, gene MA\_10694g0010 was of particular interest with regards to wood formation. SNP MA\_10694\_9101 with a partial to fully dominant mode of inheritance (Table 2) was located on the gene MA\_10694g0010 that is homologous to an enzyme involved in cell wall biosynthesis, endoglucanase 11-like, and was associated with RW (Table S1) and was expressed in the wood (phloem+cambium+xylem) component of spruce (Figure 4). This enzyme is a vital component of xylogenesis and is involved in the active digestion of the primary cell wall (Goulao *et al.*, 2011). Endoglucanases have been proposed as enzymes involved in controlling cell wall loosening (Cosgrove, 2005). Endoglucanase 11-like gene is part of the endo-1 family in which the eno-1-4- $\beta$ -glucanase *Korrigan* gene belongs. Characterization of the *Korrigan* gene in *P. glauca* has identified it to be functionally conserved and essential for cellulose synthesis (Maloney *et al.*, 2012).

Density-related phenotypes (WD, EWD, TWD and LWD) had two significant associations detected for the intercept. Both associations were detected by SNP MA\_10435406\_13733 and were both partial to fully dominant in their form of inheritance (Table 2). The SNP MA\_10435406\_13733 downstream on gene MA\_10435406g0010 was also significantly associated with the trait ENC slope latent trait. The association of this gene with the WD and EWD intercept implies that it has an impact on the overall development of density throughout the growth period. This result coincides with previous report about the influence of the earlywood component on the properties of the annual ring as a whole in Scots pine (Li *et al.*, 2014). Association of the same gene also with the slope latent trait of ENC corroborates the predictive value of number of cells for wood density. The gene is homologous to phosphoadenosine phosphosulfate reductase (PAPS), which plays a central role in the reduction of sulfur in plants. An analysis of PAPS enzymes in Arabidopsis (Klein and Papenbrock, 2004) and *Populus* (Kopriva *et al.*, 2004) revealed that enzymes involved in sulfate conjugation play an important role in plant growth and development (Klein and Papenbrock, 2004).



**Figure 3.** Box plot of the estimated genotypic effect on the phenotypes in the study. The significant SNPs associated and each one of the traits have been correlated to give the impact each genotype has on the average of the overall trait

**Slope associations.** The analysis of slope (rate of change) of wood formation traits over cambial ages, to our knowledge, has never been dissected in any Norway spruce QTL or AM analyses.

For slope latent traits two significant candidate genes concerning wood formation, PAPS and Proliferating Cell Nuclear Antigen (PCNA), were detected *across* related traits density, growth number of cells and MOE. The codominant SNP MA\_10435406\_13733 (Table 2) that is a 3'-gene variant for WD was located on a gene that is homologous to a phosphoadenosine phosphosulfate reductase gene *cysH\_2* and common *across* ENC, WD and EWD.

The SNP MA\_10434805\_21408 was located on the gene MA\_10434805g0010, which is homologous to an Arabidopsis PCNA protein (Table S1) and was ubiquitously expressed with high levels in the wood (phloem+cambium+xylem) component of spruce (Figure 4). This SNP is associated *across* LWD, RW and MOE with partial to fully dominance ( $0.50 < |d/a| < 1.25$ ) for all three associations (Table 2). The presence of these common QTL suggests that these traits might be under the control of the same genes or genetic pathways. Chen *et al.* (2014) reported a significant positive genetic correlation between wood density and MOE, which increased with tree age. However, wood volume growth and density have a negative correlation (Chen *et al.*, 2014), our study was able to detect QTLs for trees exhibiting a positive correlation for this phenomenon (MI). The common QTL observed *across* WD, EWD and ENC indicates that the number of cells during the

juvenile wood development stages has a significant impact on the overall density. The seasonal changes in EWD to LWD have been speculated to be due to a change in auxin levels leading to the initiation of wall-thickening phase, which has a direct impact on the wood quality traits such as MOE. This phase coincides with the cessation of height growth and where available resources are used for cell wall thickening (Sewell *et al.*, 2000), which may explain the common QTL between LWD, RW and MOE, as part of the same feedback loop mechanism.

$\beta_2$  and  $\beta_3$ . When analyzing QTLs associated with the two latent traits  $\beta_2$  and  $\beta_3$ , 16 significant associations were detected, with phenotypic variances ranging from 0.01 to 4.51% (Table 1). Five of the significant markers were consistent with overdominance ( $|d/a| > 1.25$ ), with the 11 markers being dominant ( $0.50 < |d/a| < 1.25$ ) (Table 2, Figure 3).

Wood density phenotypes (WD, EWD, TWD and LWD) had three significant associations with the  $\beta_2$  and  $\beta_3$  latent traits (Table 1). The upstream variant MA\_33109\_11804 associated with both the slope and  $\beta_2$  latent traits of WD was detected on the gene model MA\_33109g0100 that is homologous to the *Arabidopsis* senescence-associated gene 24 (Table S1) and the gene being expressed in shoots and buds of spruce (Figure 4). An association for the latent trait  $\beta_2$  of TWD with a codominant SNP (MA\_212523\_6278) (Table 2) was located upstream of gene MA\_212523g0010 homologous to Kinesin-related protein 13 (geneL484\_021891) and ubiquitously expressed in shoot, buds and wood component of spruce, indicating its important

Phenotype	Trait	gene_id	confidence	Ath_Nmae/description	Immature male cone	Immature female cone	Needles (2009)	Needles (2008)	Infected needles	Vegetative shoots, May 27, 2010	Vegetative shoots, June 21, 2010	Pineapple galls	Buds early season developing	Needles from vegetative shoots Aug.	Stem from vegetative shoots Aug.	Needles from vegetative shoots Sept.	Buds late season developed	Needles from dried twig	Needles from girdled twig	Stem from girdled twig	Early morning needles 5.30	Mid-day needles 12.00	Late afternoon needles 19.30	Night needles 23.30	Wood (phloem+cambium+xylem early)	Wood (phloem+cambium+xylem late)	
Width	RW	MA_879270g0010	High	Alpha-DOX1	-2	1.3	1.1	0	-1	4.6	5.3	4.9	-2.1	1.5	-2	1.5	-1	1	0.2	-2	-1	-0	0.3	-1	-2	-2	
		MA_10694g0010	High	GH9B1	3	0	0	0	0	0	0	0	0	0	0	0	0.9	0	0	0	0	0	0	0	0	1	0
		MA_10434805g0010	Medium	PCNA1	1.6	-0	-1	-1	-1	-1	-1	0.4	0.9	0.4	0.2	0.7	-0	0	0.9	0	-0	-0	1.2	-1	1	1	0
		MA_20322g0010	Medium	AP2/B3-like	2.8	-1	-2	0	0.1	-1	0.8	1.7	2.8	0.4	-2	0.5	0.2	-2	-0	-2	-2	-1	-0	-2	3	2	2
		MA_879384g0010	Medium		0.2	-2	-1	-0	0.3	-2	0.4	1.8	2	0	-2	1.3	-0	-1	-1	-1	-1	-2	1.4	0	0	0	1
	TRW	MA_214776g0010	Medium	Kinase	-3	1	-0	0	1.3	0	0.4	-1	-0.8	-2	0.7	-2	2.1	2	-2	0.9	1.1	1.6	-0	1	-4	-5	
		MA_38472g0010	High	AHDP	-3	1	0.7	1	0.7	0.9	0.5	-1	-0.4	-1	0.2	-1	-1	-1	-0	-1	0.6	-0	0.2	0.4	0	-4	-2
		MA_817099g0010	Medium	Senescence protein	0.8	-1	0.1	-1	-1	-0	-0	0.3	0.6	1.8	-0	-1	-1	0	0.7	-1	0.5	0.1	-2	-0	2	1	
		MA_10434624g0010	Medium	PME2	3.9	-1	-1	2	1	-1	-3	-3	-2.8	-1	0.2	-3	2.3	1	0.7	-1	-0	0.8	-1	1	4	3	
		MA_10434624g0010	Medium	PME2	3.9	-1	-1	2	1	-1	-3	-3	-2.8	-1	0.2	-3	2.3	1	0.7	-1	-0	0.8	-1	1	4	3	
Number of Cells	NC	MA_528074g0010	High	EFR	-2	0.5	0.5	0	0.3	0.7	0	-2	-4	-2	0.8	-3	-2	0	-2	0.2	1.2	0.6	-2	0	-2	-1	
	ENC	MA_402880g0010	High	MEE55	1	-1	-0	-1	1.4	-1	2.3	1.4	0.8	2.1	0	2.7	-1	-1	1.6	0	-1	-1	0	-1	-1	0	
	TNC	MA_9447489g0010	Medium	Metallocarboxpeptidase	-0	0.5	-0	0	0.3	-0	-0	-0	-0.2	-1	0.1	-1	0.2	0	-1	0.7	0.7	-0	-1	0	0	0	
	LNC	MA_10428744g0010	High	CLK4	-0	0.8	0.1	-0	0.4	-0	0.2	-0	-0.2	-0	-0	0	0	0	0	0.6	0.4	0.6	-1	0	0	0	
Wood percentage	EP	MA_246125g0010	Medium	TIR-NBS-LRR class	0.1	2.1	0.7	0	0.2	0.4	-0	-2	-1.7	-1	-2	-2	0.5	1	0.4	-0	0.2	0.6	-2	-1	-2	-0	
	LP	MA_10434007g0010	High	URM9	0.4	-1	-1	-0	-1	-1	-0	0.1	0.4	-0	-0	0.3	0.5	-0	-0.2	-0	0.1	0.3	0	0	0	1	
	TP	MA_462319g0010	High	Myosin-binding protein	-0	0.8	0.5	0	0.4	0.6	0.6	-0	-0.2	-0	0.1	-1	0.6	0	-0	0.5	0.1	-0	-0	-0	-0	-0	
	EP/LP	MA_80954g0010	Medium	PK54	-2	-0	-2	-2	3.6	-2	4.3	2.9	-0.4	2.5	1.7	1.3	-2	-0	-0.1	-2	-1	0.3	1.4	0	0	0	-2
		MA_98424g0010	High	STN1	-0	-1	-0	0.2	0.2	0.1	0	0.3	0.8	-0	-0	0.3	-0	-0	-1	-0.5	0	-0	-0	-0	-1	2	
		MA_8790100g0010	Medium	AAA-type protein	-0	0.2	-0	0	-0	-0	-0	-1.1	0.3	0.5	-0	0.9	0	0.8	0.5	0	-0	-1	0	0	-1	0	
		MA_96191g0010	Medium	UGT73B2	-2	1.5	0.5	-1	5.3	1.1	4.9	-2	1.3	-2	1.5	-1	0.7	0	-2	-1	-0	0.3	4.6	-1	-2	-2	
	MA_460877g0010	Medium	TIR-NBS-LRR class	-0	-2	0	-0	1.1	-1	0.3	-1	0.7	0.7	-0	-1	0.5	0.5	-0	-0	-0	-0	-0	-1	1	0	0	
Density	Ring	MA_33109g0010	Medium	SAG24	0.5	-1	-1	-0	0.1	-0	-0	0.7	0	0	0.5	0.8	-0	0.1	-0	0.1	-0	0.6	-0	0	0	-0	
	EWD	MA_10432646g0010	High	coil domain	0.2	-0	-0	0.2	-0	-0	-0.3	-0	-0.3	-0	0.3	-0	0.1	0.3	0.4	0	0.1	-0	0	0	0	-0	
		MA_20321g0010	High	BSL2	0	0	-0	-0	0.4	-0	-0	-0.2	-0	0.5	-0	0.5	0	0	0	0	0.6	0.2	-1	1	0	0	
	TWD	MA_10235390g0010	Medium	MSH5	1.3	0	0	0	0	0	0	0	0	0.9	0	0	0	0	0	0	0	0	0	0	0	0	
		MA_10433411g0010	Medium	RACK1A	0.1	-1	-0	-1	0	-0	-1	0.3	0.5	-0	-0	0.3	-0	1	0.2	-1	0.2	-0	0	1	-0	0	
		MA_212523g0010	High		3.1	-0	-1	-1	-1	0.1	1.7	2	1.4	-1	2.4	-2	-1	1.9	-1	0	-2	3.1	-0	3	2	2	
LWD	MA_10436058g0010	Medium	Spc97/Spc98 family	0	0	0	0	0	1.2	0	0	1.2	0	0	0	0	0	0	1.8	1	1	0	0	0	0		
Mass index	MI	MA_10435002g0010	Medium	OVA2	-1	0.3	0.5	1	-1	-0	-1	-1	-1	-1	0.8	-1	0	1	-1	0.6	1.2	1.4	0.2	1	-2	-2	
		MA_99004g0010	Medium	SHR	-1	0	0.5	0	1	0.1	1.3	-0	1.9	0.7	-0	0.1	-0	-1	1.2	-1	-0	-1	2.1	-1	-5	-3	
		MA_109804g0010	High	YGL	-2	0.4	-0	0	0	0	0.5	-3	-3.8	-1	1	-4	2.5	0	-0	0.8	0.9	0.5	-2	1	-5	-5	
		MA_10427214g0010	Medium		2.9	0	0	0	0	0	0.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

**Figure 4.** The heatmap showing the expression levels (VST values) of spruce candidate genes in different organs and tissues based on data of Nysted *et al.* (2013) available at <http://congenie.org>.

function in this species (Figure 4). This association is of interest because Kinesin-related proteins are known to be involved in secondary wall deposition, which can impact wood density (Zhong *et al.*, 2002), cell wall strength and oriented deposition of cellulose microfibrils. Therefore, these proteins would have a direct correlation with the increase in density at the latent trait  $\beta_2$  at age 6 years (Figure S1).

Ring width phenotypes (RW, TRW and LRW) had eight significant associations identified for the latent traits  $\beta_2$  and  $\beta_3$ , explaining PVE ranging from 0.01 to 4.51% (Table 1). The synonymous SNP MA\_2032\_28351 associated with the  $\beta_2$  latent trait for RW is located on a gene homologous with a plant-specific B3-DNA binding protein domain explaining 1.78% variation and is shared among various plant-specific transcription factors. This includes transcription factors involved in auxin and abscisic acid responsive transcription (Yamasaki *et al.*, 2004). Auxin is one of the central phytohormones in the control of plant growth and development (Abel and Theologis, 1996), and also known to be involved in cell wall loosening and

elongation (Cosgrove, 2016). This association was detected *within* the RW phenotype and detected for both  $\beta_2$  and  $\beta_3$  latent traits (Table 1). Therefore, this domain could be involved with transcription factors involved in both the decrease and increase of RW (Figure S1). Three putative genes were associated with the  $\beta_2$  and  $\beta_3$  latent traits for the TRW phenotype. Of interest a senescence-associated protein associated on the TRW  $\beta_2$  latent trait with the missense variant MA\_817099\_1105. This might be linked to the decrease in TRW at year 6 (Figure S1) due to the decline of photosynthetic rate known to be induced by the activity of senescence related proteins (Sillanpää *et al.*, 2005). The gene was highly expressed in both the early and late wood components of spruce supporting the row of these senescence genes in controlling tree growth (Figure 4). This association was also identified for the slope latent trait indicating a potential impact on the rate of change of transitional wood. The detection of senescence related genes for wood density related phenotypes for both the slope and  $\beta_2$  latent trait (MA\_33109\_11804) could indicate a possible relationship between the genes



influencing RW and wood density. Contig MA\_10434624 is homologous to a pectin esterase and was associated with the downstream variant MA\_10434624\_20686 with an over-dominant mode of inheritance for LRW ( $\beta_3$  latent trait) at year 10 (Table 2, Figure S1). This significant downstream SNP (MA\_10434624\_20686) associated with LRW on gene MA\_10434624g0020 and homologous to pectin methyl esterases (PMEs), which are cell wall associated enzymes responsible for demethylation of polygalacturonans (Phan *et al.*, 2007). This gene was observed highly expressed in developing wood (Figure 4), indicating its importance for growth in spruce. This enzyme has been shown to be linked with many developmental processes in plants, such as, cellular adhesion and stem elongation (Micheli, 2001). An association study in *Picea glauca* (Moench) Voss identified a significant nonsynonymous SNP coding for cysteine associated with earlywood and total wood cell wall thickness associated with pectin methyl esterase (Beaulieu *et al.*, 2011). Our study identified a PME SNP association in the late wood stage, supporting the importance of PMEs in wood cell development.

When analyzing QTLs detected for traits linked to the percentage of cells (EP, LP and EP/LP) we identified three putative candidate genes, DNA-3-methyladenine glycosylase II enzyme, phytochrome kinase substrate 1 and glycosyltransferase. Synonymous SNP (MA\_96191\_59480) within the gene MA\_96191g0010, which is homologous to Glucosyltransferase in *Picea sitchensis* was associated with the EP/LP,  $\beta_2$  latent trait. The gene is highly expressed in vegetative shoots (June) and during the late afternoon in needles (Figure 4). Glycosyl transferases operate by facilitating the catalytic sequential transfer of sugars from activated donors to acceptor molecules that form region and stereospecific glycosidic linkages (Lairson *et al.*, 2008). The Arabidopsis ortholog (UDP-glucosyltransferase 73B2) encodes for a putative flavonol 7-O-glucosyltransferase involved in stress responses. In our study, this significant association was associated with EP/LP, however a nonsynonymous variant in a gene coding for a glycosyl transferase in *Populus* was associated with fibre development and elongation (Porth *et al.*, 2013).

Several receptor-like kinases (TIR/NBS/LRR and serine/threonine-protein phosphatase) homologues were identified *across* traits (TRW, NC, EP, EP/LP and EWD) (Table S1). Approximately 2.5% of the annotated genes in Arabidopsis genome are RLK homologues (Shiu and Bleecker, 2001), in which they, among other functions, play an important role in the differentiation and separation of xylem and phloem cells (Fisher and Turner, 2007). Similar to our study a synonymous SNP in an RLK gene was associated with EP in white spruce (Beaulieu *et al.*, 2011), hence RLKs seem to be involved in modifying a number of different wood properties from density to cell identity and number.

Norway spruce trees that possess the ability of fast growth and high wood density are very rare, but such trees and associated SNPs were discovered in our study. Trees combining these traits are of high interest to the forestry industry. Of the seven genes significantly linked to this phenomenon of particular interest was a synonymous SNP on MA\_99004g0100 gene homologous to a transcription factor from the GRAS family (Table S1). GRAS is an important class of plant-specific proteins derived from three members: GIBBERELLIC ACID INSENSITIVE (GAI), REPRESSOR OF GAI (RGA) and SCARECROW (SCR) (GRAS) (Hirsch and Oldroyd, 2009). GRAS genes are known to be involved in the regulation of plant development through the regulation of gibberellic acid (GA) and light signalling (Hirsch and Oldroyd, 2009; Cenci and Rouard, 2017). Furthermore GA signalling has also been shown to stimulate wood formation in *Populus* (Mauriat and Moritz, 2009). Therefore, the GRAS transcription factor identified here and the other six genes positively associated with MI provide interesting genetic markers and tools to understand this phenomenon.

Wood density traits were associated with a total of 12 genes, the largest number of genes identified from the contigs. The percentage of wood was linked to 10 putative genes, cell width had nine putative genes and number of cells was associated with six genes. Two genes were shared *across* multiple traits, PCNA was common *across* RW and LWD, and phosphoadenosine phosphosulfate reductase was shared *across* WD, EWD and ENC.

## EXPERIMENTAL PROCEDURES

### Plant material and phenotype data

Plant material and phenotype data used in this study have previously been described in Chen *et al.* (2014). In brief, two progeny trials were established in 1990 in southern Sweden (S21F9021146 aka F1146 (trial1) and S21F9021147 aka F1147 (trial2)). We selected 517 families originating from 112 sampling stands to use in the investigation of wood properties. At each site, increment wood cores of 12 mm were collected at breast height from six trees of the selected families (1.3 m) (6 progeny  $\times$  2 sites = 12 progenies in total). In total, 5618 trees, 2973 and 2645 trees from the F1146 and F1147 trials respectively, were analysed. The pith to bark profiling of growth and wood physical attributes was performed using the SilviScan instrument (Evans and Ilic, 2001) at Innventia, now part of RISE, Stockholm, Sweden, where also the initial data evaluations were performed (Methods S2). These included the identification and dating of all annual rings and their compartments of early wood (EW), transition wood (TW) and late wood (LW). For this, a density-based '20-80' definition was used, described and discussed in (Lundqvist *et al.*, 2018). Traits of interest to breeders were derived from the SilviScan data, such as the radial NC and Mass Index (MI) introduced to express the relative amount of biomass at breast height.

The investigation was triggered by the observation that some trees broke the unfavourable negative correlation of the trait MI which is between density and growth. They produced, more biomass than expected, and it was therefore important. In order to

identify putative genes behind high values for this trait. MI was defined as:

$$\text{Mass index} = \frac{(\text{Individual cross} - \text{sectional average density})}{(\text{population cross} - \text{sectional average density})} \times \frac{(\text{individual cross} - \text{sectional area})}{(\text{population average cross} - \text{sectional area})}$$

The traits investigated in this study are listed in Table 3.

### Statistical analysis

EBVs were calculated for growth and wood quality traits for 14 consecutive annual growth rings. The variance and covariance components were estimated using ASREML 4.0 (Gilmour *et al.*, 2014) as described in Chen *et al.* (2014). In brief, the EBVs at each cambial age were estimated using univariate, bivariate or multivariate mixed linear models. The following univariate linear mixed model for joint-site analysis was fitted to calculate EBV:

$$Y_{ijkl} = \mu + S_i + B_{j(i)} + F_k + SF_{ik} + e_{ijkl} \quad (1)$$

where  $Y_{ijkl}$  is the observation on the  $l$ th tree from the  $k$ th family in  $j$ th block within the  $i$ th site,  $\mu$  is the general mean,  $S_i$  and  $B_{j(i)}$  are the fixed effects of the  $i$ th site and the  $j$ th block within the  $i$ th site, respectively,  $F_k$  and  $SF_{ik}$  are the random effects of the  $k$ th family and the random interactive effect of the  $i$ th site and  $k$ th family, respectively,  $e_{ijkl}$  is the random residual effect. The random family and site by family interaction effects are assumed to follow  $N(0, \sigma_f^2)$   $N(0, \sigma_{sf}^2)$  and, respectively, where  $\sigma_f^2$  and  $\sigma_{sf}^2$  are the estimated family genetic variance and site by family interaction variance, respectively. Residual variation  $e$  was assumed to  $N\left(0, \begin{bmatrix} I_{n1}\sigma_{e1}^2 & 0 \\ 0 & I_{n2}\sigma_{e2}^2 \end{bmatrix}\right)$ , where  $\sigma_{e1}^2$  and  $\sigma_{e2}^2$  are the residual variances for site 1 and site 2,  $I_{n1}$  and  $I_{n2}$  are identity matrices,  $n1$  and  $n2$  are the number of individuals in each site. The fit of different models was evaluated using the Akaike Information Criteria (AIC) and the optimal model was selected based on a compromise of model fit and complexity.

**Table 3** List of the phenotypes, their abbreviations and measurement unit

Phenotype	Abbreviation	Unit
Ring wood density	WD	kg m <sup>-3</sup>
Early wood density	EWD	kg m <sup>-3</sup>
Transition wood density	TWD	kg m <sup>-3</sup>
Late wood density	LWD	kg m <sup>-3</sup>
Ring width	RW	μm
Early wood ring width	ERW	μm
Transition wood ring width	TRW	μm
Late wood ring width	LRW	μm
Ring number of cells	NC	
Early wood number of cells	ENC	
Transition wood number of cells	TNC	
Late wood number of cells	LNC	
Early wood percentage	EP	%
Transition wood percentage	TP	%
Late wood percentage	LP	%
Early/late wood percentage	EP/LP	%
Modulus of elasticity	MOE	GPa
Mass index (density × growth)	MI	

### Latent traits

The EBVs were plotted against cambial age (annual ring number) to produce time trajectories for each trait (Figures 1 and S1). Spline model was fitted to the trajectories and their curve parameters describing the character of their development over time were used as latent traits in order to describe the dynamics of the EBVs across age.

The general definition of a linear spline with multiple knots is as follows

$$y(t) = \beta_0 + \beta_1 t + \beta_2 (t - K_1)_+ + \beta_3 (t - K_2)_+ + \dots + \beta_{1+m} (t - K_m)_+, \quad (2)$$

which is continuous and where  $K_i$  ( $i = 1, \dots, m$ ;  $K_1 < K_2 < \dots < K_m$ ) are defined as knots, and  $(t - K_i)_+ = (t - K_i)$  if  $t > K_i$  ( $K_i > 0$ ;  $i = 1, \dots, m$ ), and otherwise is equal to zero. The number of knots has to be properly defined to provide an accurate description of the data under investigation, while avoiding overadaptation to data (Li and Sillanpää, 2015; Camargo *et al.*, 2018). In our case, we found two knots most suitable to the time intervals investigated. Hence, the linear spline model to describe the growth trajectory of individual  $i$  applied in this study was defined as:

$$y(t) = \beta_0 + \beta_1 t + \beta_2 (t - K_1)_+ + \beta_3 (t - K_2)_+ + \varepsilon_i(t), \quad (3)$$

$$\varepsilon_i(t) \stackrel{\text{iid}}{\sim} N(0, \sigma^2).$$

In equation (2), the intercept  $\beta_0$ , slope parameters  $\beta_1$ ,  $\beta_2$  (at Knot 1 ( $K_1$ )) and  $\beta_3$  (at Knot 2 ( $K_2$ )) are estimated by standard least squares (Ruppert *et al.*, 2003). The four estimates were used as the latent trait in the subsequent QTL analysis conducted in RSTUDIO (Team, 2015), and then analysed using the LASSO model to identify SNPs showing significant associations to the traits.

The intercept and slopes were used to evaluate the mean and rate of change for the trait across the annual rings, respectively.  $\beta_2$  and  $\beta_3$  represent inflection points in the cambial age trajectories where the development of the EBVs enters new phases. These two points ( $\beta_2$  and  $\beta_3$ ) are therefore supposed to have biological significance, warranting a closer analysis of the genes imparting these shifts in the EBVs dynamics. The four latent traits show lower correlations compared with the direct measurements on the original scales and they also have constant variances, thereby reducing the need to account for residual dependencies in the model (Wu *et al.*, 2004; Yang and Xu, 2007; Li *et al.*, 2014).

### Sequence capture, genotyping and SNP annotation

Total genomic DNA was extracted from 517 maternal trees, using the Qiagen Plant DNA extraction (Qiagen, Hilden, Germany) protocol with DNA quantification performed using the Qubit<sup>®</sup> ds DNA Broad Range (BR) Assay Kit (Oregon, USA). Extracted DNA was submitted to RAPiD Genomics (USA) where DNA library preparation and capture sequencing were performed. Sequence capture was performed using the 40 018 diploid probes designed and evaluated for *P. abies* (Vidalis *et al.*, 2018). The Illumina sequencing compatible libraries were amplified with 14 cycles of polymerase chain reaction (PCR) and the probes were then hybridized to a pool comprising 500 ng of 8 equimolar combined libraries following Agilent's SureSelect Target Enrichment System (Agilent Technologies, <https://www.agilent.com/>). These enriched libraries were then sequenced using an Illumina HiSeq 2500 instrument (San Diego, USA) on the 2 × 100 bp sequencing mode.



Raw reads were mapped against the *P. abies* reference genome v.1.0 using BWA-MEM (Li, 2013). SAMTOOLS v.1.2 (Li et al., 2009) and Picard (<http://broadinstitute.github.io/picard>) were used for sorting and marking of PCR duplicates. Variant calling was performed using GATK HAPLOTYPECALLER v.3.6 (Van der Auwera et al., 2013) in gVCF output format. Samples were then merged into batches of approximately 200 before all 517 samples were jointly called.

Variant Quality Score Recalibration (VQSR) method was performed to avoid the use of hard filtering for exome/sequence capture data. For the VQSR analysis two datasets were created, a training subset and an input file. The training dataset was derived from the Norway spruce genetic mapping population showing expected segregation patterns (Bernhardsson et al., 2019) and assigned a prior value of 15.0. The input file was derived from the raw sequence data using GATK with the following parameters: extended probe coordinates by +100 excluding INDELS, excluding LowQual sites, and keeping only bi-allelic sites. The following annotation parameters QualByDepth (QD), MappingQuality (MQ) and BaseQRankSum, with tranches 100, 99.9, 99.0 and 90.0 were then applied for the determination of the good versus bad variant annotation profiles. After obtaining the variant annotation profiles, the recalibration was then applied to filter the raw variants. Using VCFTOOLS v.0.1.13 (Danecek et al., 2011), SNP trimming and cleaning involved the removal of any SNP with a MAF and 'missingness' of <0.05 and >20%, respectively. The resultant SNPs were annotated using default parameters for SNIPEFF 4 (Cingolani et al., 2012). Ensembl general feature format (GFF; gene sets) information was utilized to build the *P. abies* SNIPEFF database.

### Genetic structure and mode of inheritance

Linkage disequilibrium was calculated as the squared correlation coefficient between genotypes ( $r^2$ ), globally with special attention given to all the contigs with significant associations in VCFTOOLS v.0.1.13 software using the 'geno-r2' routine (Danecek et al., 2011). The trendline of LD decay with physical distance was fitted using nonlinear regression (Hill and Weir, 1988) and the regression line was displayed using RSTUDIO (Team, 2015). Non-additive effects of the significant markers was determined using the ratio of dominance ( $d$ ) to additive ( $a$ ). The ranges were: partial or complete dominance ( $-0.50 < |d/a| < 1.25$ ) and additive ( $-0.50 \leq |d/a| \leq 0.50$ ), with  $|d/a| > 1.25$  being equal to over- or underdominance (Eckert et al., 2009).

FactoMiner (Multivariate Exploratory Data Analysis and Data Mining) (Husson et al., 2017) implemented in RSTUDIO software was used to perform PCA. The covariate matrix derived from the PCA was then displayed by plotting principal component 1 scores against principal component 2 scores. The components of the PCA covariate matrix were then applied to the AM to account for population structure and correcting for any stratification within the study. Significance of each genetic principal component (PC) was determined using the Tracy-Widom (TWi) distribution and a significance threshold of  $P = 0.01$ . For population clustering, ADMIXTURE v.1.3.0 (Alexander et al., 2009) was used with five-fold cross-validation and 200 bootstrap replicates. The bestK method was implemented in RSTUDIO to determine the best K with the use of an elbow plot on the cross-validation error.

### Trait association mapping

It is natural to use LASSO method for simultaneous estimation of SNP effects and selecting a sparse subset of trait-associated SNPs to the multilocus association model. This is because LASSO has

nice properties like being able to handle high-dimensional cases with  $p \gg n$  (i.e., a number of SNPs much larger than number of individuals) and selecting only a single representative SNP from the group of highly dependent SNPs. The LASSO model as described by Li et al. (2014), was applied to all latent traits for the detection of QTLs.

The LASSO model:

$$\min_{(\alpha_0, \alpha_j)} \frac{1}{2n} \sum_{i=1}^n \left( y_i - \alpha_0 - \sum_{j=1}^p x_{ij} \alpha_j \right)^2 + \lambda \sum_{j=1}^p |\alpha_j|, \quad (4)$$

where  $y_i$  is the phenotypic value of an individual  $i$  ( $i = 1, \dots, n$ ;  $n$  is the total number of individuals) for the latent trait  $\beta_0, \beta_1, \beta_2$  or  $\beta_3$ ,  $\alpha_0$  is the population mean parameter,  $x_{ij}$  is the genotypic value of individual  $i$  and marker  $j$  coded as 0, 1 and 2 for three marker genotypes AA, AB and BB, respectively,  $\alpha_j$  is the effect of marker  $j$  ( $j = 1, \dots, n$ ;  $n$  is the total number of markers), and  $\lambda$  ( $>0$ ) is a shrinkage tuning parameter. The penalty term is able to shrink the additive effects of some of the markers exactly to zero, and select a subset of the most important markers into the model. The tuning parameter  $\lambda$  determines the degree of shrinkage, and the number of markers having non-zero effects. Cross-validation is used to decide an optimal value for  $\lambda$ .

In *stability selection* (Meinshausen and Bühlmann, 2010; Alexander and Lange, 2011): (i) several bootstrap samples are first created from the original data; and (ii) frequency over-bootstrap samples on how many times each SNP is being selected to the LASSO model is monitored and used as a stable measure of variable selection. Stability selection probability (SSP) of each SNP being selected to the model was applied as a way to control the false discovery rate and determine significant SNPs (Gao et al., 2014; Li and Sillanpää, 2015). Briefly a subsample of half the number of individuals was randomly picked up and the LASSO was performed on it to select a set of markers. This procedure was repeated 1000 times. Then the selection frequency of each marker being selected was calculated, and was used to judge the support of QTL. A decision rule suggested by Meinshausen and Bühlmann (2010) was applied to control the expected number of false positives:

$$\frac{1}{2} + \frac{q^2}{2E[V]p}, \quad (5)$$

where  $q$  is the number of selected markers,  $E[V]$  is the expected number of false positives, and  $p$  is the total number of markers. For a marker to be declared as a significant QTL, a SSP inclusion frequency of at least 0.52 (i.e. derived based on formula 5) was used for all traits. This frequency was inferred conditional on the expected number of false selected markers being less than one (Bühlmann et al., 2014).

Population structure was accounted for in all analyses by including the first five PCs based on the genotype data as covariates into the model. An adaptive LASSO approach (Zou, 2006) was used to determine the percentage of phenotypic variance (PVE) ( $H^2_{QT}$ ) of all the QTLs (Methods S1). These analysis were all performed in RSTUDIO (Team, 2015).

### Candidate gene mining

To assess putative functionality of SNPs with significant associations, a gene enrichment analysis of putative genes and their associated orthologs was performed against the NORWOOD v1.0 database (<http://norwood.congenie.org>) hosted by CONGENIE (<http://congenie.org/>). The complete *P. abies* contigs that harboured the QTLs that were not annotated in the CONGENIE were

used to perform a nucleotide BLAST (BLASTN) search, using the option for only highly similar sequences (MEGABLAST) in the National Center for Biotechnology Information (NCBI) nucleotide collection database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi?>).

## CONCLUSION

This work has dissected the genetic basis of wood properties in Norway spruce with use of functional AM. In total, we identified 52 significant QTLs for wood properties and mining of candidate genes located in the vicinity of significant QTLs identified genes that could be directly or indirectly responsible for variations in the observed traits. Functional mapping analyses allowed us to utilize all the longitudinal data for a trait simultaneously and may better account for the temporal trends and correlation structures across years for the complex traits associated with wood formation. It can therefore be applied to the detection of QTLs stable over time (i.e. the QTLs associated with intercept traits) with greater statistical evidence. The slope latent trait over cambial ages or the rate of juvenile-to-mature wood transition has allowed for the dissection the dynamics of the transition process itself and can be applied to other important plant breeding traits. The significance of our results is provided by the identification of QTLs associated to both high wood density and fast growth, therefore larger biomass. These QTLs can now be a basis for future functional genomics in Norway spruce.

However, the direct use of QTLs for marker-assisted breeding has not been successful, mainly due to the difficulty in transferring the associations across populations and species of forest trees. With the small percentage variances detected and no direct information about the developmental change of QTL expression, breeders will be unable to make use of these QTLs in direct early selection. Non-additive interactions especially epistasis, play an important role in accounting for the total genetic variance of a trait. Therefore this study will be a good basis for initiating the detection and estimation of possible epistatic influence on these complex traits. Future work should focus on replicated sampling from a larger number of representative genotypes across different environments, which take into consideration genotype  $\times$  environment interactions. Additional support for marker-assisted tree breeding may also be provided by the functional genetics studies, systems mapping and consideration of biological mechanisms (Liu and Yan, 2019) of the identified candidate genes in model trees like *Populus* sp.

## DATA AVAILABILITY

All the latent traits, genotypic data, SNP position files the AM scripts used for the analysis are publicly available at zenodo.org at <https://doi.org/10.5281/zenodo.1480536>. Raw sequence data for all the samples utilized in the study are found through the European Nucleotide Archive under accession number PRJEB29652. The Norway spruce genome assemblies and resources are available from <http://congenie.org/pabiesgenome>.

## ACKNOWLEDGEMENTS

We acknowledge the support of the Bio4Energy research organization for wood property analyses and evaluations. All genetic data were obtained through funding from the Knut and Alice Wallenberg foundation. JB is supported through a postdoc position funded by the Kempe foundation.

## CONFLICT OF INTEREST

The authors declare no competing interests.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

**Figure S1.** Phenotype trajectories representing the main traits.

**Figure S2.** Significant contigs LD heatmap.

**Figure S3.** ADMIXTURE plot of the entire population.

**Figure S4.** Data are structured into three categories.

**Methods S1.** PVE evaluation of a QTL.

**Methods S2.** Trait data set used for GWAS identification of novel candidate loci affecting number of tracheids formed, radial growth, density, stiffness and mass at breast height of young Norway spruce.

**Table S1.** ConGenIE BLAST search of contigs with significant QTLs.

**Table S2.** Ring-related data (B): list of variables and examples of data.

**Table S3.** Curve shape data (A): list of variables for each property and example of data.

## REFERENCES

- Abel, S. and Theologis, A. (1996) Early Genes and Auxin Action. *Plant Physiology*. **111**(1), 9.
- Alexander, D.H. and Lange, K. (2011) Stability selection for genome-wide association. *Genet. Epidemiol.* **35**(7), 722–728. <https://doi.org/10.1002/gepi.20623>.
- Alexander, D., Novembre, J. and Lange, K. (2009) Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**(9), 1655–1664.
- Al-Tamimi, N., Brien, C., Oakey, H., Berger, B., Saade, S., Ho, Y.S., Schmöckel, S.M., Tester, M. and Negrão, S. (2016) Salinity tolerance loci revealed in rice using high-throughput non-invasive phenotyping. *Nat. Commun.* **7**, 13342. <https://doi.org/10.1038/ncomms13342>. <https://www.nature.com/articles/ncomms13342#supplementary-information>.
- Anderegg, W.R.L. (2015) Spatial and temporal variation in plant hydraulic traits and their relevance for climate change impacts on vegetation. *New Phytol.* **205**(3), 1008–1014. <https://doi.org/doi:10.1111/nph.12907>.
- Beaulieu, J., Doerksen, T., Boyle, B. et al. (2011) Association genetics of wood physical traits in the conifer white spruce and relationships with gene expression. *Genetics*, **188**(1), 197–214. <https://doi.org/10.1534/genetics.110.125781>.
- Beavis, W.D. (1998) QTL analyses: power, precision, and accuracy. *Mol Dissect Complex Traits*, **1998**, 145–162.
- Bentley, A.R., Scutari, M., Gosman, N. et al. (2014) Applying association mapping and genomic selection to the dissection of key traits in elite European wheat. *Theor. Appl. Genet.* **127**(12), 2619–2633. <https://doi.org/10.1007/s00122-014-2403-y>.
- Bergelson, J. and Roux, F. (2010) Towards identifying genes underlying ecologically relevant traits in *Arabidopsis thaliana*. *Nat. Rev. Genet.* **11**, 867. <https://doi.org/10.1038/nrg2896>.
- Bernhardsson, C., Vidalis, A., Wang, X., Scofield, D.G., Schiffthaler, B., Baisson, J., Street, N.R., García-Gil, M.R. and Ingvarsson, P.K. (2019) An ultra-dense haploid genetic map for evaluating the highly fragmented genome assembly of Norway spruce (*Picea abies*). *G3: Genes - Genomes - Genetics*, **9**(5), 1623–1632. [g3.200840.202018](https://doi.org/10.1534/g3.118.200840). <https://doi.org/10.1534/g3.118.200840>.
- Bertaud, F. and Holmbom, B. (2004) Chemical composition of earlywood and latewood in Norway spruce heartwood, sapwood and transition zone wood. *Wood Sci. Technol.* **38**(4), 245–256. <https://doi.org/10.1007/s00226-004-0241-9>.
- Bühlmann, P., Kalisch, M. and Meier, L. (2014) High-dimensional statistics with a view toward applications in biology. *Annu. Rev. Stat. Appl.* **1**(1), 255–278. <https://doi.org/10.1146/annurev-statistics-022513-115545>.
- Camargo, A.V., Mackay, I., Mott, R., Han, J., Doonan, J.H., Askew, K., Corke, F., Williams, K. and Bentley, A.R. (2018) Functional mapping of quantitative trait loci (QTLs) associated with plant performance in a wheat MAGIC mapping population. *Front. Plant Sci.* **9**, 887. <https://doi.org/10.3389/fpls.2018.00887>.

- Cenci, A. and Rouard, M. (2017) Evolutionary analyses of GRAS transcription factors in angiosperms. *Front. Plant Sci.* **8**, 273. <https://doi.org/10.3389/fpls.2017.00273>.
- Chen, Z.-Q., Gil, M.R.G., Karlsson, B., Lundqvist, S.-O., Olsson, L. and Wu, H.X. (2014) Inheritance of growth and solid wood quality traits in a large Norway spruce population tested at two locations in southern Sweden. *Tree Genet. Genomes*, **10**(5), 1291–1303.
- Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X. and Ruden, D.M. (2012) A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly*, **6**(2), 80–92.
- Cosgrove, D.J. (2005) Growth of the plant cell wall. *Nat. Rev. Mol. Cell Biol.* **6**(11), 850–861. [https://doi.org/http://www.nature.com/nrm/journal/v6/n11/supplinfo/nrm1746\\_S1.html](https://doi.org/http://www.nature.com/nrm/journal/v6/n11/supplinfo/nrm1746_S1.html).
- Cosgrove, D.J. (2016) Catalysts of plant cell wall loosening. *F1000Res.* **5**, F1000. Faculty Rev-1119. <https://doi.org/10.12688/f1000research.7180.1>
- Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T. and Sherry, S.T. (2011) The variant call format and VCFtools. *Bioinformatics*, **27**(15), 2156–2158.
- De La Torre, A.R., Puiu, D., Crepeau, M.W., Stevens, K., Salzberg, S.L., Langley, C.H. and Neale, D.B. (2019) Genomic architecture of complex traits in loblolly pine. *New Phytol.* **221**(4), 1789–1801. <https://doi.org/doi:10.1111/nph.15535>.
- Du, Q., Pan, W., Xu, B., Li, B. and Zhang, D. (2013) Polymorphic simple sequence repeat (SSR) loci within cellulose synthase (PtoCesA) genes are associated with growth and wood properties in *Populus tomentosa*. *New Phytol.* **197**(3), 763–776. <https://doi.org/10.1111/nph.12072>.
- Du, Q., Tian, J., Yang, X., Pan, W., Xu, B., Li, B., Ingvarsson, P.K. and Zhang, D. (2015) Identification of additive, dominant, and epistatic variation conferred by key genes in cellulose biosynthesis pathway in *Populus tomentosa*. *DNA Res.* **22**(1), 53–67. <https://doi.org/10.1093/dnares/dsu040>.
- Du, Q., Lu, W., Quan, M., Xiao, L., Song, F., Li, P., Zhou, D., Xie, J., Wang, L. and Zhang, D. (2018) Genome-wide association studies to improve wood properties: challenges and prospects. *Front. Plant Sci.* **9**, 1912. <https://doi.org/10.3389/fpls.2018.01912>
- Dutilleul, P., Herman, M. and Avella-Shaw, T. (1998) Growth rate effects on correlations among ring width, wood density, and mean tracheid length in Norway spruce (*Picea abies*). *Can. J. For. Res.* **28**(1), 56–68.
- Eckert, A.J., Bower, A.D., Wegrzyn, J.L., Pande, B., Jermstad, K.D., Krutovsky, K.V., Clair, J.B.S. and Neale, D.B. (2009) Association genetics of coastal Douglas fir (*Pseudotsuga menziesii* var. *menziesii*, Pinaceae). I. Cold-hardiness related traits. *Genetics*, **182**(4), 1289–1302.
- Emebiri, L.C., Devey, M.E., Matheson, A.C. and Slee, M.U. (1998) Age-related changes in the expression of QTLs for growth in radiata pine seedlings. *Theor. Appl. Genet.* **97**(7), 1053–1061. <https://doi.org/10.1007/s001220050991>.
- Evans, R. and Ilic, J. (2001) Rapid prediction of wood stiffness from microfibril angle and density. *Forest Products J.* **51**(3), 53.
- Fisher, K. and Turner, S. (2007) PXY, a receptor-like kinase essential for maintaining polarity during plant vascular-tissue development. *Curr. Biol.* **17**(12), 1061–1066.
- Gao, H., Wu, Y., Li, J., Li, H., Li, J. and Yang, R. (2014) Forward LASSO analysis for high-order interactions in genome-wide association study. *Brief. Bioinform.* **15**(4), 552–561.
- Gilmour, A., Gogel, B., Cullis, B., Welham, S., Thompson, R., Butler, D., Cherry, M., Collins, D., Dutkowsky, G. and Harding, S. (2014). ASReml user guide. Release 4.1, structural specification. *VSN International Ltd, Hemel Hempstead, HP1 1ES, UK* [www.vsnl.co.uk](http://www.vsnl.co.uk).
- Gong, C., Du, Q., Xie, J., Quan, M., Chen, B. and Zhang, D. (2018) Dissection of insertion-deletion variants within differentially expressed genes involved in wood formation in *Populus*. *Front. Plant Sci.* **8**, 2199. <https://doi.org/10.3389/fpls.2017.02199>.
- González-Martínez, S.C., Wheeler, N.C., Ersoz, E., Nelson, C.D. and Neale, D.B. (2007) Association Genetics in *Pinus taeda* L. I. Wood Property Traits. *Genetics*, **175**(1), 399–409. <https://doi.org/10.1534/genetics.106.061127>.
- Goulao, L.F., Vieira-Silva, S. and Jackson, P.A. (2011) Association of hemi-cellulose- and pectin-modifying gene expression with *Eucalyptus globulus* secondary growth. *Plant Physiol. Biochem.* **49**(8), 873–881. <https://doi.org/10.1016/j.plaphy.2011.02.020>.
- Groover, A., Devey, M., Fiddler, T., Lee, J., Megraw, R., Mitchel-Olds, T., Sherman, B., Vujcic, S., Williams, C. and Neale, D. (1994) Identification of quantitative trait loci influencing wood specific gravity in an outbred pedigree of loblolly pine. *Genetics*, **138**(4), 1293–1300.
- Hall, D., Hallingbäck, H.R. and Wu, H.X. (2016) Estimation of number and size of QTL effects in forest tree traits. *Tree Genet. Genomes*, **12**(6), 110. <https://doi.org/10.1007/s11295-016-1073-0>.
- Hallingbäck, H.R., Sánchez, L. and Wu, H.X. (2014) Single versus subdivided population strategies in breeding against an adverse genetic correlation. *Tree Genet. Genomes*, **10**(3), 605–617.
- Hamblin, M.T., Buckler, E.S. and Jannink, J.-L. (2011) Population genetics of genomics-based crop improvement methods. *Trends Genet.* **27**(3), 98–106. <https://doi.org/10.1016/j.tig.2010.12.003>.
- Hannrup, B., Cahalan, C., Chantre, G., Grabner, M., Karlsson, B., Bayon, I.L., Jones, G.L., Müller, U., Pereira, H. and Rodrigues, J.C. (2004) Genetic parameters of growth and wood quality traits in *Picea abies*. *Scand. J. For. Res.* **19**(1), 14–29.
- Hauksson, J.B., Bergqvist, G., Bergsten, U., Sjöström, M. and Edlund, U. (2001) Prediction of basic wood properties for Norway spruce. Interpretation of Near Infrared Spectroscopy data using partial least squares regression. *Wood Sci. Technol.* **35**(6), 475–485. <https://doi.org/10.1007/s00226-001-0123-3>.
- Heuven, H.C. and Janss, L.L. (2010). Bayesian multi-QTL mapping for growth curve parameters. Paper presented at the BMC proceedings, 4(1), S12.
- Hill, W. and Weir, B. (1988) Variances and covariances of squared linkage disequilibria in finite populations. *Theor. Popul. Biol.* **33**(1), 54–78.
- Hirsch, S. and Oldroyd, G.E.D. (2009) GRAS-domain transcription factors that regulate plant development. *Plant Signal. Behav.* **4**(8), 698–700.
- Hirsch, C.D., Evans, J., Buell, C.R. and Hirsch, C.N. (2014) Reduced representation approaches to interrogate genome diversity in large repetitive plant genomes. *Brief. Funct. Genomics*, **13**(4), 257–267.
- Huang, X. and Han, B. (2014) Natural variations and genome-wide association studies in crop plants. *Annu. Rev. Plant Biol.* **65**, 531–551.
- Huang, W., Richards, S., Carbone, M.A. et al. (2012) Epistasis dominates the genetic architecture of quantitative traits. *Proc. Natl Acad. Sci. USA*, **109**(39), 15553–15559. <https://doi.org/10.1073/pnas.1213423109>.
- Husson, F., Lê, S. and Pagès, J. (2017). Exploratory multivariate analysis by example using R: Chapman and Hall/CRC.
- Johannes, F., Porcher, E., Teixeira, F.K. et al. (2009) Assessing the impact of transgenerational epigenetic variation on complex traits. *PLoS Genet.* **5**(6), e1000530. <https://doi.org/10.1371/journal.pgen.1000530>.
- Khan, M.A. and Korban, S.S. (2012) Association mapping in forest trees and fruit crops. *J. Exp. Bot.* **63**(11), 4045–4060. <https://doi.org/10.1093/jxb/ers105>.
- Klein, M. and Papenbrock, J. (2004) The multi-protein family of Arabidopsis sulphotransferases and their relatives in other plant species. *J. Exp. Bot.* **55**(404), 1809–1820. <https://doi.org/10.1093/jxb/erh183>.
- Kopriva, S., Hartmann, T., Massaro, G., Hönické, P. and Rennenberg, H. (2004) Regulation of sulfate assimilation by nitrogen and sulfur nutrition in poplar trees. *Trees*, **18**(3), 320–326. <https://doi.org/10.1007/s00468-003-0309-4>.
- Lairson, L., Henrissat, B., Davies, G. and Withers, S. (2008) Glycosyltransferases: structures, functions, and mechanisms. *Annu. Rev. Biochem.* **77**, 521–555.
- Lamara, M., Raherison, E., Lenz, P., Beaulieu, J., Bousquet, J. and MacKay, J. (2016) Genetic architecture of wood properties based on association analysis and co-expression networks in white spruce. *New Phytol.* **210**(1), 240–255. <https://doi.org/doi:10.1111/nph.13762>.
- Larocque, G.R. and Marshall, P.L. (1995) Wood relative density development in red pine (*Pinus resinosa* Ait.) stands as affected by different initial spacings. *For. Sci.* **41**(4), 709–728.
- Larsson, H., Källman, T., Gyllenstrand, N. and Lascoux, M. (2013) Distribution of Long-Range Linkage Disequilibrium and Tajima's D Values in Scandinavian Populations of Norway Spruce (*Picea abies*). *G3: Genes - Genomes - Genetics*, **3**(5), 795–806. <https://doi.org/10.1534/g3.112.005462>.
- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv preprint arXiv:1303.3997*.
- Li, Z. and Sillanpää, M.J. (2013) A Bayesian nonparametric approach for mapping dynamic quantitative traits. *Genetics*, **194**(4), 997–1016.
- Li, Z. and Sillanpää, M.J. (2015) Dynamic quantitative trait locus analysis of plant phenomic data. *Trends Plant Sci.* **20**(12), 822–833. <https://doi.org/10.1016/j.tplants.2015.08.012>.



- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. and Durbin, R. (2009) The sequence alignment/map format and SAMtools. *Bioinformatics*, **25**(16), 2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>.
- Li, Z., Hallingbäck, H.R., Abrahamsson, S., Fries, A., Gull, B.A., Sillanpää, M.J. and Garcia-Gil, M.R. (2014) Functional multi-locus QTL mapping of temporal trends in Scots pine wood traits. *G3: Genes - Genomes - Genetics*, **4**(12), 2365–2379. <https://doi.org/10.1534/g3.114.014068>.
- Liu, H.-J. and Yan, J. (2019) Crop genome-wide association study: a harvest of biological relevance. *Plant J.* **97**(1), 8–18. <https://doi.org/10.1111/tpj.14139>.
- Lu, M., Krutovsky, K.V., Nelson, C.D., Koralewski, T.E., Byram, T.D. and Loopstra, C.A. (2016) Exome genotyping, linkage disequilibrium and population structure in loblolly pine (*Pinus taeda* L.). *BMC Genom.* **17**(1), 730.
- Lundqvist, S.-O., Seifert, S., Grahn, T., Olsson, L., Garcia-Gil, M.R., Karlsson, B. and Seifert, T. (2018) Age and weather effects on between and within ring variations of number, width and coarseness of tracheids and radial growth of young Norway spruce. *Eur. J. Forest Res.* **137**(5), 719–743.
- Ma, C.-X., Casella, G. and Wu, R. (2002) Functional mapping of quantitative trait loci underlying the character process: a theoretical framework. *Genetics*, **161**(4), 1751–1762.
- Mackay, T.F.C. (2013) Epistasis and quantitative traits: using model organisms to study gene–gene interactions. *Nat. Rev. Genet.* **15**, 22. <https://doi.org/10.1038/nrg3627>.
- Maloney, V.J., Samuels, A.L. and Mansfield, S.D. (2012) The endo-1, 4- $\beta$ -glucanase Korrigan exhibits functional conservation between gymnosperms and angiosperms and is required for proper cell wall formation in gymnosperms. *New Phytol.* **193**(4), 1076–1087.
- Mauriat, M. and Moritz, T. (2009) Analyses of GA20ox-and GID1-over-expressing aspen suggest that gibberellins play two distinct roles in wood formation. *Plant J.* **58**(6), 989–1003.
- McKown, A.D., Klápšte, J., Guy, R.D., Gerald, A., Porth, I., Hannemann, J., Friedmann, M., Muchero, W., Tuskan, G.A. and Ehrling, J. (2014) Genome-wide association implicates numerous genes underlying ecological trait variation in natural populations of *Populus trichocarpa*. *New Phytol.* **203**(2), 535–553. <https://doi.org/10.1111/nph.12815>.
- Meinshausen, N. and Bühlmann, P. (2010) Stability selection. *J. R. Stat. Soc. Series B Stat. Methodol.* **72**(4), 417–473. <https://doi.org/10.1111/j.1467-9868.2010.00740.x>.
- Micheli, F. (2001) Pectin methylesterases: cell wall enzymes with important roles in plant physiology. *Trends Plant Sci.* **6**(9), 414–419. [https://doi.org/10.1016/S1360-1385\(01\)02045-3](https://doi.org/10.1016/S1360-1385(01)02045-3).
- Namroud, M.-C., Guillet-Claude, C., Mackay, J., Isabel, N. and Bousquet, J. (2010) Molecular evolution of regulatory genes in spruces from different species and continents: heterogeneous patterns of linkage disequilibrium and selection but correlated recent demographic changes. *J. Mol. Evol.* **70**(4), 371–386.
- Neale, D.B. and Savolainen, O. (2004) Association genetics of complex traits in conifers. *Trends Plant Sci.* **9**(7), 325–330.
- Novaes, E., Osorio, L., Drost, D.R. et al. (2009) Quantitative genetic analysis of biomass and wood chemistry of *Populus* under different nitrogen levels. *New Phytol.* **182**(4), 878–890. <https://doi.org/10.1111/j.1469-8137.2009.02785.x>.
- Nystedt, B., Street, N.R., Wetterbom, A., Zuccolo, A., Lin, Y.-C., Scofield, D.G., Vezzi, F., Delhomme, N., Giacomello, S. and Alexeyenko, A. (2013) The Norway spruce genome sequence and conifer genome evolution. *Nature*, **497**(7451), 579–584.
- Olesen, P. (1977) The variation of the basic density level and tracheid width within the juvenile and mature wood of Norway spruce. *For. Tree Improv.* **12**, 1–22.
- Paine, C.E.T., Marthews, T.R., Vogt, D.R., Purves, D., Rees, M., Hector, A. and Turnbull, L.A. (2012) How to fit nonlinear plant growth models and calculate growth rates: an update for ecologists. *Methods Ecol. Evol.* **3**(2), 245–256. <https://doi.org/10.1111/j.2041-210X.2011.00155.x>.
- Parchman, T.L., Gompert, Z., Mudge, J., Schilkey, F.D., Benkman, C.W. and Buerkle, C. (2012) Genome-wide association genetics of an adaptive trait in lodgepole pine. *Mol. Ecol.* **21**(12), 2991–3005. <https://doi.org/10.1111/j.1365-294X.2012.05513.x>.
- Peltola, H., Gort, J., Pulkkinen, P., Gerendiain, A.Z., Karppinen, J. and Ikonen, V.-P. (2009) Differences in growth and wood density traits in Scots Pine (*Pinus sylvestris* L.) genetic entries grown at different spacing and sites. *Silva Fenn.* **43**(3), 339–354.
- Phan, T.D., Bo, W., West, G., Lycett, G.W. and Tucker, G.A. (2007) Silencing of the major salt-dependent isoform of Pectinesterase in tomato alters fruit softening. *Plant Physiol.* **144**(4), 1960–1967. <https://doi.org/10.1104/pp.107.096347>.
- Porth, I., Klápšte, J., Skyba, O. et al. (2013) Genome-wide association mapping for wood characteristics in *Populus* identifies an array of candidate single nucleotide polymorphisms. *New Phytol.* **200**(3), 710–726. <https://doi.org/10.1111/nph.12422>.
- Resende, R.T., Resende, M.D.V., Silva, F.F., Azevedo, C.F., Takahashi, E.K., Silva-Junior, O.B. and Grattapaglia, D. (2017a) Regional heritability mapping and genome-wide association identify loci for complex growth, wood and disease resistance traits in *Eucalyptus*. *New Phytol.* **213**(3), 1287–1300. <https://doi.org/10.1111/nph.14266>.
- Resende, R.T., Resende, M.D.V., Silva, F.F., Azevedo, C.F., Takahashi, E.K., Silva-Junior, O.B. and Grattapaglia, D. (2017b) Regional heritability mapping and genome-wide association identify loci for complex growth, wood and disease resistance traits in *Eucalyptus*. *New Phytol.* **213**(3), 1287–1300.
- Ruppert, D., Wand, M.P. and Carroll, R.J. (2003). *Semiparametric Regression* Vol. 12. Cambridge University Press: UK.
- Sewell, M.M., Bassoni, D.L., Megraw, R.A., Wheeler, N.C. and Neale, D.B. (2000) Identification of QTLs influencing wood property traits in loblolly pine (*Pinus taeda* L.). I. Physical wood properties. *Theor. Appl. Genet.* **101**(8), 1273–1281. <https://doi.org/10.1007/s001220051607>.
- Shiu, S.-H. and Bleeker, A.B. (2001) Plant receptor-like kinase gene family: diversity, function, and signaling. *Sci. STKE*, **2001**(113), re22.
- Sillanpää, M., Kontunen-Soppela, S., Luomala, E.-M., Sutinen, S., Kangasjärvi, J., Häggman, H. and Vapaavuori, E. (2005) Expression of senescence-associated genes in the leaves of silver birch (*Betula pendula*). *Tree Physiol.* **25**(9), 1161–1172.
- Storey, J.D., Akey, J.M. and Kruglyak, L. (2005) Multiple locus linkage analysis of genomewide expression in yeast. *PLoS Biol.* **3**(8), e267. <https://doi.org/10.1371/journal.pbio.0030267>.
- Strauss, S., Lande, R. and Namkoong, G. (1992) Limitations of molecular-marker-aided selection in forest tree breeding. *Can. J. For. Res.* **22**(7), 1050–1061.
- Team, R. (2015). RStudio: integrated development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com>.
- Thavamanikumar, S., Southerton, S.G., Bossinger, G. and Thumma, B.R. (2013) Dissection of complex traits in forest trees—opportunities for marker-assisted selection. *Tree Genet. Genomes*, **9**(3), 627–639.
- Thumma, B.R., Southerton, S.G., Bell, J.C., Owen, J.V., Henery, M.L. and Moran, G.F. (2010) Quantitative trait locus (QTL) analysis of wood quality traits in *Eucalyptus nitens*. *Tree Genet. Genomes*, **6**(2), 305–317.
- Tibshirani, R. (1996) Regression shrinkage and selection via the LASSO. *J. R. Stat. Soc. Series B Methodol.* **58**, 267–288.
- Trujillo-Moya, C., George, J., Fluch, S., Geburek, T., Grabner, M., Karanitsch-Ackerl, S., Konrad, H., Mayer, K., Sehr, E. and Wischnitzki, E. (2018) Drought sensitivity of Norway Spruce at the Species' warmest fringe: quantitative and molecular analysis reveals high genetic variation among and within provenances. *G3 (Bethesda, Md.)*, **8**(4), 1225–1245.
- Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine, A., Jordan, T., Shakir, K., Roazen, D. and Thibault, J. (2013) From FastQ data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. *Curr. Protoc. Bioinformatics*, **43**(1), 11.10.1-11.10.33.
- Verhaegen, D., Plomion, C., Gion, J.-M., Poitel, M., Costa, P. and Kremer, A. (1997) Quantitative trait dissection analysis in *Eucalyptus* using RAPD markers: 1. Detection of QTL in interspecific hybrid progeny, stability of QTL expression across different ages. *Theor. Appl. Genet.* **95**(4), 597–608. <https://doi.org/10.1007/s001220050601>.
- Vidalis, A., Scofield, D.G., Neves, L.G., Bernhardsson, C., Garcia-Gil, M.R. and Ingvarsson, P. (2018) Design and evaluation of a large sequence-capture probe set and associated SNPs for diploid and haploid samples of Norway spruce (*Picea abies*). *bioRxiv*. <https://doi.org/10.1101/291716>.
- Wegrzyn, J.L., Eckert, A.J., Choi, M., Lee, J.M., Stanton, B.J., Sykes, R., Davis, M.F., Tsai, C.-J. and Neale, D.B. (2010) Association genetics of

- traits controlling lignin and cellulose biosynthesis in black cottonwood (*Populus trichocarpa*, Salicaceae) secondary xylem. *New Phytol.* **188**(2), 515–532. <https://doi.org/doi:10.1111/j.1469-8137.2010.03415.x>.
- Wu, W.-R., Li, W.-M., Tang, D.-Z., Lu, H.-R. and Worland, A.** (1999) Time-related mapping of quantitative trait loci underlying tiller number in rice. *Genetics*, **151**(1), 297–303.
- Wu, R., Ma, C.-X., Lin, M. and Casella, G.** (2004) A general framework for analyzing the genetic architecture of developmental characteristics. *Genetics*, **166**(3), 1541–1551.
- Xing, J.U.N., Li, J., Yang, R., Zhou, X. and Xu, S.** (2012) Bayesian B-spline mapping for dynamic quantitative traits. *Genet. Res.* **94**(2), 85–95. <https://doi.org/10.1017/S0016672312000249>.
- Yamasaki, K., Kigawa, T., Inoue, M. et al.** (2004) Solution structure of the B3 DNA binding domain of the Arabidopsis cold-responsive transcription factor RAV1. *Plant Cell*, **16**(12), 3448–3459.
- Yang, R. and Xu, S.** (2007) Bayesian shrinkage analysis of quantitative trait loci for dynamic traits. *Genetics*, **176**(2), 1169–1185. <https://doi.org/10.1534/genetics.106.064279>.
- Yang, R., Tian, Q. and Xu, S.** (2006) Mapping quantitative trait loci for longitudinal traits in line crosses. *Genetics*, **173**(4), 2339–2356.
- Yang, J., Li, L., Jiang, H., Nettleton, D. and Schnable, P.S.** (2014) Dominant gene action accounts for much of the missing heritability in a gwas and provides insight into heterosis. *Genome-wide association studies to dissect the genetic architecture of yield-related traits in maize and the genetic basis of heterosis*, **1001**, 44.
- Zhong, R., Burk, D.H., Morrison, W.H. and Ye, Z.-H.** (2002) A kinesin-like protein is essential for oriented deposition of cellulose microfibrils and cell wall strength. *Plant Cell*, **14**(12), 3101–3117.
- Zhou, G., Chen, Y., Yao, W., Zhang, C., Xie, W., Hua, J., Xing, Y., Xiao, J. and Zhang, Q.** (2012) Genetic composition of yield heterosis in an elite rice hybrid. *Proc. Natl Acad. Sci. USA*, **109**(39), 15847–15852. <https://doi.org/10.1073/pnas.1214141109>.
- Zou, H.** (2006) The adaptive LASSO and its oracle properties. *J. Am. Stat. Assoc.* **101**(476), 1418–1429.