

19. Sui X, LaMonte MJ, Laditka JN, Hardin JW, Chase N, Hooker SP, Blair SN. Cardio-respiratory fitness and adiposity as mortality predictors in older adults. *JAMA* 2007; **298**:2507–2516.
20. Wannamethee SG, Shaper AG, Alberti KG. Physical activity, metabolic factors, and the incidence of coronary heart disease and type 2 diabetes. *Arch Intern Med* 2000; **160**:2108–2116.
21. Duvigneaud N, Matton L, Wijndaele K, Deriemaeker P, Lefevre J, Philippaerts R, Thomis M, Delecluse C, Duquet W. Relationship of obesity with physical activity, aerobic fitness and muscle strength in Flemish adults. *J Sports Med Phys Fitness* 2008; **48**:201–210.
22. Whaley MH, Blair SN. Epidemiology of physical activity, physical fitness and coronary heart disease. *J Cardiovasc Risk* 1995; **2**:289–295.
23. Andersen LB. Physical activity and physical fitness as protection against premature disease or death. *Scand J Med Sci Sports* 1995; **5**:318–328.
24. Rankinen T, Rauramaa R, Vaisanen S, Penttilä IM, Uusitupa M. Relation of habitual diet and cardiorespiratory fitness to blood coagulation and fibrinolytic factors. *Thromb Haemost* 1994; **71**:180–183.
25. Kullo IJ, Khaleghi M, Hensrud DD. Markers of inflammation are inversely associated with VO₂ max in asymptomatic men. *J Appl Physiol* 2007; **102**:1374–1379.
26. Weiss C, Seitel G, Bartsch P. Coagulation and fibrinolysis after moderate and very heavy exercise in healthy male subjects. *Med Sci Sports Exerc* 1998; **30**:246–251.
27. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 2005; **352**:1685–1695.

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehu249
Online publish-ahead-of-print 18 June 2014

Chest pain 9 months after interventional atrial septal defect occlusion: do not forget the worst!

Manfred Otto Vogt^{1*}, Christian Nöbauer², Christian Meierhofer¹, and Rüdiger Lange²

¹Department of Pediatric Cardiology and Congenital Heart Disease, Deutsches Herzzentrum München, Technische Universität München, Munich, Germany and

²Department of Cardiovascular Surgery, Deutsches Herzzentrum München, Technische Universität München, Munich, Germany

* Corresponding author. Tel: +49 8912183012, Fax: +49 8912183013, Email: drvogt@dhm.mhn.de

Nine months after interventional occlusion of a 27 mm atrial septal defect (ASD) with an Amplatzer 32 mm septal occluder (ASO; AGA Medical), a 45-year-old man presented with chest pain. On transthoracic echocardiography, a pericardial effusion of 5–7 mm could be demonstrated.

Transoesophageal echocardiography showed fluid at the roof of the left atrium (arrow on Panel A) towards the posterior aortic wall (Panel A: LA, left atrium; RA, right atrium; AO, Aorta; *, ASO). Under the suspicion of an erosion of the LA by ASO, a CT scan was done demonstrating the density of blood in the retroaortic fluid (Panel B: arrow, retroaortic fluid, LA, left atrium; RA, right atrium; AO, Aorta; *, ASO). The indication for a surgical intervention was established after proof of haemopericardium. Through a midline sternotomy, the chest was opened and 500 mL of blood was removed. From outside the heart, two fibrotic erosions of the LA (Panel C: *) and on the posterior aortic wall (Panel C: #) could be seen macroscopically. Inside the heart, a white strain of fibrotic tissue on the right atrial side of the ASO could be seen (Panel D: arrow). The device was explanted and the ASD closed with a patch.

Erosion of the LA roof and posterior aortic wall are possible lethal complications after ASD device occlusion with ASO. As there might be a long interval between implantation and appearance of symptoms the diagnosis might be missed. Deficient anterior superior rims and the use of oversized devices are known as risk factors. Any fluid around the heart even months/years after ASO should be a striking warning not to miss a haemopericardium.

