

RESEARCH ARTICLE

Blunt Chest Trauma in Mice after Cigarette Smoke-Exposure: Effects of Mechanical Ventilation with 100 % O₂

Katja Wagner^{1,2}, Michael Gröger¹, Oscar McCook¹, Angelika Scheuerle³, Pierre Asfar⁴, Bettina Stahl¹, Markus Huber-Lang⁵, Anita Ignatius⁶, Birgit Jung⁷, Matthias Duechs⁷, Peter Möller³, Michael Georgieff², Enrico Calzia¹, Peter Radermacher^{1*}, Florian Wagner^{1,2}

1 Institut für Anästhesiologische Pathophysiologie und Verfahrensentwicklung, Ulm, Germany, **2** Klinik für Anästhesiologie, Universitätsklinikum, Ulm, Germany, **3** Institut für Pathologie, Universitätsklinikum, Ulm, Germany, **4** Laboratoire HIFIH, UPRES EA 3859, PRES l'UNAM, IFR 132, CNRS UMR 6214, INSERM U1083, Université Angers, Département de Réanimation Médicale et de Médecine Hyperbare, Centre Hospitalier Universitaire, Angers, France, **5** Klinik für Unfall-, Hand-, Plastische und Wiederherstellungschirurgie, Universitätsklinikum, Ulm, Germany, **6** Institut für Unfallchirurgische Forschung und Biomechanik, Universitätsklinikum, Ulm, Germany, **7** Abteilung Respiratory Diseases Research, Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach/Riss, Germany



OPEN ACCESS

Citation: Wagner K, Gröger M, McCook O, Scheuerle A, Asfar P, Stahl B, et al. (2015) Blunt Chest Trauma in Mice after Cigarette Smoke-Exposure: Effects of Mechanical Ventilation with 100 % O₂. PLoS ONE 10(7): e0132810. doi:10.1371/journal.pone.0132810

Editor: Martijn van Griensven, Klinikum rechts der Isar - Technical University Munich - TUM, GERMANY

Received: May 11, 2015

Accepted: June 18, 2015

Published: July 30, 2015

Copyright: © 2015 Wagner et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: Supported by a grant of the Deutsche Forschungsgemeinschaft (Klinische Forschergruppe 200 "Die Entzündungsantwort nach muskuloskelettalem Trauma"; DFG RA 396/9-2) and a grant from the Ministry of Science, Research and the Arts of Baden-Württemberg (Az:32-729.55-0/239-5/32-7533.-6-10/15/1) to P.R. (Boehringer Ingelheim Ulm University BioCenter). The Deutsche Forschungsgemeinschaft had no role in study design, data collection and analysis, decision to publish, or

* peter.radermacher@uni-ulm.de

Abstract

Cigarette smoking (CS) aggravates post-traumatic acute lung injury and increases ventilator-induced lung injury due to more severe tissue inflammation and apoptosis. Hyperinflammation after chest trauma is due to the physical damage, the drop in alveolar PO₂, and the consecutive hypoxemia and tissue hypoxia. Therefore, we tested the hypotheses that 1) CS exposure prior to blunt chest trauma causes more severe post-traumatic inflammation and thereby aggravates lung injury, and that 2) hyperoxia may attenuate this effect. Immediately after blast wave-induced blunt chest trauma, mice (n=32) with or without 3-4 weeks of CS exposure underwent 4 hours of pressure-controlled, thoraco-pulmonary compliance-titrated, lung-protective mechanical ventilation with air or 100 % O₂. Hemodynamics, lung mechanics, gas exchange, and acid-base status were measured together with blood and tissue cytokine and chemokine concentrations, heme oxygenase-1 (HO-1), activated caspase-3, and hypoxia-inducible factor 1-α (HIF-1α) expression, nuclear factor-κB (NF-κB) activation, nitrotyrosine formation, purinergic receptor 2X₄ (P2XR₄) and 2X₇ (P2XR₇) expression, and histological scoring. CS exposure prior to chest trauma lead to higher pulmonary compliance and lower PaO₂ and Horovitz-index, associated with increased tissue IL-18 and blood MCP-1 concentrations, a 2-4-fold higher inflammatory cell infiltration, and more pronounced alveolar membrane thickening. This effect coincided with increased activated caspase-3, nitrotyrosine, P2XR₄, and P2XR₇ expression, NF-κB activation, and reduced HIF-1α expression. Hyperoxia did not further affect lung mechanics, gas exchange, pulmonary and systemic cytokine and chemokine concentrations, or histological scoring, except for some patchy alveolar edema in CS exposed mice. However, hyperoxia attenuated tissue HIF-1α, nitrotyrosine, P2XR₇, and P2XR₄ expression, while it increased HO-1 formation in CS exposed mice. Overall, CS exposure aggravated post-traumatic

preparation of the manuscript. Boehringer Ingelheim Pharma GmbH & Co. KG provided support in the form of salaries for authors B.J. and M.D., but did not have any additional role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of B. J. and M.D. are articulated in the "author contributions" section.

Competing Interests: B.J. and M.D. are employees of Boehringer Ingelheim Pharma GmbH & Co. KG, a company developing drugs for the treatment of chronic obstructive pulmonary disease (COPD). This employment does not alter the authors' adherence to PLOS ONE policies on sharing data and materials. The other authors have declared that no competing interests exist.

inflammation, nitrosative stress and thereby organ dysfunction and injury; short-term, lung-protective, hyperoxic mechanical ventilation have no major beneficial effect despite attenuation of nitrosative stress, possibly due to compensation of by regional alveolar hypoxia and/or consecutive hypoxemia, resulting in down-regulation of HIF-1 α expression.

Introduction

Blunt chest trauma is frequently associated with poly-trauma, and independently contributes to mortality if acute lung injury (ALI) develops [1]. Epidemiological data demonstrate that active or passive cigarette smoking (CS) is associated with the development of ALI after blunt trauma [2], and that active cigarette smoking increases the susceptibility to develop Acute Respiratory Distress Syndrome (ARDS) despite younger age and better overall general health status [3]. Scarce data, however, are only available in experimental animals, and the results are conflicting: in mechanically ventilated rats, pre-challenge CS exposure aggravated tissue inflammation and apoptosis, but had only marginal effects in spontaneously breathing animals [4]. Data on lung mechanics, gas exchange or histological changes were not reported in that study. Moreover, CS exposure even suppressed the pro-inflammatory responses of alveolar macrophages during halothane and isoflurane anaesthesia [5].

Lung contusion due to blunt chest trauma induced both pulmonary and systemic hyper-inflammation, oxidative stress, and enhanced apoptosis [6–8]. The pulmonary and systemic hyper-inflammatory response is due to the physical damage *per se*, the trauma-related drop in alveolar O₂ tension [6], and the consecutive hypoxemia and tissue hypoxia [9]. Chronic obstructive pulmonary disease (COPD) is also associated with pulmonary and systemic inflammation, oxidative and nitrosative stress, and apoptosis [10,11], at least in part as a result of alveolar hypoxia and hypoxemia [12–15]. Finally, in mice, CS exposure-induced COPD [16,17] lead to a similar degree of pulmonary inflammation [18] as that induced by blunt chest trauma in otherwise healthy littermates [8].

It is well-established that long-term hyperoxia causes ALI characterised by oxidative stress and enhanced cell death [19]. Scarce data, however, are only available on the interaction between CS exposure and hyperoxia: *pre*-natal exposure to the CS component benzo[*a*]pyrene potentiated immediate *post*-natal hyperoxic lung injury [20], and long-term (over five days) *post*-natal exposure to hyperoxia had an additive effect on the histological lung damage and organ dysfunction in CS-induced COPD during adulthood [21]. However, in various animal models resulting from haemorrhage [22–25], ischemia/reperfusion injury [26,27] and polymicrobial sepsis [28–30] short-term ventilation with 100% O₂ resulted in attenuated inflammation and reduced apoptosis [26,27,29–32], and ultimately improved organ function and survival. Therefore, in anesthetized, resuscitated, and mechanically ventilated mice we tested the hypotheses that 1) CS exposure prior to blunt chest trauma aggravates the post-traumatic pulmonary and systemic inflammatory response and thereby organ dysfunction and injury, and that 2) short-term hyperoxia may attenuate this effect.

Materials and Methods

The study protocol was approved by the University Animal Care Committee and the federal authorities for animal research of the Regierungspräsidium Tübingen, Baden- Württemberg, Germany (protocol no. 1046). The experiments were performed in accordance with the National Institutes of Health Guidelines on the Use of Laboratory Animals. A total of n = 36

C57BL/6J mice of either gender at an age of 10–16 weeks and weighing 23 ± 2 g were obtained from Charles River (Kisslegg, Germany), housed in isolated, ventilated cages under a 12-hours light-dark cycle, and received food and water *ad libitum*. Four of these mice did not undergo CS exposure, anesthesia, chest trauma, and surgery, and served as controls for immunoblotting and electrophoretic mobility shifts (EMSA).

Cigarette smoke-induced pulmonary inflammation

In order to address the 1st hypothesis, 16 mice were exposed to cigarette smoke (“CS”) over 3–4 weeks for 5 days per week in an exposure box as described previously [18]. Thereafter, 1 week was allowed as a recovery period prior to the blast experiment. This approach was chosen to avoid any acute stress effect induced by the CS exposure procedure *per se*. Mice received 4 cigarettes (Roth-Händle without filters, tar 10 mg, nicotine 1.0 mg, carbon monoxide 6 mg, Badische Tabakmanufaktur Roth-Händle®, Lahr, Germany) on day 1, 6 ones day 2, and 8 ones for the following days of the exposure period lasting for 15 min for each cigarette, which was followed by 8-minutes with fresh air ($15 \text{ L}\cdot\text{min}^{-1}$), and an additional 24-minutes break after each second cigarette. A semi-automatic cigarette lighter and smoke generator with an electronic timer was used to control the exposure (Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach, Germany). Particle concentration was monitored by a real time ambient particle monitor (MicroDust Pro, Casella, Amherst, NH, USA). In pilot experiments, this CS-exposure procedure had not caused any effect on the behavior, body weight, or respiratory pattern. Control animals (“Non-CS”, $n = 16$) were exposed to room air.

Anaesthesia, blast wave, surgery, and experimental protocol

Mice were anesthetized with a mixture of 2.5% sevoflurane (Sevorane, Abbott, Wiesbaden, Germany) in 50% O₂ and N₂ and buprenorphine i.p. ($1 \mu\text{g}\cdot\text{g}^{-1}$). Blunt chest trauma was induced by a single blast wave centred on the thorax as described previously [8]. Briefly, compressed air rapidly ruptures a Mylar polyester film (Du Pont de Nemur, Bad Homburg, Germany), which releases a reproducible single blast wave toward the animal’s mid-sternal chest, and thus induces a reproducible contusion of the lungs without remote organ damage. Immediately after trauma, animals received i.p. ketamine ($85 \mu\text{g}\cdot\text{g}^{-1}$), midazolam ($0.9 \mu\text{g}\cdot\text{g}^{-1}$), and fentanyl ($0.18 \mu\text{g}\cdot\text{g}^{-1}$), and were placed on a procedure bench equipped with a closed-loop system to control body temperature [8,33,34]. An incision was made in the anterior neck to expose trachea, right internal jugular vein, and right carotid artery. The trachea was intubated, and the lungs were mechanically ventilated with a pressure-controlled, lung-protective ventilation strategy using a specially designed small animal ventilator (FlexiVent, Scireq, Montreal, Canada). After a lung recruitment manoeuvre consisting of an inspiratory hold at 18 cm H₂O over 5 seconds, the initial respirator settings were: tidal volume $8 \mu\text{L}\cdot\text{g}^{-1}$, respiratory rate 150 breaths $\cdot\text{min}^{-1}$, inspiratory/expiratory time ratio 1:2, PEEP 5 cm H₂O. Recruitment manoeuvres were repeated hourly, because this approach allowed maintaining thoraco-pulmonary compliance in the physiological range by preventing atelectasis, and reduced pulmonary inflammation [35]. In order to address the 2nd hypothesis $n = 8$ animals in each group were randomly assigned to mechanical ventilation with 100% O₂ (FiO₂ 1.0), whereas the other mice were ventilated with air (FiO₂ 0.21). Catheters were inserted into the jugular vein, the carotid artery, and the bladder. Anaesthesia was maintained with continuous i.v. ketamine ($100\text{--}150 \mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$), midazolam ($0.2\text{--}0.3 \mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$), and fentanyl ($1\text{--}1.5 \mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$). Anaesthetic drugs were titrated to reach deep sedation and analgesia as documented by complete tolerance against noxious stimuli. To maintain mean arterial pressure > 50 mmHg animals received $12 \mu\text{L}\cdot\text{g}^{-1}$.

h^{-1} of hydroxyethyl starch in a balanced electrolyte solution (Tetraspan 6%, 130, 0.42, Braun Medical, Melsungen, Germany).

Measurements

All animals were studied over four hours of mechanical ventilation. Systemic haemodynamics (heart rate, mean blood pressure) and body temperature were recorded hourly. The static thoraco-pulmonary compliance was measured hourly by incrementally increasing the airway pressure up to a maximum inspiratory pressure of 20 cm H₂O and using the indwelling software of the respirator that allows automatic recording of the inspiratory and expiratory pressure-volume loop. Arterial blood samples were taken hourly for blood gases and pH. The respiratory rate was titrated to maintain arterial PCO₂ at 30–40 mmHg, the PEEP level was titrated according to the arterial PO₂: if the PaO₂/FiO₂ ratio was > 300 mmHg PEEP was decreased to 3 cmH₂O, and if 100 mmHg < PaO₂/FiO₂ < 200 mmHg, it was increased to 8 cm H₂O. At the end of the observation period, animals were killed through blood withdrawal via the carotid artery. Immediately thereafter the respiratory tubes were clamped at end-expiration, i.e. PEEP level, the thorax was opened, and the lungs were removed.

Blood and lung tissue preparation

Whole blood was immediately spun, and plasma was stored at -80°C until analysis. The right lung was sampled, immediately frozen in liquid nitrogen and stored at -80°C for cytokine measurements, immunoblotting and EMSA. The left lung was fixed in formalin and paraffin-embedded for histology and immunohistochemistry.

Cytokine concentrations

Plasma and lung tissue levels of the cytokines and chemokines tumour necrosis factor (TNF)- α , interleukin (IL)-1 β , IL-6, IL-10, IL-18, keratinocyte chemoattractant (KC), and monocyte chemoattractant protein-1 (MCP-1) were measured by a mouse multiplex cytokine kit (Bio-Plex Pro Cytokine Assay, Bio-Rad, Hercules, CA) in accordance to the manufacturer's instructions [8,33,34]. In brief, the appropriate standards and samples were added to a filter plate. The samples were incubated with antibodies chemically attached to fluorescent-labelled micro beads. Thereafter, premixed detection antibodies were added to each well, and streptavidin-phycoerythrin was added. Beads were then re-suspended, and the cytokines reaction mixture was quantified using the Bio-Plex protein array reader. Data were automatically processed and analysed by Bio-Plex Manager Software 4.1 using the standard curve produced from recombinant cytokine standards. Levels below the detection limit of the assays were set to zero for statistical purposes. Due to technical problems, plasma cytokine concentrations are from $n = 7$ animals in the "CS" groups, while lung tissue levels are from $n = 6$ and $n = 7$ of the air-ventilated "non-CS" and "CS" mice, respectively.

Cell extracts, EMSA, and immune-blot

Lung tissue was homogenized and lysed in lysing buffer. For cell extract preparation, cells were re-suspended, lysed on ice and centrifuged. The supernatant (protein extract) was stored at -80°C. For the assessment of the expression of haeme oxygenase-1 (HO-1), cleaved caspase-3 and HIF-1 α , protein concentrations were determined, and equal total protein aliquots (20–60 μg) were separated by SDS-PAGE and transferred by Western blotting [8,33]. After blocking, the membranes were incubated with commercially available primary antibodies (anti-HO-1, Abcam, Cambridge, NY; anti-cleaved caspase 3, Cell Signaling, Danvers, MA; anti-

HIF-1 α , Thermo Fisher Scientific, Waltham, MA). The primary antibodies were detected using horseradish peroxidase-conjugated secondary antibodies (Cell Signaling, Danvers, MA or Santa Cruz, Dallas, TX). The membranes were subjected to chemo-luminescence using SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific). Exposed films were scanned, and intensity of immune-reactivity was measured using NIH Image J software (<http://rsb.info.nih.gov/nih-image>). Actin and vinculin (Santa Cruz Biotechnology, Inc., Santa Cruz, CA) served as the loading controls. All immune-blot were repeated twice.

Activation of the nuclear transcription factor κ B (NF- κ B) was determined using electrophoretic mobility shift assay (EMSA) as described recently [8,33,34]: cell extracts were incubated with Poly-deoxy-inosinic-deoxy-cytidylic acid (poly-dI-dC) and 32 P-labeled double stranded oligonucleotide containing the NF- κ B (HIV κ B-site) (5' -AGT TGA GGG GAC TTT CCC AGG C-3' , Biomers, Ulm). Complexes were separated in polyacrylamide gels and exposed to X-ray films. A phosphor-imager and image analysis software (AIDA Image Analyzer, Raytest, Straubenhardt) allowed quantifying the radioactively labeled NF- κ B.

For comparison between individual gels, the intensity of each band was related to that of two loaded control animals which had not undergone surgical instrumentation and trauma. Therefore, the immune-blot and EMSA data are expressed as fold increase over the mean of the two control values using the mean value of the three individual gels for each experimental animal.

Histology and immunohistochemistry

The left lung was formalin fixed and embedded in paraffin for histological and immunohistochemistry analysis. Adapting recently published scoring systems [36] the haematoxylin eosin-stained lung sections were analysed by two experienced pathologist (A.S., P.M.) blinded for the group assignment for alveolar collapse (i.e. dystelectasis or atelectasis), emphysematous over-distension, inflammatory cell (i.e. neutrophils, macrophages, and lymphocytes, respectively) infiltration, thickening of the alveolar membranes, protein debris in the airspaces, and alveolar oedema. These parameters were scored on a scale ranging from 0 (normal lung histology without pathological findings), 0.5 (minor histological injury), 1 (moderate and patchy histological injury), 1.5 (major histological injury < 25% of the lung involved), and 2 (major histological injury 25–50% of the lung involved).

Immunohistochemistry detection of nitrotyrosine formation and expression of the purinergic receptors 2X₇ and 2X₄ (P2XR₇, P2XR₄) was performed as follows: the paraffin sections were deparaffinized in xylene and graded ethanols, boiled twice in sodium citrate buffer for heat-induced epitope retrieval before being exposed to the primary antibody (anti-nitrotyrosine, Millipore, Schwalbach, Germany; anti-P2X₇-Receptor and -P2X₄-Receptor, Alomone Labs, Jerusalem, Israel.). Primary antibody detection was performed by a secondary antibody and visualized with a red chromogen (Alkaline Phosphatase-conjugated Goat-Anti-Rabbit IgG, Jackson Immuno Research, West Grove, USA). Slides were visualized using a Zeiss Axio Imager A1 microscope using a 10x objective (EC Plan-NEOFLUAR). Four distinct random 800 μ m² square regions were quantified for intensity of signal using the image analysis AxioVision 4.8 software (Zeiss, Jena, Germany). Therefore, results are presented as densitometric sum red [8,34].

Statistical analysis

All data are presented as median (quartiles) unless otherwise stated. The sample sizes were based on our previous experience [8], for which a statistical power analysis using the Horowitz-index (PaO₂/FiO₂ ratio), thoraco-pulmonary compliance and lung tissue NF- κ B activation as

main criteria and based on two-sided testing, $\alpha = 0.05$, power 80% and non-parametric analysis of variance had yielded a minimum of $n = 8-10$ for eight experimental groups. The scope of the present study was to assess the effects of CS-exposure prior to and hyperoxia immediately after *blunt chest trauma*. Therefore, in an attempt to reduce the number of animals, we did not study sham-operated mice and used $n = 8$ in each of the four experimental groups. After exclusion of normal distribution using the Kolmogorov-Smirnov-test, intergroup differences were analysed with a Kruskal-Wallis one way ANOVA on ranks, and a subsequent Dunn's test for multiple comparisons using a two-tailed hypothesis testing. Differences were considered statistically significant when $p < 0.05$. All quantitative graphical presentations and statistical analyses were performed using the GraphPad Prism 5, version 5.04, software (GraphPad Software Inc., La Jolla, CA).

Results

Effects of cigarette smoke-exposure

[Table 1](#) demonstrates that systemic haemodynamics, metabolism, and acid-base status did not differ between the four experimental groups. However, CS exposure prior to chest trauma was associated with a higher static thoraco-pulmonary compliance post trauma and lower PaO₂ and Horovitz-index.

[Table 2](#) shows that CS exposure prior to trauma increased post-traumatic lung tissue IL-18 and blood MCP-1 concentrations, the latter, however, did not reach statistical significance ($p < 0.1$ each during air and O₂ ventilation).

[Table 3](#) demonstrates that the CS exposure-induced alterations in pulmonary gas exchange and lung cytokine concentrations coincided with a two- to four-fold higher tissue infiltration of neutrophils, smokers' macrophages, and lymphocytes, as well as with more pronounced alveolar wall thickening.

Table 1. Parameters of systemic hemodynamics, lung mechanics, pulmonary gas exchange, metabolism, and acid-base. Parameters of systemic hemodynamics, lung mechanics, pulmonary gas exchange, metabolism, and acid-base status in mice without (Non-CS) and with (CS) cigarette smoke exposure over three to four weeks prior to blunt chest trauma and after four hours mechanical ventilation with air (FiO₂ 0.21) or 100% O₂ (FiO₂ 1.0) ($n = 8$ in each group). All data are median (quartiles).

	Non-CS FiO ₂ 0.21	CS FiO ₂ 0.21	Non-CS FiO ₂ 1.0	CS FiO ₂ 1.0	p-value K-W ANOVA
Heart rate [min ⁻¹]	340 (340;348)	355 (320;375)	395 (364;443)	322 (318;350)	0.089
Mean arterial pressure [mmHg]	54 (54;58)	56 (52;57)	57 (56;60)	58 (55;60)	0.326
Glucose [mmol·L ⁻¹]	5.7 (5.4;6.6)	4.5 (3.7;4.8) §	5.0 (4.4;6.4)	5.2 (4.8;5.8)	0.024
Lactate [mmol·L ⁻¹]	1.8 (1.1;2.0)	1.1 (0.9;1.1)	1.2 (0.8;1.3)	0.9(0.8;1.5)	0.199
Minute ventilation [L·min ⁻¹]	750 (745;810)	810 (768;828)	800 (760;830)	750 (748;758)	0.227
Compliance [L·cm H ₂ O ⁻¹]	79 (75;93)	97 § (82;106)	75 (65;85)	90 § (87;96)	0.028
PaO ₂ [mmHg]	102 (97;106)	78 § (75;82)	379 § (318;395)	370 § (329;424)	< 0.001
PEEP [cm H ₂ O]	3.0 (3.0;3.0)	3.0 (3.0;3.5)	3.0 (3.0;3.0)	3.0 (3.0;3.5)	0.272
PaO ₂ /FiO ₂ ratio [mmHg]	491 (460;505)	368 § (358;391)	379 § (318;395)	370 (329;424)	0.007
PaCO ₂ [mmHg]	35 (32;38)	32 (32;35)	37 (29;37)	37 (34;43)	0.460
Arterial pH	7.32 (7.30;7.34)	7.30 (7.23;7.33)	7.32 (7.31;7.35)	7.30 (7.28;7.34)	0.565
Arterial base excess [mmol·L ⁻¹]	-7.4 (-8.5;-6.9)	-8.1 (-9.7;-6.8)	-6.8 (-8.0;-6.5)	-7.0 (-10.0;-5.4)	0.617

§ depicts $p < 0.05$ vs. the corresponding Non-CS group,

§ depicts $p < 0.05$ vs. the corresponding air group (K-W ANOVA Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.t001

Table 2. Plasma and lung tissue cytokine and chemokine concentrations. Plasma (in pg·mL⁻¹) and lung tissue (in pg·mg_{protein}⁻¹) cytokine and chemokine concentrations at the end of the four-hours observation period (n = 8 in each group) obtained from mice without (Non-CS) and with (CS) cigarette smoke exposure over 3–4 weeks prior to blunt chest trauma and mechanical ventilation with air (FiO₂ 0.21) or 100% O₂ (FiO₂ 1.0). All data are median (quartiles).

	Non-CS FiO ₂ 0.21	CS FiO ₂ 0.21	Non-CS FiO ₂ 1.0	CS FiO ₂ 1.0	p-value K-W ANOVA
IL-1β [pg·mL ⁻¹]	32 (25;36)	37 (22;49)	23 (13;29)	13 (9;19)	0.215
IL-1β [pg·mg _{protein} ⁻¹]	309 (209;457)	531 (204;631)	223 (118;332)	285 (245;325)	0.413
IL-6 [pg·mL ⁻¹]	212 (71;475)	511 (310;863)	123 (93;275)	177 (122;434)	0.304
IL-6 [pg·mg _{protein} ⁻¹]	5 (5;20)	9 (8;11)	11 (8;18)	5 (4;7)	0.116
IL-10 [pg·mL ⁻¹]	99 (72;171)	153 (92;255)	85 (53;126)	95 (80;177)	0.467
IL-10 [pg·mg _{protein} ⁻¹]	47 (43;53)	38 (34;40)	42 (39;45)	48 (45;51)	0.153
KC [pg·mL ⁻¹]	275 (146;421)	265 (156;644)	115 (97;136)	223 (212;324)	0.369
KC [pg·mg _{protein} ⁻¹]	570 (313;833)	606 (469;671)	320 (224;497)	450 (391;586)	0.204
MCP-1 [pg·mL ⁻¹]	772 (528;2517)	1395 (889;7011)	265 (208;779)	3060 (1859;3389)	0.099
MCP-1 [pg·mg _{protein} ⁻¹]	141 (135;156)	137 (123;166)	136 (114;150)	129 (121;135)	0.622
TNF-α [pg·mL ⁻¹]	90 (84;104)	111 (82;137)	86 (61;93)	93 (86;108)	0.456
TNF-α [pg·mg _{protein} ⁻¹]	113 (75;133)	113 (102;121)	122 (111;145)	113 (88;144)	0.707
IL-18 [pg·mL ⁻¹]	234 (165;464)	189 (169;217)	179 (147;220)	148 (105;208)	0.429
IL-18 [pg·mg _{protein} ⁻¹]	603 (533;673)	973 (795;1045) §	563 (440;600)	882 (629;1088) §	0.019

§ depicts p < 0.05 vs. the corresponding Non-CS group,

§ depicts p < 0.05 vs. the corresponding air group (K-W ANOVA Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.t002

Typical examples of the histological items are shown in Fig 1. Whereas CS exposure did not affect post-traumatic lung tissue HO-1 expression (Fig 2), it markedly increased activated caspase-3 (Fig 3) expression and NF-κB activation (Fig 4), and reduced HIF-1α formation (Fig 5). CS exposure was also associated with increased tissue nitrotyrosine formation (Fig 6), P2XR₇ and P2XR₄ expression (Figs 7 and 8).

Table 3. Parameters of lung histopathology. Parameters of lung histopathology at the end of the four-hours observation period (n = 8 in each group) obtained from mice without (Non-CS) and with (CS) cigarette smoke exposure over 3–4 weeks prior to blunt chest trauma and mechanical ventilation with air (FiO₂ 0.21) or 100% O₂ (FiO₂ 1.0) (n = 8 in each group). Parameters were scored as 0 (normal lung histology without pathological findings), 0.5 (minor histological injury), 1 (moderate and patchy histological injury), 1.5 (major histological injury < 25% of the lung involved), and 2 (major histological injury 25–50% of the lung involved). All data are median (quartiles).

	Non-CS FiO ₂ 0.21	CS FiO ₂ 0.21	Non-CS FiO ₂ 1.0	CS FiO ₂ 1.0	p-value K-W ANOVA
Dystelectasis/atelectasis	1.0 (0.5;1.0)	0.5 (0.5;1.0)	1.0 (0.5;1.0)	0.5 (0.5;0.6)	0.683
Emphysema	1.5 (1.5;2.0)	1.5 (1.0;1.5)	1.3 (1.0;1.5)	1.1 § (1.0;1.3)	0.011
Alveolar membrane thickening	1.0 (1.0;1.0)	1.5 § (1.5;1.8)	1.0 (1.0;1.5)	1.5 § (1.4;1.6)	0.047
Lymphocytes	1.0 (1.0;1.0)	2.0 § (1.8;2.3)	1.5 (1.5;2.0)	2.0 § (1.9;2.5)	< 0.001
Neutrophils	0.5 (0.5;0.8)	2.0 § (2.0;2.5)	0.8 (0.5;1.0)	1.5 § (1.5;2.0)	< 0.001
Macrophages	3.5 (3.3;3.8)	12.0 § (7.0;13.5)	3.5 (3.0;5.6)	14.0 § (11.5;15.8)	< 0.001
Protein debris in the airspaces	1.0 (0.5;1.0)	0.5 (0.5;0.8)	1.0 (1.0;1.0)	0.8 (0.5;1.0)	0.074
Alveolar edema	0.5 (0.3;0.5)	0.5 (0.5;1.0)	1.0 (0.9;1.1)	1.5 § (0.9;2.0)	0.022

§ depicts p < 0.05 vs. the corresponding Non-CS group,

§ depicts p < 0.05 vs. the corresponding air group (K-W ANOVA Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.t003




Fig 1. Typical examples of the histopathological items scored. *Upper panel:* Lung region with little histopathological abnormalities, i.e. no dystelectasis/ atelectasis, normal thickness of alveolar membranes, and little lymphocyte immigration. Dotted arrows show some degree of protein debris in the airspaces. *Lower panel:* Lung region with major histological injury (alveolar membrane thickening and lymphocyte immigration grade 1.5 each). Solid arrows show alveolar smokers' macrophages within the airways.

doi:10.1371/journal.pone.0132810.g001

Effects of hyperoxia

Tables 1 and 2 demonstrate that lung-protective mechanical ventilation at FiO_2 1.0 did not further affect lung mechanics, gas exchange or pulmonary and systemic cytokine and chemokine concentrations. Table 3 shows that ventilation with 100% O_2 did not affect lung histology except for some patchy alveolar edema in CS exposed mice, however, without any deleterious effect on organ function (see Table 1). While ventilation with 100% O_2 markedly increased lung tissue HO-1 expression (Fig 2) in CS exposed mice, it did not further affect activated caspase-3 (Fig 3) and HIF-1 α expression (Fig 4), or NF- κ B activation (Fig 5), and even attenuated nitrotyrosine formation (Fig 6), and P2XR₇, and P2XR₄ expression (Figs 7 and 8).




Fig 2. Results of the immune blotting for HO-1. Original western blots and quantitative analysis of lung tissue expression of HO-1 from mice without (open boxplots; n = 8 each) and with (hatched boxplots; n = 7 each) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) together with two blots each (right part of blot panel) from control animals that did not undergo cigarette smoke exposure, anaesthesia, chest trauma, and surgery. All data are median (quartiles, range) as fold increase over values from control animals; § p < 0.05 vs. corresponding cigarette smoke exposure group, \$ p < 0.05 vs. corresponding air ventilation group (Kruskall-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g002

Discussion

This study investigated the effect of short-term hyperoxia on post-traumatic pulmonary and systemic inflammation and lung injury in mice that had undergone CS exposure prior to blunt chest trauma. The major findings were that i) CS exposure prior to trauma aggravated trauma-induced lung dysfunction and organ injury due to more pronounced pulmonary inflammation and nitrosative stress, ultimately resulting in enhanced apoptosis and tissue damage, and that ii) lung-protective mechanical ventilation with 100% O₂ partially attenuated this effect.

Effects of cigarette smoke-exposure on the acute trauma response

Smoking is directly related to the development of ALI after severe blunt trauma [2]. Our findings well agree with this observation: mice that had undergone CS exposure prior to trauma showed impaired pulmonary gas exchange and more pronounced lung histological damage. Despite the comparable histological evidence of emphysematous lung over-distension, the higher static thoraco-pulmonary compliance in the CS exposed mice further supports this notion: CS abuse is the most important etiological factor for COPD [37], which in turn, is




Fig 3. Results of the immune blotting for activated caspase-3. Original western blots and quantitative analysis of lung tissue expression of activated caspase-3 from mice without (open boxplots; n = 8 each) and with (hatched boxplots; n = 7 each) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) together with two blots each (right part of blot panel) from control animals that did not undergo cigarette smoke exposure, anaesthesia, chest trauma, and surgery. All data are median (quartiles, range) as fold increase over values from control animals; \$ p < 0.05 vs. corresponding cigarette smoke exposure group, \$ p < 0.05 vs. corresponding air ventilation group (Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g003

associated with a hyperinflation-induced increase of the static compliance [38], even when total lung capacity is still normal [39].

Murine models of CS-induced COPD are characterised by pulmonary inflammation and mediator release: after four days of CS-exposure, animals presented with a several-fold increase of neutrophil accumulation and pro-inflammatory cytokine and chemokine concentrations in the bronchoalveolar lavage (BAL) fluid [18,40]. In addition, CS exposure caused a two- to three-fold increase of markers of oxidative and nitrosative stress in the BAL fluid [41]. This effect resulted in enhanced apoptosis as documented by increased expression of the activated caspase-3 [41,42]. Finally, CS exposure increased ventilator-induced lung injury due to aggravated tissue inflammation and apoptosis in alveolar type II cells [4]. Our findings well agree with these observations: despite the 1-week recovery period aiming to avoid any procedure-related stress, CS-exposed mice presented with higher NF-κB activation, nitrotyrosine formation, and tissue IL-18 concentrations. The latter finding is of particular interest in the context of the CS exposure-related higher expression of the P2XR₇ and P2XR₄: inflammation-regulated production of IL-1β and IL-18 has been referred to as a key mechanism of injurious ventilation- [43] and hyperoxia-induced [44] ALI *via* P2XR₇ and subsequent inflammasome activation and IL-1β secretion: Genetic P2XR₇ deletion reduced the endotoxin-induced release of IL-1β and thereby attenuated the impairment of lung mechanics and the




Fig 4. Results of the EMSA for NF-κ activation. Original gel shifts and quantitative analysis of lung tissue expression of NF-κB activation from mice without (open boxplots; n = 8 each) and with (hatched boxplots; n = 7 each) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) together with two blots each (right part of blot panel) from control animals that did not undergo cigarette smoke exposure, anaesthesia, chest trauma, and surgery. All data are median (quartiles, range) as fold increase over values from control animals; § p < 0.05 vs. corresponding cigarette smoke exposure group, § p < 0.05 vs. corresponding air ventilation group (Kruskall-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g004

histological organ damage *in vivo* [45], and P2XR₇ activation aggravated the endotoxin-related vascular hypo-reactivity *in vitro* [46]. Co-expression of the P2XR₄ enhanced the P2XR₇-related inflammatory response [47], and up-regulation of P2XR₄ was referred to compensate for P2XR₇ depletion [48]. Finally, in addition to their role for the development of ALI, P2XR₇ and P2XR₄ activation was shown to assume major importance for CS-related lung injury: CS exposure caused up-regulation of both the P2XR₄ and P2XR₇ [49], and either pharmacological blockade or genetic deletion of the P2XR₇ attenuated the pulmonary IL-1β and IL-18 accumulation after CS exposure [50]. Consequently, our experiments support P2XR₄ and P2XR₇ as potential therapeutic targets both in ALI [51] and CS-induced COPD [52,53].

Pre-traumatic CS exposure reduced lung tissue HIF-1α expression after chest trauma. At first glance, this finding is in contrast to data from both CS exposed animals [54] and patients with COPD [55,56]. The increased nitrosative stress in CS exposed mice may have assumed importance in this context: under normoxic conditions, hyper-inflammation-induced nitric oxide (NO) release impairs HIF-1α degradation due to inhibition of prolyl-hydroxylases (PHD) [57]. However, excess NO formation under hypoxic conditions may also reduce HIF-1α accumulation because of an NO-mediated feedback with expression of PHD and/or O₂-redistribution to PHD resulting from NO-induced inhibition of mitochondrial respiration [58].




Fig 5. Results of the immune blotting for HIF-1 α . Original western blots and quantitative analysis of lung tissue expression of HIF-1 α expression from mice without (open boxplots; n = 8 each) and with (hatched boxplots; n = 7 each) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) together with two blots each (right part of blot panel) from control animals that did not undergo cigarette smoke exposure, anaesthesia, chest trauma, and surgery. All data are median (quartiles, range) as fold increase over values from control animals; § p < 0.05 vs. corresponding cigarette smoke exposure group, \$ p < 0.05 vs. corresponding air ventilation group (Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g005

Effects of hyperoxia on the acute trauma response

In mice without pre-traumatic CS exposure, pure O₂ ventilation was associated with a significantly lower PaO₂/FiO₂ ratio than in the corresponding air-ventilated animals. This effect was most likely due to O₂ breathing-induced instability of lung regions with low ventilation/perfusion-ratios [59] rather than due to O₂ toxicity: O₂ ventilation i) did not affect gas exchange or lung mechanics in mice after CS exposure, and ii) was not associated with any biochemical or morphological sign of aggravated pulmonary inflammation. Equivocal data are available whether short-term mechanical ventilation with 100% O₂ induces a pulmonary and systemic inflammatory response: in rats, 120 minutes of pure O₂ ventilation caused a marked increase in extravascular lung water and the BAL fluid neutrophil, chemokine and cytokine content, but this effect required the use of injurious tidal volumes (20 mL·kg⁻¹). In contrast, tidal volumes similar to our study (7 mL·kg⁻¹) had no effect [60]. In rabbits, four hours of mechanical ventilation with 100% O₂ significantly increased BAL polymorphonuclear leukocytes and MCP-1 concentrations, associated with increased alveolar-capillary permeability. Again, this effect was only present with high (25 mL·kg⁻¹) tidal volumes, while clinically more relevant (10 mL·kg⁻¹) tidal volumes had no effects [61]. Finally, in mice incrementally increasing the FiO₂ during 120 minutes of mechanical ventilation (tidal volume 7–8 mL·kg⁻¹) did not alter airway resistance, tissue




Fig 6. Results of the immunohistochemistry for nitrotyrosine. Typical examples (upper panel) and quantitative analysis (lower panel) of immunohistochemistry for lung tissue nitrotyrosine formation from mice without (dotted boxplots) and with (hatched boxplots) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) (n = 8 in each group). All data median (quartiles, range), \$ p < 0.05 vs. corresponding cigarette smoke exposure group, \$ p < 0.05 vs. corresponding air ventilation group (Kruskall-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g006

elastance nor BAL fluid concentrations of chemokines and cytokines [62]. Scarce data are only available on the effects of non-injurious mechanical ventilation with O₂ in the presence of sepsis- or trauma-induced ALI: low-tidal volume (7 mL·kg⁻¹) mechanical ventilation with 100% O₂ caused more pronounced pulmonary and systemic inflammation than spontaneous air breathing, but the effects of hyperoxia *per se* remain open, since mechanically ventilation with lower FiO₂ values was not studied [63]. In swine with fecal peritonitis-induced septic shock, lung-protective, pure O₂ mechanical ventilation up to 24 hours did not cause any aggravation of pulmonary or systemic inflammation or histological damage [29,30].

Hyperoxia may have attenuated pulmonary and systemic inflammation by compensating regional pulmonary hypoxia as well as tissue hypoxia resulting from chest trauma-induced hypoxemia: hypoxic hypoxia (FiO₂ 0.1) of incremental duration caused a time-dependent increase of the BAL fluid neutrophil count and albumin content reflecting alveolar-capillary leakage [64], and further aggravated endotoxin-induced ALI [65]. MCP-1 is a crucial mediator of this inflammatory response [66], and in our experiments, blood MCP-1 levels were three




Fig 7. Results of the immunohistochemistry for P2X₇. Typical examples (upper panel) and quantitative analysis (lower panel) of immunohistochemistry for lung tissue expression of the purinergic receptor P2X₇ from mice without (dotted boxplots) and with (hatched boxplots) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) (n = 8 in each group). All data median (quartiles, range), § p < 0.05 vs. corresponding cigarette smoke exposure group, \$ p < 0.05 vs. corresponding air ventilation group (Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g007

times lower in pure O₂-ventilated mice without CS exposure prior to trauma (772 (528;2517) vs. 265 (208;779) pg·mL⁻¹, p = 0.08). Pure O₂ ventilation was also associated with reduced lung tissue nitrotyrosine formation, no matter whether mice had been exposed to CS or not. This finding well agrees with previous findings on the effects of short-term exposure to hyperoxia on oxidative and nitrosative stress: two or three hours of intermittent O₂ breathing after zymosan injection increased the activity of tissue antioxidant enzymes [31], and long-term exposure to FiO₂ 0.8 attenuated lung nitrotyrosine formation with intra-tracheal carrageenan-induced pneumonitis [67]. Hence, our findings suggest that under acute stress, short-term hyperoxia can counteract the nitrosative stress associated with CS exposure-induced COPD. Hyperoxia attenuated the P2XR₇ expression both in mice with and without CS exposure prior to chest trauma, whereas it reduced P2XR₄ expression only in animals with CS exposure. Hence, short-term hyperoxia may be of particular interest in acute stress under conditions of chronic hypox (aem)ia: hypoxic hypoxaemia induced by exposure to simulated altitude not only increased lung tissue P2XR₄ mRNA, but also in the right ventricle, which went alongside with pulmonary artery hypertension and consecutive right ventricular hypertrophy [68].




Fig 8. Results of the immunohistochemistry for P2X₄. Typical examples (upper panel) and quantitative analysis (lower panel) of immunohistochemistry for lung tissue expression of the purinergic receptor P2X₄ from mice without (dotted boxplots) and with (hatched boxplots) cigarette smoke exposure prior to blunt chest trauma and mechanically ventilated with air (white boxplots) and 100% O₂ (grey boxplots) (n = 8 in each group). All data median (quartiles, range), § p < 0.05 vs. corresponding cigarette smoke exposure group, \$ p < 0.05 vs. corresponding air ventilation group (Kruskal-Wallis analysis of variance on ranks with post-hoc Dunn's test for multiple comparisons).

doi:10.1371/journal.pone.0132810.g008

Albeit we lack a direct proof, it is tempting to speculate that the hyperoxia-related attenuation of pulmonary and systemic inflammation was due to down-regulation of HIF-1 α expression: Both genetic deletion [69,70] and pharmacological blockade [71] of HIF-1 α attenuated the local and systemic inflammatory response and thereby reduced the severity of ALI after remote [69,70] or direct [71] organ damage. Finally, the neuro-protective effect of exposure to both hyper- and normobaric O₂ was associated with a two to three-fold reduction of HIF-1 α expression [72].

Limitations of the study

It could be argued that despite the marked differences in the inflammatory response, the CS-related effects on lung mechanics, gas exchange, and histology lack physiological significance: overall they were moderate, and the Horowitz-index always remained above 300 mmHg, i.e. above the definition threshold of ALI. Of note, we [8,51] and others [73] found a similar dissociation between significantly increased levels of inflammatory biomarkers in the lung and

lacking effects on lung mechanics and gas exchange in mice after chest trauma [8,51] and polymicrobial sepsis [73]. Only few studies reported Horovitz-indices compatible with the definition of ALI (< 300 mmHg) in mechanically ventilated mice: animals were either ventilated with injurious tidal volumes ($12\text{--}40$ mL \cdot kg $^{-1}$) [74–78], or 24 hours after injection of endotoxin, i.e. in the presence of prolonged, severe ALI [79]. Moreover, we used pressure-controlled, lung-protective ventilation that also comprised an initial lung recruitment manoeuvre. Consequently, any further damage beyond the effect of lung contusion *per se* or due to injurious mechanical ventilation was avoided. In fact, neither pulmonary mechanics nor gas exchange deteriorated during the 4-hour observation period. Our approach is in contrast to previous experiments, which explicitly studied the effect of CS on tissue inflammation and apoptosis related to ventilator-induced lung injury [4]. However, in that study, data on lung mechanics, gas exchange or histological changes were not reported at all.

Clearly, the short duration of the mechanical ventilation precludes any conclusion on the long-term effects. However, a recent review article showed that only two studies described mechanical ventilation in mice of up to eight hours, while in the other reports only four to six hours of mechanical ventilation were used [80]. Finally, we cannot exclude that correcting trauma-related hypoxaemia using an FiO_2 of 0.3–0.4 rather than air ventilation in the control group may have produced similar or even better results when compared to the hyperoxia group.

Conclusion

In a murine model of COPD induced by CS exposure, pulmonary and systemic inflammation as well as nitrosative stress were aggravated after blunt chest trauma, which ultimately resulted in increased severity of post-traumatic organ dysfunction and injury as documented by impaired gas exchange and more pronounced histological damage. Overall, short-term, lung-protective mechanical ventilation with 100% O_2 did not have a major therapeutic effect despite attenuation of nitrosative stress. The latter was possibly due to correction of regional alveolar hypoxia and/or consecutive hypoxemia, resulting in HIF-1 α down-regulation.

Acknowledgments

Very special thanks are dedicated to Rosemarie Mayer and Rosa Maria Engelhardt for their skillful technical assistance.

Author Contributions

Conceived and designed the experiments: KW PA MH AI BJ M. Georgieff EC FW PR. Performed the experiments: KW M. Gröger FW. Analyzed the data: KW OM AS M. Gröger BS FW. Contributed reagents/materials/analysis tools: AS OM MH BS MD. Wrote the paper: KW PA AI PM M. Georgieff BJ FW PR. Cigarette smoke exposure: BJ MD.

References

1. Shah CV, Locallo AR, Lanken PN, Kahn JM, Bellamy S, Gallop R, et al. The impact of development of acute lung injury on hospital mortality in critically ill trauma patients. *Crit Care Med* 2008; 36:2309–2315 doi: [10.1097/CCM.0b013e318180dc74](https://doi.org/10.1097/CCM.0b013e318180dc74) PMID: [18664786](https://pubmed.ncbi.nlm.nih.gov/18664786/)
2. Calfee CS, Matthay MA, Eisner MD, Benowitz N, Mariah C, Pittet JF, et al. Active and passive cigarette smoking and acute lung injury after severe blunt trauma. *Am J Respir Crit Care Med* 2011; 183:160–165
3. Hsieh SJ, Zhuo H, Benowitz NL, Thompson BT, Liu KD, Matthay MA, et al. Prevalence and impact of active and passive cigarette smoking in acute respiratory distress syndrome. *Crit Care Med* 2014; 42:2058–2068 doi: [10.1097/CCM.0000000000000418](https://doi.org/10.1097/CCM.0000000000000418) PMID: [24942512](https://pubmed.ncbi.nlm.nih.gov/24942512/)

4. Hirsch J, Chalkley RJ, Bentley T, Burlingame AL, Frank JA. Double impact of cigarette smoke and mechanical ventilation on the alveolar epithelial type II cell. *Crit Care* 2014; 18:R50 doi: [10.1186/cc13795](https://doi.org/10.1186/cc13795) PMID: [24666941](https://pubmed.ncbi.nlm.nih.gov/24666941/)
5. Kotani N, Hashimoto H, Sessler DI, Yatsu Y, Muraoka M, Matsuki A. Exposure to cigarette smoke impairs alveolar macrophage functions during halothane and isoflurane anesthesia in rats. *Anesthesiology* 1999; 91:1823–1833 PMID: [10598627](https://pubmed.ncbi.nlm.nih.gov/10598627/)
6. Knöferl MW, Liner UC, Seitz DH, Perl M, Brückner UB, Kinzl L, et al. Cardiopulmonary, histological, and inflammatory alterations after lung contusion in a novel mouse model of blunt chest trauma. *Shock* 2003; 19:519–525 PMID: [12785006](https://pubmed.ncbi.nlm.nih.gov/12785006/)
7. Hoth JJ, Welss JD, Brownlee NA, Hiltbold EM, Meredeith JW, McCall CE, et al. Toll-like receptor 4-dependent responses to lung injury in a murine model of pulmonary contusion. *Shock* 2009; 31:376–381 doi: [10.1097/SHK.0b013e3181862279](https://doi.org/10.1097/SHK.0b013e3181862279) PMID: [18665044](https://pubmed.ncbi.nlm.nih.gov/18665044/)
8. Wagner F, Scheuerle A, Weber S, Stahl B, McCook O, Knöferl MW, et al. Cardiopulmonary, histologic, and inflammatory effects of intravenous Na₂S after blunt chest trauma-induced lung contusion in mice. *J Trauma* 2011; 71:1659–1667 doi: [10.1097/TA.0b013e318228842e](https://doi.org/10.1097/TA.0b013e318228842e) PMID: [21857260](https://pubmed.ncbi.nlm.nih.gov/21857260/)
9. Eltzschig HK, Carmeliet P. Hypoxia and inflammation. *N Engl J Med* 2011; 364:656–665 doi: [10.1056/NEJMr0910283](https://doi.org/10.1056/NEJMr0910283) PMID: [21323543](https://pubmed.ncbi.nlm.nih.gov/21323543/)
10. Tuder RM, Petrache I. Pathogenesis of chronic obstructive pulmonary disease. *J Clin Invest* 2012; 122:2749–2755
11. Schmidt EP, Tuder RM. Role of apoptosis in amplifying inflammatory responses in lung diseases. *J Cell Death* 2010; 2010:41–53 PMID: [22081757](https://pubmed.ncbi.nlm.nih.gov/22081757/)
12. Seimetz M, Parajuli N, Pichl A, Veit F, Kwapiszewska G, Weisel FC, et al. Inducible NOS inhibition reverses tobacco-smoke-induced emphysema and pulmonary hypertension in mice. *Cell* 2011; 147:293–305 doi: [10.1016/j.cell.2011.08.035](https://doi.org/10.1016/j.cell.2011.08.035) PMID: [22000010](https://pubmed.ncbi.nlm.nih.gov/22000010/)
13. Chao J, Wood JG, Gonzalez NC. Alveolar hypoxia, alveolar macrophages, and systemic inflammation. *Respir Res* 2009; 10:54 doi: [10.1186/1465-9921-10-54](https://doi.org/10.1186/1465-9921-10-54) PMID: [19545431](https://pubmed.ncbi.nlm.nih.gov/19545431/)
14. Gonzalez NC, Wood JG. Alveolar hypoxia-induced systemic inflammation: what low PO₂ does and does not do. *Adv Exp Med Biol* 2010; 662:27–32 doi: [10.1007/978-1-4419-1241-1_3](https://doi.org/10.1007/978-1-4419-1241-1_3) PMID: [20204767](https://pubmed.ncbi.nlm.nih.gov/20204767/)
15. Fröhlich S, Boylan J, McLoughlin P. Hypoxia-induced inflammation in the lung? A potential therapeutic target in acute lung injury? *Am J Respir Cell Mol Biol* 2013; 48:271–279 doi: [10.1165/rcmb.2012-0137TR](https://doi.org/10.1165/rcmb.2012-0137TR) PMID: [23087053](https://pubmed.ncbi.nlm.nih.gov/23087053/)
16. Vlahos R, Bozinovski S, Gualano RC, Ernst M, Anderson GP. Modelling COPD in mice. *Pulm Pharmacol Ther* 2006; 19:12–17 PMID: [16286233](https://pubmed.ncbi.nlm.nih.gov/16286233/)
17. Gould NS, Min E, Gauthier S, Chu HW, Martin R, Day BJ. Aging adversely affects the cigarette smoke-induced glutathione adaptive response in the lung. *Am J Respir Crit Care Med* 2010; 182:1114–1122 doi: [10.1164/rccm.201003-0442OC](https://doi.org/10.1164/rccm.201003-0442OC) PMID: [20622027](https://pubmed.ncbi.nlm.nih.gov/20622027/)
18. Wollin L, Pieper M. Tiotropium bromide exerts anti-inflammatory activity in a cigarette smoke mouse model of COPD. *Pulm Pharmacol Ther* 2010; 23:345–354 doi: [10.1016/j.pupt.2010.03.008](https://doi.org/10.1016/j.pupt.2010.03.008) PMID: [20362689](https://pubmed.ncbi.nlm.nih.gov/20362689/)
19. Kallett RH, Matthay MA. Hyperoxic acute lung injury. *Respir Care* 2013; 58:123–141 doi: [10.4187/respcare.01963](https://doi.org/10.4187/respcare.01963) PMID: [23271823](https://pubmed.ncbi.nlm.nih.gov/23271823/)
20. Thakur VS, Liang YW, Lingappan K, Jiang W, Wang L, Barrios R, et al. Increased susceptibility to hyperoxic lung injury and alveolar simplification in newborn rats by prenatal administration of benzo[a]pyrene. *Toxicol Lett* 2014; 230:322–332 PMID: [24657529](https://pubmed.ncbi.nlm.nih.gov/24657529/)
21. McGrath-Morrow SA, Lauer T, Collaco JM, Yee M, O'Reilly M, Mitzner W, et al. Neonatal hyperoxia contributes additively to cigarette smoke-induced chronic obstructive pulmonary disease changes in adult mice. *Am J Respir Cell Mol Biol* 2011; 45:610–616 doi: [10.1165/rcmb.2010-0259OC](https://doi.org/10.1165/rcmb.2010-0259OC) PMID: [21239606](https://pubmed.ncbi.nlm.nih.gov/21239606/)
22. Bitterman H, Brod V, Weisz G, Kushnir D, Bitterman N. Effects of oxygen on regional hemodynamics in hemorrhagic shock. *Am J Physiol Heart Circ Physiol* 1996; 271:H203–H211
23. Efrati S, Berman S, Aharon GB, Siman-Tov Y, Averbukh Z, Weissgarten J. Application of normobaric hyperoxia therapy for amelioration of haemorrhagic shock-induced acute renal failure. *Nephrol Dial Transplant* 2008; 23:2213–2222 doi: [10.1093/ndt/gfn093](https://doi.org/10.1093/ndt/gfn093) PMID: [18400820](https://pubmed.ncbi.nlm.nih.gov/18400820/)
24. Meier J, Kemming GI, Kisch-Wedel H, Blum J, Pape A, Habler OP. Hyperoxic ventilation reduces six-hour mortality after partial fluid resuscitation from hemorrhagic shock. *Shock* 2004; 22:240–247 PMID: [15316394](https://pubmed.ncbi.nlm.nih.gov/15316394/)
25. Blasiolo B, Bayir H, Vagni VA, Janesko-Feldman K, Cheikhi A, Wisniewski SR, et al. Effect of hyperoxia on resuscitation of experimental combined traumatic brain injury and hemorrhagic shock in mice. *Anesthesiology* 2013; 118:649–663 doi: [10.1097/ALN.0b013e318280a42d](https://doi.org/10.1097/ALN.0b013e318280a42d) PMID: [23299361](https://pubmed.ncbi.nlm.nih.gov/23299361/)

26. Waisman D, Brod V, Wolff R, Sanop E, Chernin M, Weintraub Z, et al. Effects of hyperoxia on local and remote microcirculatory inflammatory response after splanchnic ischemia and reperfusion. *Am J Physiol Heart Circ Physiol* 2003; 285:H643–H652 PMID: [12714329](#)
27. Sukhotnik I, Brod V, Lurie M, Rahat MA, Shnizer S, Lahat N, et al. The effect of 100% oxygen on intestinal preservation and recovery following ischemia-reperfusion injury in rats. *Crit Care Med* 2009; 37:1054–1061 doi: [10.1097/CCM.0b013e31819d0f5c](#) PMID: [19237917](#)
28. Buras JA, Holt D, Orlow D, Bellkoff B, Pavlides S, Reenstra WR. Hyperbaric oxygen protects from sepsis mortality via an interleukin-10-dependent mechanism. *Crit Care Med* 2006; 34:2624–2629 PMID: [16932233](#)
29. Barth E, Bassi G, Maybauer DM, Simon F, Gröger M, Öter S, et al. Effects of ventilation with 100% oxygen during early hyperdynamic porcine fecal peritonitis. *Crit Care Med* 2008; 36:495–503 PMID: [18091553](#)
30. Hauser B, Barth E, Bassi G, Simon F, Gröger M, Öter S, et al. Hemodynamic, metabolic and organ function effects of pure O₂ ventilation during established fecal peritonitis-induced septic shock. *Crit Care Med* 2009; 37:2465–2469 doi: [10.1097/CCM.0b013e3181aee8ad](#) PMID: [19531939](#)
31. Hou L, Xie K, Li N, Qin M, Lu Y, Ma S, et al. 100% oxygen inhalation protects against zymosan-induced sterile sepsis in mice: the roles of inflammatory cytokines and antioxidant enzymes. *Shock* 2009; 32:451–461 doi: [10.1097/SHK.0b013e31819c391a](#) PMID: [19174736](#)
32. Waisman D, Brod V, Rahat MA, Amit-Cohen BC, Lahat N, Rimar D, et al. Dose-related effects of hyperoxia on the lung inflammatory response in septic rats. *Shock* 2012; 37:95–102 doi: [10.1097/SHK.0b013e3182356fc3](#) PMID: [21921827](#)
33. Wagner F, Wagner K, Weber S, Stahl B, Knöferl MW, Huber-Lang M, et al. Inflammatory effects of hypothermia and inhaled H₂S during resuscitated, hyperdynamic murine septic shock. *Shock* 2011; 35:396–402 doi: [10.1097/SHK.0b013e3181ffff0e](#) PMID: [20938376](#)
34. Wagner K, Wachter U, Vogt JA, Scheuerle A, McCook O, Weber S, et al. Adrenomedullin binding improves catecholamine responsiveness and kidney function in resuscitated murine septic shock. *Intensive Care Med Exp* 2013; 1:2
35. Reiss LK, Kowallik A, Uhlig S. Recurrent recruitment manoeuvres improve lung mechanics and minimize lung injury during mechanical ventilation of healthy mice. *PLoS ONE* 2011; 6:e24527 doi: [10.1371/journal.pone.0024527](#) PMID: [21935418](#)
36. Matute-Bello G, Downey G, Moore BB, Groshong SD, Matthay MA, Slutsky AS, et al. An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. *Am J Respir Crit Care Med* 2011; 44:725–738
37. Geldmacher H, Biller H, Herbst A, Urbanski K, Allison M, Buist AS, et al. The prevalence of chronic obstructive pulmonary disease (COPD) in Germany. Results of the BOLD study. *Dtsch Med Wochenschr* 2008; 133:2609–2614 PMID: [19052996](#)
38. Papandrinopoulou D, Tzouda V, Tsoukalas G. Lung compliance and chronic obstructive pulmonary disease. *Pulm Med* 2012; 2012:542769 doi: [10.1155/2012/542769](#) PMID: [23150821](#)
39. O'Donnell DE, Lavenezia P. Physiology and consequences of lung hyperinflation in COPD. *Eur Respir Rev* 2006; 15:61–67
40. Geraghty P, Wyman AE, Garcia-Arcos I, Dabo AJ, Gadhvi S, Foronjy R. STAT3 modulates cigarette smoke-induced inflammation and protease expression. *Front Physiol* 2013; 4:267 doi: [10.3389/fphys.2013.00267](#) PMID: [24101903](#)
41. Guan SP, Tee W, Ng DSW, Chan TK, Peh HY, Ho WE, et al. Andrographolide protects against cigarette smoke-induced oxidative lung injury via augmentation of Nrf2 activity. *Br J Pharmacol* 2013; 168:1707–1718 doi: [10.1111/bph.12054](#) PMID: [23146110](#)
42. Park JW, Ryter SW, Kyung SY, Lee SP, Jeong SH. The phosphodiesterase 4 inhibitor rolipram protects against cigarette smoke extract-induced apoptosis in human lung fibroblasts. *Eur J Pharmacol* 2013; 706:76–83 doi: [10.1016/j.ejphar.2013.02.049](#) PMID: [23499692](#)
43. Dolinay T, Kim YS, Howrylak J, Hunninghake GM, An CH. Inflammasome-regulated cytokines are critical mediators of acute lung injury. *Am J Respir Crit Care Med* 2012; 185:1225–1234 doi: [10.1164/rccm.201201-0003OC](#) PMID: [22461369](#)
44. Kolliputi N, Shaik RS, Waxman AB. The inflammasome mediates hyperoxia-induced alveolar cell permeability. *J Immunol* 2010; 184: 5819–5826 doi: [10.4049/jimmunol.0902766](#) PMID: [20375306](#)
45. Monção-Ribeiro L, Cagido VR, Lima-Murad G, Teixeira-Santana P, Riva DR, Borojevic R, et al. Lipopolysaccharide-induced lung injury: Role of P2X7 receptor. *Respir Physiol Neurobiol* 2011; 179:314–325 PMID: [21982752](#)

46. Chiao CW, Tostes RC, Webb RC. P2X₇ receptor activation amplifies lipopolysaccharide-induced vascular hyporeactivity via interleukin-1 β release. *J Pharmacol Exp Ther* 2008; 326:864–870 doi: [10.1124/jpet.107.135350](https://doi.org/10.1124/jpet.107.135350) PMID: [18559654](https://pubmed.ncbi.nlm.nih.gov/18559654/)
47. Kawano A, Tsukimoto M, Mori D, Noguchi T, Harada H, Takenouchi T, et al. Regulation of P2X₇-dependent inflammatory functions by P2X₄ receptor in mouse macrophages. *Biochem Biophys Res Comm* 2012; 420:102–107 doi: [10.1016/j.bbrc.2012.02.122](https://doi.org/10.1016/j.bbrc.2012.02.122) PMID: [22405772](https://pubmed.ncbi.nlm.nih.gov/22405772/)
48. Weinhold K, Krause-Buchholz U, Rödel G, Kasper M, Barth K. Interaction and interrelation of P2X₇ and P2X₄ receptor complexes in mouse lung epithelial cells. *Cell Mol Life Sci* 2010; 67:2631–2642 doi: [10.1007/s00018-010-0355-1](https://doi.org/10.1007/s00018-010-0355-1) PMID: [20405163](https://pubmed.ncbi.nlm.nih.gov/20405163/)
49. Cicko S, Lucattelli M, Müller T, Lommatzsch M, De Cunto G, Cardini S, et al. Purinergic receptor inhibition prevents the development of smoke-induced lung injury and emphysema. *J Immunol* 2010; 185:688–697 doi: [10.4049/jimmunol.0904042](https://doi.org/10.4049/jimmunol.0904042) PMID: [20519655](https://pubmed.ncbi.nlm.nih.gov/20519655/)
50. Eltom S, Stevenson CS, Rastrick J, Dale N, Raemdonck K, Wong S, et al. P2X₇ receptor and caspase 1 activation are central to airway inflammation observed after exposure to tobacco smoke. *PLoS ONE* 2011; 6:e24097 doi: [10.1371/journal.pone.0024097](https://doi.org/10.1371/journal.pone.0024097) PMID: [21915284](https://pubmed.ncbi.nlm.nih.gov/21915284/)
51. Hafner S, Wagner K, Wepler M, Matallo J, Gröger M, McCook O, et al. Physiological and immune-biological characterization of a long-term murine model of blunt chest trauma. *Shock* 2015; 43:140–147 doi: [10.1097/SHK.0000000000000277](https://doi.org/10.1097/SHK.0000000000000277) PMID: [25526372](https://pubmed.ncbi.nlm.nih.gov/25526372/)
52. Lucattelli M, Cicko S, Müller T, Lommatzsch M, De Cunto G, Cardini S, et al. P2X₇ receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am J Respir Cell Mol Biol* 2011; 44:423–429 doi: [10.1165/rcmb.2010-0038OC](https://doi.org/10.1165/rcmb.2010-0038OC) PMID: [20508069](https://pubmed.ncbi.nlm.nih.gov/20508069/)
53. Müller T, Viera RP, Grimm M, Dürk T, Cicko S, Zeiser R, et al. A potential role for P2X₇R in allergic airway inflammation in mice and humans. *Am J Respir Cell Mol Biol* 2011; 44:456–464 doi: [10.1165/rcmb.2010-0129OC](https://doi.org/10.1165/rcmb.2010-0129OC) PMID: [20508067](https://pubmed.ncbi.nlm.nih.gov/20508067/)
54. Jiang H, Zhu Y, Xu H, Sun Y, Li Q. Activation of hypoxia-inducible factor-1 α via nuclear factor- κ B in rats with chronic obstructive pulmonary disease. *Acta Biochem Biophys Sin* 2010; 42:483–488
55. Yasuo M, Mizuno S, Kraskauskas D, Bogaard HJ, Natarajan R, Cool CD, et al. Hypoxia inducible factor-1 α in human emphysema lung tissue. *Eur Respir J* 2011; 37:775–783 doi: [10.1183/09031936.00022910](https://doi.org/10.1183/09031936.00022910) PMID: [20562128](https://pubmed.ncbi.nlm.nih.gov/20562128/)
56. Lee SH, Lee SH, Kim CH, Yang KS, Lee EJ, Min KH, et al. Increased expression of vascular endothelial growth factor and hypoxia inducible factor-1 α in lung tissue of patients with chronic bronchitis. *Clin Biochem* 2014; 47:552–559 PMID: [24463065](https://pubmed.ncbi.nlm.nih.gov/24463065/)
57. Metzen E, Zhou J, Jelkmann W, Fandrey J, Brüne B. Nitric oxide impairs normoxic degradation of HIF-1 α by inhibition of prolyl hydroxylases. *Mol Biol Cell* 2003; 14:3470–3481 PMID: [12925778](https://pubmed.ncbi.nlm.nih.gov/12925778/)
58. Berchner-Pfannenschmidt U, Tug S, Kirsch M, Fandrey J. Oxygen-sensing under the influence of nitric oxide. *Cell Signal* 2010; 22:349–356 doi: [10.1016/j.cellsig.2009.10.004](https://doi.org/10.1016/j.cellsig.2009.10.004) PMID: [19861159](https://pubmed.ncbi.nlm.nih.gov/19861159/)
59. Dantzker DR, Wagner PD, West JB. Instability of lung units with low V_A/Q ratios during O₂ breathing. *J Appl Physiol* 1975; 38:886–895
60. Quinn DA, Moufarrej RK, Volokhov A, Hales CA. Interactions of lung stretch, hyperoxia, and MIP-2 production in ventilator-induced lung injury. *J Appl Physiol* 2002; 930:517–525
61. Sinclair SE, Altemeier WA, Matute-Bello G, Chi EY. Augmented lung injury due to interaction between hyperoxia and mechanical ventilation. *Crit Care Med* 2004; 32:2496–2501 PMID: [15599157](https://pubmed.ncbi.nlm.nih.gov/15599157/)
62. Cannizzaro V, Berry LJ, Zosky GR, Turner DJ, Hantos Z, Sly PD. Impact of supplemental oxygen in mechanically ventilated adult and infant mice. *Respir Physiol Neurobiol* 2009; 165:61–65 doi: [10.1016/j.resp.2008.10.006](https://doi.org/10.1016/j.resp.2008.10.006) PMID: [18992373](https://pubmed.ncbi.nlm.nih.gov/18992373/)
63. Müller HC, Witzernath M, Tscherning T, Gutbier B, Hippenstiel S, Santel A, et al. Adrenomedullin attenuates ventilator-induced lung injury in mice. *Thorax* 2010; 65:1077–1084 doi: [10.1136/thx.2010.135996](https://doi.org/10.1136/thx.2010.135996) PMID: [20971983](https://pubmed.ncbi.nlm.nih.gov/20971983/)
64. Madjdpour C, Jewell UR, Kneller S, Ziegler U, Schwendener R, Booy C, et al. Decreased alveolar oxygen induces lung inflammation. *Am J Physiol Lung Cell Mol Physiol* 2002; 284:L360–L367. PMID: [12388372](https://pubmed.ncbi.nlm.nih.gov/12388372/)
65. Vuichard D, Ganter MT, Schimmer RC, Suter D, Booy C, Reyes L, et al. Hypoxia aggravates lipopolysaccharide-induced lung injury. *Clin Exp Immunol* 2005; 141:248–60 PMID: [15996189](https://pubmed.ncbi.nlm.nih.gov/15996189/)
66. Chao J, Donham P, van Rooijen N, Wood JG, Gonzalez NC. Monocyte chemoattractant protein-1 released from alveolar macrophages mediates the systemic inflammation of acute alveolar hypoxia. *Am J Respir Cell Mol Biol* 2011; 45:53–61 doi: [10.1165/rcmb.2010-0264OC](https://doi.org/10.1165/rcmb.2010-0264OC) PMID: [20813992](https://pubmed.ncbi.nlm.nih.gov/20813992/)
67. Fišárková B, Vytášek R, Miková D, Vízek M. Hyperoxia attenuated nitrotyrosine concentration in the lung tissue of rats with experimental pneumonia. *Physiol Res* 2004; 53:487–492 PMID: [15479126](https://pubmed.ncbi.nlm.nih.gov/15479126/)

68. Ohata Y, Ogata S, Nakanishi K, Kanazawa F, Uenoyama M, Tominaga S, et al. Expression of P2X4R mRNA and protein in rats with hypobaric hypoxia-induced pulmonary hypertension. *Circ J* 2011; 75:945–954 PMID: [21378451](#)
69. Feinman R, Deitch EA, Watkins AC, Abungu B, Colorado I, Kannan KB, et al. HIF-1 mediates pathogenic inflammatory responses to intestinal ischemia-reperfusion injury. *Am J Physiol Gastrointest Liver Physiol* 2010; 299:G833–G843 doi: [10.1152/ajpgi.00065.2010](#) PMID: [20689059](#)
70. Suresh MV, Ramakrishnan SK, Thomas B, Machado-Arnada D, Bi Y, Talarico N, et al. Activation of hypoxia-inducible factor 1- α in type 2 alveolar epithelial cell is a major driver of acute inflammation following lung contusion. *Crit Care Med* 2014; 42:e642–e653 doi: [10.1097/CCM.0000000000000488](#) PMID: [25014067](#)
71. Jiang H, Huang Y, Xu H, Li QF. Inhibition of hypoxia inducible factor-1 α ameliorates lung injury induced by trauma and hemorrhagic shock in rats. *Acta Pharmacol Sin* 2012; 33:635–643 doi: [10.1038/aps.2012.5](#) PMID: [22465950](#)
72. Calvert JW, Cahill J, Yamaguchi-Okada M, Zhang JH. Oxygen treatment after experimental hypoxia-ischemia in neonatal rats alters the expression of HIF-1 α and its downstream target genes. *J Appl Physiol* 2006; 101:853–865 PMID: [16728520](#)
73. Uematsu S, Engelberts D, Peltekova V, Otulakowski G, Post M, Kavanagh BP. Dissociation of inflammatory mediators and function: experimental lung injury in nonpulmonary sepsis. *Crit Care Med* 2013; 41:151–158 doi: [10.1097/CCM.0b013e318267606f](#) PMID: [23128385](#)
74. Albeceita G, Fernandez A, Parra D, Gonzalo JA, García-Prieto E, Taboada F. Mechanical ventilation causes monocyte deactivation in intact and endotoxin-treated mice. *J Trauma* 2008; 64:470–476 doi: [10.1097/TA.0b013e31814931ac](#) PMID: [18301217](#)
75. Wilson MR, Patel BV, Takata M. Ventilation with 'clinically relevant' high tidal volumes does not promote stretch-induced lung injury in the lungs of healthy mice. *Crit Care Med* 2012; 40:2850–2857 PMID: [22890257](#)
76. Wolthuis EK, Vlaar APJ, Choi G, Roeleofs JJTH, Juffermans NP, Schultz MJ. Mechanical ventilation using non-injurious ventilation settings causes lung injury in the absence of pre-existing lung injury in healthy mice. *Crit Care* 2009; 13:R1 doi: [10.1186/cc7688](#) PMID: [19152704](#)
77. Vlaar APJ, Wolthuis EK, Hofstra JJ, Roeleofs JJTH, Boon L, Schultz MJ, et al. Mechanical ventilation aggravates transfusion-related acute lung injury induced by MHC-1 class antibodies. *Intensive Care Med* 2010; 36:879–887 doi: [10.1007/s00134-010-1802-z](#) PMID: [20221752](#)
78. Müller-Redetzky HC, Will D, Hellwig K, Kummer W, Tschernig T, Pfeil U, et al. Mechanical ventilation drives pneumococcal pneumonia into lung injury and sepsis in mice: protection by adrenomedullin. *Crit Care* 2014; 18:R73 doi: [10.1186/cc13830](#) PMID: [24731244](#)
79. Rossi JL, Velentza AV, Steinhorn DM, Watterson DM, Wainwright MS. MLCK210 gene knockout or kinase inhibition preserves lung function following endotoxin-induced lung injury in mice. *Am J Physiol Lung Cell Mol Physiol* 2007; 292:L1327–L1334 PMID: [17307813](#)
80. Reiss LK, Kowallik A, Uhlig S. Recurrent recruitment manoeuvres improve lung mechanics and minimize lung injury during mechanical ventilation of healthy mice. *PLoS ONE* 2011; 6:e24527 doi: [10.1371/journal.pone.0024527](#) PMID: [21935418](#)