

Profiling meteorite soluble organic matter using (ultra)high resolution organic spectroscopy

Philippe Schmitt-Kopplin (1,2), Alexander Ruf (3), Norbert Hertkorn (1), Mourad Harir (1), Basem Kanawati (1), Marianna Lucio (1), Eric Quirico (4), Ansgar Greshake (6), Zelimir Gabelica (5)

(1) Analytical BioGeoChemistry, Helmholtz Zentrum Muenchen, Munich, Germany, (2) Technische Universität München, Munich, Germany (3) Université de Marseille, France (4) Université Grenoble Alpes, Institut de Planétologie de Grenoble (IPAG), Grenoble, France (5) Museum für Naturkunde, Berlin, Germany (6) Université de Haute Alsace, Mulhouse, France. (schmitt-kopplin@helmholtz-muenchen.de)

Abstract

We present here a review of the state of the art in non-targeted high resolution analysis of the soluble organic fraction of various meteorites; these include carbonaceous chondrites as well as achondrites and ordinary chondrites.

1. Introduction

Understanding the origin and evolution of organic matter, is linked to observationally-derived astrochemistry (telescopic observations) and the laboratory wet chemical analysis of return objects and meteorites. The molecular composition and diversity of non-terrestrial organic matter in carbonaceous chondrites was studied by means of both, targeted [1] and non-targeted [2,3] chemical analytical approaches, leading to new insights. Targeted chemical analyses are hypothesis-driven and are largely focused on molecules of biological/prebiotic interest. In a non-targeted approach, all analytes are globally profiled within the analytical possibilities without biased or constrained hypothesis in order to gain comprehensive information. We review in this presentation the state of the art in using non-targeted high resolution organic spectroscopy.

2. The analytical approaches

Ultrahigh-resolving analytics, like high field Fourier transform ion cyclotron resonance mass spectrometry (FT-ICR-MS) and nuclear magnetic resonance spectroscopy (NMR), represent a powerful tool to allow insights into the holistic complex compositional space to tens of thousands of different molecular compositions and functional groups and likely millions of diverse structures. This could be observed in solvent extracts of pristine carbonaceous

meteorites [2, 3], and suggests that interstellar chemistry is extremely active and rich. Since then we studied the chemical composition of thousands of individual components out of complex organic mixtures, as accessed in the solvent-accessible organic fraction, extracted under mild conditions, from diversely-classified and heated meteorites

3. Results

We described that heteroatomic organic molecules play an important role in the description of non-terrestrial chemical evolution. The thermally and shock-stressed Chelyabinsk (LL5) [4] showed high number of nitrogen counts within CHNO molecular formulas, especially in the melt region. This match of the organic molecular profile with the petrologic character could be also observed for Novato (L6) [5], Braunschweig (L6) and the latest German fall Stubenberg (LL6) [6]. Additionally, the extremely thermally altered Sutter's mill (C-type) [7] reflects a loss in the organic diversity, but an increase in the polysulphur domain, as compared to other CM2-analyzed falls. The increase of polynitrogen and polysulphur compounds could be simulated in laboratory experiments by heating Murchison (CM2). Recently we reported the discovery of a previously unrecognized chemical class, dihydroxymagnesium carboxylates, $[(HO)_2MgO_2CR]^-$, gained from nonterrestrial meteoritic analyses [8]. These thermostable compounds might have contributed to the stabilization of organic molecules on a geological time scale, which emphasizes their potential astrobiological relevance.

The resulted extreme richness in chemical diversity offers information on the meteoritic parent body history and help in expanding our knowledge or astrochemistry towards higher molecular masses and complex molecular structures.

References

- [1] Pizzarello, S. et al. PNAS, 110(39):15614–15619.
- [2] Schmitt-Kopplin, P. et al. 2010 PNAS 107(7):2763–2768.
- [3] Hertkorn, N. et al. 2015. Magnetic Resonance in Chemistry, 53(9):754–768.
- [4] Popova, O.P. et al. 2013 Science, 342(6162): 1069–1073.
- [5] Jenniskens, P. et al. 2014. Meteoritics & Planetary Science, 49(8): 1388–1425.
- [6] Bischoff, A. 2017 Meteoritics & Planetary Science, 52(8):1683–1703.
- [7] Jenniskens P. et al. Science 2012, 338: 1583-1587.
- [8] Ruf A. et al. 2017, PNAS, 114(11): 2819-2824.