



Flavor violating muon decay into an electron and a light gauge boson

Alejandro Ibarra^a, Marcela Marín^b, Pablo Roig^{b,*}

^a Physik-Department, Technische Universität München, James-Frank-Straße, 85748 Garching, Germany

^b Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apartado Postal 14-740, 07000, Ciudad de México, Mexico

ARTICLE INFO

Article history:

Received 2 November 2021

Received in revised form 19 January 2022

Accepted 25 January 2022

Available online 29 January 2022

Editor: J. Hisano

ABSTRACT

We analyze the flavor violating muon decay $\mu \rightarrow e\chi$, where χ is a massive gauge boson, with emphasis in the regime where χ is ultralight. We first study this process from an effective field theory standpoint in terms of form factors. We then present two explicit models where $\mu \rightarrow e\chi$ is generated at tree level and at the one-loop level. We also comment on the prospects of observing the process $\mu \rightarrow e\chi$ in view of the current limits on $\mu \rightarrow 3e$ from the SINDRUM collaboration.

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The Standard Model of Particle Physics [1–3] predicts the conservation of lepton flavor. The discovery of neutrino oscillations [4–6] provided conclusive evidence for the violation of lepton flavor in Nature, and therefore for the existence of new physics beyond the Standard Model. However, and aside for neutrino oscillations, no other lepton flavor violating process has been observed up to this day.

Experimental searches for lepton flavor violation in muon decays date back to the late 1940's [7,8]. The most sensitive searches as of today have set the upper limits $\text{Br}(\mu^+ \rightarrow e^+\gamma) \leq 4.2 \times 10^{-13}$ by the MEG collaboration [9] and $\text{Br}(\mu^+ \rightarrow e^+e^-e^+) \leq 1.0 \times 10^{-12}$ by the SINDRUM collaboration [10]. The searches for rare muon decays are complemented by those for other lepton flavor violating processes, such as $\mu - e$ conversion in nuclei [11–13], muonium-antimuonium conversion [14], or in neutral meson decays, such as $K_L^0 \rightarrow \mu^\pm e^\mp$ [15] or $B^0 \rightarrow \mu^\pm e^\mp$ [16] among others (see e.g. [17]).

In recent years there has been interest in Physics at the low energy frontier (for a review, see [18]). This possibility may lead to new lepton flavor violating muon decays, such as $\mu \rightarrow e\chi$, with χ an invisible boson. The non-observation of this decay by the TWIST collaboration allows to set the 90% C.L. upper limit $\text{BR}(\mu \rightarrow e\chi) < 8.1 \times 10^{-6}$ [19]. Further, the light boson χ could lead to the three-body lepton flavor violating decay $\mu^+ \rightarrow e^+e^-e^+$, when χ is off-shell, resulting into complementary constraints on this scenario.

In this work we will focus on the possibility that χ is a light gauge boson, associated to the spontaneous breaking of an Abelian gauge symmetry, $U(1)_\chi$ (the case where χ is a light scalar or a pseudoscalar has been extensively studied, see e.g. [20–35]).

The simplest Lagrangian describing the lepton flavor interacting interaction is $\mathcal{L}_{\text{LFV}} = g\bar{\mu}\gamma^\rho\chi_\rho e + \text{h.c.}$, with χ_ρ the 4-potential associated to the $U(1)_\chi$ symmetry. With this effective description, the rate for $\mu \rightarrow e\chi$ contains terms proportional to g^2/m_χ^2 , due to the emission of the longitudinal component of the gauge boson (a similar behavior is found for other effective interactions). Naively, the rate diverges as $m_\chi \rightarrow 0$, therefore the effective theory cannot be matched to the well studied decay $\mu \rightarrow e\gamma$. Further, one may wonder whether the decays into several gauge bosons $\mu \rightarrow e\chi \cdots \chi$, could also contribute significantly to the total decay width, reminiscent of the “hyperphoton catastrophe” for the electron decay into a neutrino and an ultralight photon [36–38].

In this paper we present a detailed analysis of the decay rate $\mu \rightarrow e\chi$, with special emphasis in the regime where $m_\chi \rightarrow 0$ (for previous works, see e.g. [39–41]). In Section 2 we present the most general effective interaction leading to the decay $\mu \rightarrow e\chi$ in terms of a number of form factors, and we identify the conditions that the form factors must fulfill in order to render a finite rate in the limit $m_\chi \rightarrow 0$. In Sections 3 and 4 we present two gauge invariant and renormalizable models where the process $\mu \rightarrow e\chi$ is generated, respectively, at tree level and at the one-loop level. We calculate the rate for $\mu \rightarrow e\chi$ and we explicitly show that the rate remains finite as $m_\chi \rightarrow 0$. For these two models, we also calculate the rate for the rare decay $\mu \rightarrow 3e$. Finally, in Section 5 we present our conclusions.

* Corresponding author.

E-mail address: proig@fis.cinvestav.mx (P. Roig).

Table 1

Spins and charges under $SU(2)_L \times U(1)_Y \times U(1)_\chi$ of the particles of the model described in Section 3, leading to the decay $\mu \rightarrow e\chi$ at tree level. All fields are assumed to be singlets under $SU(3)_C$.

	L_1	L_2	e_{R_1}	e_{R_2}	ϕ_{11}	ϕ_{12}	ϕ_{21}	ϕ_{22}
spin	1/2	1/2	1/2	1/2	0	0	0	0
$SU(2)_L$	2	2	1	1	2	2	2	2
$U(1)_Y$	-1/2	-1/2	-1	-1	Y_{11}	Y_{12}	Y_{21}	Y_{22}
$U(1)_\chi$	q_{L_1}	q_{L_2}	q_{e_1}	q_{e_2}	$q_{\phi_{11}}$	$q_{\phi_{12}}$	$q_{\phi_{21}}$	$q_{\phi_{22}}$

2. Effective theory

We consider first the effective theory describing the decay $\mu \rightarrow e\chi$, where χ is a light gauge boson, $m_\chi \ll m_\mu$. We denote the four-momenta of the muon, electron and χ as p_μ , p_e and p_χ , respectively. The transition amplitude is given by $M = \bar{u}(p_e)\Gamma^\alpha(p_\mu, p_e)u(p_\mu)\epsilon_\alpha^*(p_\chi)$, where $\Gamma^\alpha(p_\mu, p_e)$ can be written in terms of six dimensionless scalar form factors $F_i(p_\chi^2)$, $G_i(p_\chi^2)$, $i = 1, 2, 3$, as:

$$\Gamma^\alpha = \left(\gamma^\alpha - \frac{\not{p}_\chi \not{p}_\chi^\alpha}{p_\chi^2} \right) F_1(p_\chi^2) + i \frac{\sigma^{\alpha\beta} p_{\chi\beta}}{m_\mu + m_e} F_2(p_\chi^2) + \frac{2p_\chi^\alpha}{m_\mu + m_e} F_3(p_\chi^2) + \left(\gamma^\alpha - \frac{\not{p}_\chi \not{p}_\chi^\alpha}{p_\chi^2} \right) \gamma^5 G_1(p_\chi^2) + i \frac{\sigma^{\alpha\beta} \gamma^5 p_{\chi\beta}}{m_\mu + m_e} G_2(p_\chi^2) + \frac{2p_\chi^\alpha}{m_\mu + m_e} \gamma^5 G_3(p_\chi^2), \quad (1)$$

where $\sigma^{\alpha\beta} = -\frac{i}{2}[\gamma^\alpha, \gamma^\beta]$. For the decay process $\mu \rightarrow e\chi$, where the gauge boson χ is on-shell, $p_\chi^2 = m_\chi^2$. The conservation of the $U(1)_\chi$ charge requires the form factor $F_3(p_\chi^2)$ to vanish. Moreover the Ward identities imply that $p_\chi^\alpha \cdot \epsilon_\alpha^*(p_\chi) = 0$, so the terms proportional to $G_3(p_\chi^2)$ and $\frac{p_\chi p_\chi^\alpha}{p_\chi^2}$ will not contribute to the amplitude.

The decay rate can then be expressed in terms of four form factors and reads:

$$\Gamma(\mu \rightarrow e\chi) = \frac{\lambda^{1/2}[m_\mu^2, m_e^2, m_\chi^2]}{16\pi m_\mu} \left[\left(1 - \frac{m_e}{m_\mu} \right)^2 \left(1 - \frac{m_\chi^2}{(m_\mu - m_e)^2} \right) \left(2|F_1(m_\chi^2) - F_2(m_\chi^2)|^2 + \left| F_1(m_\chi^2) \frac{(m_\mu + m_e)}{m_\chi} - F_2(m_\chi^2) \frac{m_\chi}{(m_\mu + m_e)} \right|^2 \right) + \left(1 + \frac{m_e}{m_\mu} \right)^2 \left(1 - \frac{m_\chi^2}{(m_\mu + m_e)^2} \right) \left(2|G_1(m_\chi^2) - G_2(m_\chi^2) \frac{(m_\mu - m_e)}{(m_\mu + m_e)}|^2 + \left| G_1(m_\chi^2) \frac{(m_\mu - m_e)}{m_\chi} + G_2(m_\chi^2) \frac{m_\chi}{(m_\mu + m_e)} \right|^2 \right) \right], \quad (2)$$

where $\lambda[m_\mu^2, m_e^2, m_\chi^2]$ is the usual Källén function. Analogous expressions hold for the decays $\tau \rightarrow \mu\chi$ and $\tau \rightarrow e\chi$, with the appropriate substitutions.

The term proportional to $1/m_\chi$ corresponds to the emission of the longitudinal component of the vector boson, and apparently makes the limit $m_\chi \rightarrow 0$ divergent and not continuously matched to the well studied result from $\mu \rightarrow e\gamma$ [42–44]. Therefore, in an effective field theory approach, great care should be taken when considering decays into ultralight gauge bosons, since in a gauge invariant and renormalizable theory one generically expects the rate of $\mu \rightarrow e\chi$ to be finite.

In the following sections we present two explicit models where the lepton flavor violating effective interaction Eq. (1) is generated either at tree-level or at the one-loop level. Apart from the interest of the models in themselves, we will show explicitly that the form factors $F_1(m_\chi^2)$ and $G_1(m_\chi^2)$ contain an implicit dependence on m_χ , rooted in gauge invariance, so that the rate for $\mu \rightarrow e\chi$ is finite in the limit $m_\chi \rightarrow 0$.

3. $\mu \rightarrow e\chi$ at tree level

The particle content of the model, and the corresponding spins and charges under $SU(2)_L \times U(1)_Y \times U(1)_\chi$, are summarized in Table 1.¹ Here, $L_i = (\nu_{L_i}, e_{L_i})$ and e_{R_i} , $i = 1, 2$, denote the Standard Model $SU(2)_L$ lepton doublets and singlets, respectively (we have restricted ourselves to the two generation case, although the extension to three generations is straightforward). They have a generation independent hypercharge, Y_L and Y_e respectively, although we allow for generation dependent charges under $U(1)_\chi$. Further, ϕ_{jk} , $i, j = 1, 2$ denote complex scalar fields, doublets under $SU(2)_L$, with hypercharge Y_{jk} and charge under $U(1)_\chi$ equal to $q_{\phi_{jk}}$.

The kinetic terms of the particles of the model read:

$$\mathcal{L}_{\text{kin}} = \sum_{j=1}^2 i(\bar{L}_j \not{D} L_j + \bar{e}_{R_j} \not{D} e_{R_j}) + \sum_{j,k=1}^2 (D_\mu \phi_{jk})^\dagger (D^\mu \phi_{jk}), \quad (3)$$

where D_μ denotes the covariant derivative, given by

¹ The model can be made anomaly-free adding heavy particles with suitable charges, without modifying the discussion that follows.

$D_\mu = \partial_\mu + igW_\mu^a T_a + ig'YB_\mu + ig_\chi q\chi_\mu$ for the $SU(2)_L$ doublets ,

$D_\mu = \partial_\mu + ig'YB_\mu + ig_\chi q\chi_\mu$ for the $SU(2)_L$ singlets , (4)

with g , g' and g_χ the coupling constants of $SU(2)_L$, $U(1)_Y$ and $U(1)_\chi$ respectively.

We also assume $Y_{jk} = 1/2$. Then, for j, k such that $q_{\phi_{jk}} = q_{L_j} - q_{e_k}$ the following Yukawa couplings arise in the Lagrangian:

$$-\mathcal{L}_{\text{Yuk}} = \sum_{j,k=1}^2 y_{jk} \bar{L}_j \phi_{jk} e_{R_k} + \text{h.c.} \quad (5)$$

We also assume that the doublet scalars acquire a vacuum expectation value for some i, j , $\langle \phi_{jk} \rangle = v_{jk}$. To keep the discussion general, we consider that the charges of the particles allow all Yukawa couplings, and that all ϕ_{jk} acquire a vacuum expectation value; the different subcases follow straightforwardly by setting the corresponding y_{jk} and/or v_{jk} to zero.

The non-zero expectation values for ϕ_{jk} generate a mass for the χ boson:

$$m_\chi^2 = g_\chi^2 (q_{\phi_{11}}^2 v_{11}^2 + q_{\phi_{12}}^2 v_{12}^2 + q_{\phi_{21}}^2 v_{21}^2 + q_{\phi_{22}}^2 v_{22}^2) . \quad (6)$$

Furthermore, since ϕ_{jk} have charge under $SU(2)_L \times U(1)_Y$, their expectation value would also contribute to the Z and W masses. Since we are assuming $m_\chi \ll m_\mu$, this contribution can be safely neglected.

The expectation value of the doublet scalars generates a mass term for the charged leptons, $-\mathcal{L}_{\text{mass}} \supset \bar{e}_L M_{jk} e_{R_k} + \text{h.c.}$, with

$$M = \begin{pmatrix} y_{11} v_{11} & y_{12} v_{12} \\ y_{21} v_{21} & y_{22} v_{22} \end{pmatrix} . \quad (7)$$

We now rotate the fields to express the Lagrangian in the mass eigenstate basis:

$$\begin{pmatrix} e_L \\ \mu_L \end{pmatrix} = \begin{pmatrix} \cos \theta_L & \sin \theta_L \\ -\sin \theta_L & \cos \theta_L \end{pmatrix} \begin{pmatrix} e_{L1} \\ e_{L2} \end{pmatrix} , \quad \begin{pmatrix} e_R \\ \mu_R \end{pmatrix} = \begin{pmatrix} \cos \theta_R & \sin \theta_R \\ -\sin \theta_R & \cos \theta_R \end{pmatrix} \begin{pmatrix} e_{R1} \\ e_{R2} \end{pmatrix} \quad (8)$$

so that $-\mathcal{L}_{\text{mass}} \supset \bar{e}_L m_e e_R + \bar{\mu}_L m_\mu \mu_R + \text{h.c.}$, with

$$\begin{aligned} m_\mu^2 &\simeq y_{11}^2 v_{11}^2 + y_{12}^2 v_{12}^2 + y_{21}^2 v_{21}^2 + y_{22}^2 v_{22}^2 , \\ m_e^2 &\simeq \frac{(y_{11} v_{11} y_{22} v_{22} - y_{12} v_{12} y_{21} v_{21})^2}{y_{11}^2 v_{11}^2 + y_{12}^2 v_{12}^2 + y_{21}^2 v_{21}^2 + y_{22}^2 v_{22}^2} , \\ \sin 2\theta_L &\simeq -2 \frac{y_{11} v_{11} y_{21} v_{21} + y_{12} v_{12} y_{22} v_{22}}{y_{11}^2 v_{11}^2 + y_{12}^2 v_{12}^2 + y_{21}^2 v_{21}^2 + y_{22}^2 v_{22}^2} , \\ \sin 2\theta_R &\simeq -2 \frac{y_{11} v_{11} y_{12} v_{12} + y_{21} v_{21} y_{22} v_{22}}{y_{11}^2 v_{11}^2 + y_{12}^2 v_{12}^2 + y_{21}^2 v_{21}^2 + y_{22}^2 v_{22}^2} , \end{aligned} \quad (9)$$

where we have used that empirically $m_\mu \gg m_e$.

Finally, we recast the kinetic Lagrangian Eq. (4) in terms of the mass eigenstates. We find flavor violating terms of the form

$$-\mathcal{L} \supset \bar{e}_R i g_{e\mu}^{RR} \gamma^\rho \chi_\rho \mu_R + \bar{e}_L i g_{e\mu}^{LL} \gamma^\rho \chi_\rho \mu_L + \text{h.c.} , \quad (10)$$

with

$$\begin{aligned} g_{e\mu}^{RR} &= g_\chi (q_{e1} - q_{e2}) \sin \theta_R \cos \theta_R , \\ g_{e\mu}^{LL} &= g_\chi (q_{L1} - q_{L2}) \sin \theta_L \cos \theta_L . \end{aligned} \quad (11)$$

Clearly, if the $U(1)_\chi$ charges are generation independent, the flavor violation is absent at tree-level (as is the case for the photon and Z flavor violating couplings). Further, if the interaction eigenstates are aligned to the mass eigenstates, the tree-level flavor violation is also absent. This happens in particular for some choices of the expectation values of the fields ϕ_{ij} , for example, when $v_{ii} = 0$, but $v_{12}, v_{21} \neq 0$.

Comparing to the general form of the lepton flavor violating interaction vertex, Eq. (1), one can identify

$$\begin{aligned} F_1 &= \frac{1}{2} (g_{e\mu}^{RR} + g_{e\mu}^{LL}) , \\ G_1 &= \frac{1}{2} (g_{e\mu}^{RR} - g_{e\mu}^{LL}) , \end{aligned} \quad (12)$$

while all other form factors vanish at tree-level. The rate for $\mu \rightarrow e\chi$ then reads:

$$\Gamma(\mu \rightarrow e\chi) = \frac{m_\mu}{32\pi} (|g_{e\mu}^{LL}|^2 + |g_{e\mu}^{RR}|^2) \left(2 + \frac{m_\mu^2}{m_\chi^2}\right) \left(1 - \frac{m_\chi^2}{m_\mu^2}\right)^2 , \quad (13)$$

where we have neglected the electron mass against the muon mass. Naively, the term m_μ^2/m_χ^2 would enhance the rate as $m_\chi \rightarrow 0$. However, if the gauge and fermion masses arise as a consequence of the spontaneous breaking of the $U(1)_\chi$ symmetry, the limit $m_\chi \rightarrow 0$ requires $v_{jk} \rightarrow 0$ for all i, j (which in turn implies $m_\mu \rightarrow 0$), or $g_\chi \rightarrow 0$ (which in turn implies $g_{e\mu}^{LL}, g_{e\mu}^{RR} \rightarrow 0$). One can explicitly check

from Eqs. (6), (9) and (11) that indeed when $m_\chi \rightarrow 0$ the term $(|g_{e\mu}^{LL}|^2 + |g_{e\mu}^{RR}|^2)m_\mu^2/m_\chi^2$ is finite (as expected from the Goldstone boson equivalence theorem [45–48]), and depends on a function of the Yukawa couplings, the gauge coupling, and the charges and vacuum expectation values of the fields ϕ_{jk} .²

For example, for the specific case where the Yukawa couplings satisfy $y_{22} \gg y_{11} \gg y_{12}, y_{21}$ and all $v_{jk} = v$, so that the mass matrix M is almost diagonal, and $q_{jk} = Q$, the relevant parameters of the model after the spontaneous breaking of the gauge symmetries are:

$$\begin{aligned} m_\mu^2 &\simeq y_{22}^2 v^2, \quad m_e^2 \simeq y_{11}^2 v^2, \quad m_\chi^2 \simeq 4g_\chi^2 Q^2 v^2, \\ \sin 2\theta_L &\simeq -2\frac{y_{12}}{y_{22}}, \quad \sin 2\theta_R \simeq -2\frac{y_{21}}{y_{22}}. \end{aligned} \quad (14)$$

Therefore, the rate for $\mu \rightarrow e\chi$ in the limit $m_\chi \rightarrow 0$ is given by

$$\Gamma(\mu \rightarrow e\chi) \Big|_{m_\chi \rightarrow 0} \simeq \frac{m_\mu}{32\pi} \frac{g_\chi^2}{y_{22}^2} \left(2 + \frac{y_{22}^2}{4g_\chi^2 Q^2}\right) \left(1 - \frac{4g_\chi^2 Q^2}{y_{22}^2}\right)^2 \left[y_{12}^2 (q_{L1} - q_{L2})^2 + y_{21}^2 (q_{e1} - q_{e2})^2\right]. \quad (15)$$

The rate is maximal when $Qg_\chi/y_{22} \rightarrow 0$, and zero when $Qg_\chi/y_{22} \simeq 1/2$, which corresponds to $m_\chi/m_\mu \simeq 1$, i.e. when the phase space available for the decay closes. For most values of g_χ/y_{22} , the prefactor is $\sim 10^{-3}$, and therefore the rate can only be suppressed by invoking small couplings y_{12}, y_{21} , or by postulating intergenerational universality of the $U(1)_\chi$ charges. In the latter case, the process $\mu \rightarrow e\chi$ could be generated at the one loop level, as we will discuss in the next section.

A complementary probe of the μ - e flavor violation is the three-body decay $\mu^+ \rightarrow e^+ e^- e^+$, which is generated in this model at tree-level via the exchange of a virtual χ . The flavor conserving interaction vertex of the electron with the χ -boson has a similar form as Eq. (10):

$$-\mathcal{L} \supset \bar{e}_R i g_{ee}^{RR} \gamma^\rho \chi_\rho e_R + \bar{e}_L i g_{ee}^{LL} \gamma^\rho \chi_\rho e_L, \quad (16)$$

where

$$\begin{aligned} g_{ee}^{RR} &= g_\chi (q_{e2} \sin^2 \theta_R + q_{e1} \cos^2 \theta_R), \\ g_{ee}^{LL} &= g_\chi (q_{L2} \sin^2 \theta_L + q_{L1} \cos^2 \theta_L). \end{aligned} \quad (17)$$

The doubly differential decay width for $\mu \rightarrow 3e$ can be calculated from the interaction Lagrangians Eqs. (16) and (10) and reads:

$$\begin{aligned} \frac{d^2\Gamma(\mu \rightarrow 3e)}{ds dt} &= \frac{1}{128\pi^3 m_\mu^3} \left[(|g_{ee}^{LL}|^2 |g_{e\mu}^{RR}|^2 + |g_{ee}^{RR}|^2 |g_{e\mu}^{LL}|^2) \left(\frac{t(m_\mu^2 - t)}{(m_\chi^2 - s)^2 + m_\chi^2 \Gamma_\chi^2} + \frac{s(m_\mu^2 - s)}{(m_\chi^2 - t)^2 + m_\chi^2 \Gamma_\chi^2} \right) \right. \\ &\quad \left. + (|g_{ee}^{LL}|^2 |g_{e\mu}^{LL}|^2 + |g_{ee}^{RR}|^2 |g_{e\mu}^{RR}|^2) \left(\frac{(s+t)(m_\mu^2 - s - t)(4\Gamma_\chi^2 m_\chi^2 + (s+t-2m_\chi^2)^2)}{((m_\chi^2 - s)^2 + m_\chi^2 \Gamma_\chi^2)((m_\chi^2 - t)^2 + m_\chi^2 \Gamma_\chi^2)} \right) \right], \end{aligned} \quad (18)$$

where we have defined $s \equiv (p_\mu - p_{e1})^2$ and $t \equiv (p_\mu - p_{e2})^2$, with p_{e1} and p_{e2} the electron momenta, and which are kinematically restricted to be in the range:

$$0 \leq t \leq (m_\mu^2 - s) \quad \text{and} \quad 0 \leq s \leq m_\mu^2. \quad (19)$$

Further, Γ_χ is the total width of the χ -boson. We focus in what follows in a scenario where $1 \text{ MeV} \lesssim m_\chi \lesssim m_\mu$. In this mass range, the dominant decay channels are $\chi \rightarrow e^- e^+, \bar{\nu}_{L1} \nu_{L1}, \bar{\nu}_{L2} \nu_{L2}$. Using the electron interaction vertex from Eq. (16) and the neutrino interaction vertex from Eq. (4), we find that the total decay width is:

$$\Gamma_\chi = \frac{m_\chi}{24\pi} \left(|g_{ee}^{LL}|^2 + |g_{ee}^{RR}|^2 + |g_\chi q_{L1}|^2 + |g_\chi q_{L2}|^2 \right), \quad (20)$$

which satisfies $\Gamma_\chi \ll m_\chi$.

We show in Fig. 1 the ratio between $\Gamma(\mu \rightarrow e\chi)$ and $\Gamma(\mu \rightarrow 3e)$ as a function of m_χ , for a representative case where χ couples only to the right-handed leptons (i.e. $q_{L1} = q_{L2} = 0$; and $q_{e2} = 0$, $g_\chi q_{e1} = 2$, and $\tan \theta_R = 1$) or when χ couples only to the left-handed leptons (i.e. $q_{e1} = q_{e2} = 0$; and $q_{L2} = 0$, $g_\chi q_{L1} = 2$, and $\tan \theta_L = 1$). In the former case, the ratio is $\simeq 1$, and in the latter it is $\simeq 5$; with a mild sensitivity to the concrete choices of the charges and mixing angles. This result can be understood using the narrow width approximation (NWA), which holds when χ is produced close to the mass shell. Under this approximation, one can replace in the propagators:

$$\frac{1}{(x - m_\chi^2)^2 + m_\chi^2 \Gamma_\chi^2} \rightarrow \frac{\pi}{m_\chi \Gamma_\chi} \delta(x - m_\chi^2), \quad (21)$$

where x is any Mandelstam variable. Under this approximation, the decay rate for $\mu \rightarrow 3e$ reads:

² An analogous behavior occurs in the top decay $t \rightarrow bW^+$. The decay rate is $\Gamma(t \rightarrow bW^+) \sim m_t^2/m_W^2$ and naively diverges when $m_W \rightarrow 0$. However, since both masses arise as a consequence of the spontaneous breaking of the electroweak symmetry, $\Gamma(t \rightarrow bW^+) \sim m_t y_t^2/g^2$ and is finite.




Fig. 1. Ratio of rates of $\mu \rightarrow e\chi$ and $\mu \rightarrow 3e$ as a function of m_χ for the tree-level model presented in Section 3, for the cases described in the text $q_{L_i} = 0$, $q_{e_2} = 0$, $g_\chi q_{e_1} = 2$, and $\tan \theta_R = 1$ (magenta line); and $q_{e_1} = 0$, $q_{L_2} = 0$, $g_\chi q_{L_1} = 2$, and $\tan \theta_L = 1$ (blue line).

Table 2

Spins and charges under $SU(2)_L \times U(1)_Y \times U(1)_\chi$ of the particles of the model described in Section 4, leading to the decay $\mu \rightarrow e\chi$ at the one loop level. All fields are assumed to be singlets under $SU(3)_C$.

	L_1	L_2	e_{R_1}	e_{R_2}	ϕ	ψ	η
spin	1/2	1/2	1/2	1/2	0	1/2	0
$SU(2)_L$	2	2	1	1	2	1	1
$U(1)_Y$	-1/2	-1/2	-1	-1	+1/2	Y_ψ	Y_η
$U(1)_\chi$	q_L	q_L	q_e	q_e	q_ϕ	q_ψ	q_η

$$\Gamma(\mu \rightarrow 3e) \simeq \frac{m_\mu}{32\pi} \frac{(|g_{e\mu}^{LL}|^2 + |g_{e\mu}^{RR}|^2)(|g_{ee}^{LL}|^2 + |g_{ee}^{RR}|^2)}{|g_{ee}^{LL}|^2 + |g_{ee}^{RR}|^2 + |g_\chi q_{L_1}|^2 + |g_\chi q_{L_2}|^2} \left(2 + \frac{m_\mu^2}{m_\chi^2}\right) \left(1 - \frac{m_\chi^2}{m_\mu^2}\right)^2 + \frac{m_\chi}{64\pi} (|g_{e\mu}^{LL}|^2 |g_{e\mu}^{LL}|^2 + |g_{e\mu}^{RR}|^2 |g_{e\mu}^{RR}|^2) \frac{m_\chi}{m_\mu} \left(1 - 2 \frac{m_\chi^2}{m_\mu^2}\right). \quad (22)$$

As for the decay $\mu \rightarrow e\chi$ the rate apparently diverges as $m_\chi \rightarrow 0$, but is in fact finite since m_μ and m_χ are both generated after the breaking of the $U(1)_\chi$ symmetry. Further, and using Eq. (13), one reproduces the result $\Gamma(\mu \rightarrow e\chi)/\Gamma(\mu \rightarrow 3e) \simeq 1$ or $\simeq 5$ that we obtained numerically for our two representative scenarios. As m_χ becomes larger, the ratio becomes sensitive to the underlying model parameters, although this sensitivity is suppressed by a factor m_χ^2/m_μ^2 , and is hence typically weak, in agreement with the numerical results of Fig. 1.

Given the current experimental limits on $\text{Br}(\mu \rightarrow e\chi)$ and $\text{Br}(\mu \rightarrow 3e)$, the most stringent constraints on the model will stem from the latter process.³ For the case $m_\chi \ll m_\mu$, and using the upper limit $\text{Br}(\mu \rightarrow 3e) \leq 1.0 \times 10^{-12}$ from SINDRUM, one finds very stringent constraints on the strength of the effective couplings. Concretely, when $q_{L_i} = 0$, one finds $|g_{e\mu}^{RR}|/m_\chi \lesssim 1.6 \times 10^{-16}/\text{MeV}$. This limit on the effective parameters can in turn be translated into limits on the Yukawa couplings, $U(1)_\chi$ -charges and gauge coupling, and expectation values of the scalar doublets ϕ_{ij} , with the restriction of reproducing the correct muon mass $m_\mu \simeq 105 \text{ MeV}$. Let us finish this section noting that the rare muon decay $\mu \rightarrow e\gamma$ could occur in this model at the one loop level. However, the strong constraints from the tree-level decay $\mu \rightarrow 3e$ preclude the observation of this process.

4. $\mu \rightarrow e\chi$ at the one loop level

In this section we present a renormalizable model with generation independent $U(1)_\chi$ charges, and containing new fields that generate the process $\mu \rightarrow e\chi$ at the one loop level. To provide masses to charged leptons we introduce a doublet scalar, with hypercharge +1/2, and with $U(1)_\chi$ -charge $q_\phi = q_L - q_e$, such that the Yukawa coupling $y_{jk} \bar{L}_j e_{Rk} \phi + \text{h.c.}$ is allowed. This choice leads to the conservation of the electron and the muon flavors, akin to the Standard Model. To violate the lepton flavor numbers, we introduce a new Dirac fermion ψ and a new complex scalar η , both singlets under $SU(2)_L$, with hypercharges and Y_ψ and Y_η , and $U(1)_\chi$ -charges q_ψ and q_η respectively. The spins and charges of the particles of the model are listed in Table 2.



We assume that $q_e = q_\psi + q_\eta$ and $Y_e = Y_\psi + Y_\eta$, so that the Yukawa couplings $y_i' \bar{e}_{Ri} \psi \eta$ are allowed by the symmetries. We also assume that ϕ acquires a vacuum expectation value, but η does not. The gauge boson of the $U(1)_\chi$ symmetry then acquires a mass

$$m_\chi = g_\chi q_\phi \langle \phi \rangle, \quad (23)$$

which we assume to be $m_\chi \ll m_\mu$.⁴ Further, a mass matrix for the charged leptons is generated, of the form Eq. (7). Let us note that if η acquires an expectation value, a mixing between e_{Ri} and ψ is generated, and the mass matrix becomes instead 3×3 . The analysis in that case would be analogous, although we disregard that possibility for simplicity and assume that $\langle \eta \rangle = 0$.

³ We note that the on-shell χ decays into e^+e^- with a decay length $L_\chi \simeq 7.9 \times 10^{-10} \text{ m} (m_\chi/\text{MeV})^{-2} g^{-2}$, where g is a combination of couplings, cf. Eq. (20), and therefore the decay occurs inside the detector.

⁴ In this simple model, m_μ , m_e and m_χ are all proportional to $\langle \phi \rangle$. However, one can completely uncorrelate the fermion masses and the gauge boson masses by imposing $q_\phi = 0$ and by postulating the existence of another scalar field, whose expectation value contributes to m_χ , but not to the fermion masses.

Fig. 2. One loop diagrams contributing to the decay $\mu \rightarrow e\chi$.Fig. 3. Moduli of the functions $\mathcal{F}_{i\alpha}(x)$, with $i = 1, 2$ and $\alpha = \eta, \psi$, as defined in Eq. (27).

Recasting the Lagrangian in terms of the mass eigenstates of the theory, $e_{L,R}$ and $\mu_{L,R}$, one finds interaction terms with the massive gauge boson χ of the form

$$\begin{aligned} \mathcal{L} \supset & -ig_\chi q_L (\bar{e}_L \gamma^\nu e_L + \bar{\mu}_L \gamma^\nu \mu_L + \bar{\nu}_{L1} \gamma^\nu \nu_{L1} + \bar{\nu}_{L2} \gamma^\nu \nu_{L2}) \chi_\nu - ig_\chi q_e (\bar{e}_R \gamma^\nu e_R + \bar{\mu}_R \gamma^\nu \mu_R) \chi_\nu \\ & - ig_\chi q_\psi \bar{\Psi} \gamma^\nu \Psi \chi_\nu - iq_\eta g_\chi [\eta^* (\partial^\nu \eta) - (\partial^\nu \eta^*) \eta] \chi_\nu, \end{aligned} \quad (24)$$

as well as a Yukawa coupling to the right-handed leptons:

$$\mathcal{L} \supset h_e \bar{e}_R \eta \psi + h_\mu \bar{\mu}_R \eta \psi + \text{h.c.} \quad (25)$$

The process $\mu \rightarrow e\chi$ is generated in this model at the one loop-level, through the four diagrams shown in Fig. 2. The form factors are finite and read:

$$\begin{aligned} F_1(m_\chi^2) = G_1(m_\chi^2) &= \frac{g_\chi h_e h_\mu}{384\pi^2} \frac{m_\chi^2}{M_\eta^2} \left[q_\eta \mathcal{F}_{1\eta} \left(\frac{M_\psi^2}{M_\eta^2} \right) + q_\psi \mathcal{F}_{1\psi} \left(\frac{M_\psi^2}{M_\eta^2} \right) \right], \\ F_2(m_\chi^2) = -G_2(m_\chi^2) &= \frac{g_\chi h_e h_\mu}{384\pi^2} \frac{m_\mu^2}{M_\eta^2} \left[q_\eta \mathcal{F}_{2\eta} \left(\frac{M_\psi^2}{M_\eta^2} \right) + q_\psi \mathcal{F}_{2\psi} \left(\frac{M_\psi^2}{M_\eta^2} \right) \right], \end{aligned} \quad (26)$$

where

$$\begin{aligned} \mathcal{F}_{1\eta}(x) &= \frac{-2 + 9x - 18x^2 + x^3(11 - 6\ln x)}{3(1-x)^4}, \\ \mathcal{F}_{1\psi}(x) &= \frac{16 - 45x + 36x^2 - 7x^3 + 6(2-3x)\ln x}{3(1-x)^4}, \\ \mathcal{F}_{2\eta}(x) &= \frac{1 - 6x + 3x^2(1 - 2\ln x) + 2x^3}{(1-x)^4}, \\ \mathcal{F}_{2\psi}(x) &= \frac{-2 - 3x(1 + 2\ln x) + 6x^2 - x^3}{(1-x)^4}. \end{aligned} \quad (27)$$

The absolute values of these functions are shown in Fig. 3, and are regular at $x = 1$.

Using Eq. (2), and that $F_1 = G_1$, $F_2 = -G_2$, the decay rate can be recast as

$$\Gamma(\mu \rightarrow e\chi) \simeq \frac{m_\mu}{8\pi} \left(1 - \frac{m_\chi^2}{m_\mu^2} \right)^2 \left[\left| F_1(m_\chi^2) \frac{m_\mu}{m_\chi} - F_2(m_\chi^2) \frac{m_\chi}{m_\mu} \right|^2 + 2 \left| F_1(m_\chi^2) - F_2(m_\chi^2) \right|^2 \right], \quad (28)$$




Fig. 4. Branching ratio of the process $\mu \rightarrow e\chi$ as a function of m_χ for the one loop model presented in Section 4, assuming $q_\eta = 1$ and $q_\psi = 0$ (left plot), and $q_\eta = 0$ and $q_\psi = 1$ (right plot); in both cases it was assumed $h_e h_\mu = 1$ and $g_\chi = 1$. The solid lines show the full result obtained from Eq. (28), while the dashed lines neglect the contribution from F_1 . The gray dotted line indicates the current upper limit on $Br(\mu \rightarrow e\chi)$ from the TWIST collaboration.

where we have neglected the electron mass. The form factor F_1 (and G_1) is proportional to m_χ^2/M_η^2 , while F_2 (and G_2) is proportional to m_μ^2/M_η^2 . Inserting these form factors in the rate Eq. (28), one finds that the factors $1/m_\chi$ from the emission of the longitudinal polarization cancel with the factors m_χ^2 implicit in the form factors F_1 and G_1 , yielding a finite rate for $\mu \rightarrow e\chi$ in the limit $m_\chi \rightarrow 0$. Further, one finds that $F_1 m_\mu/m_\chi \propto m_\chi m_\mu/M_\eta^2$, and $F_2 m_\chi/m_\mu \propto m_\chi m_\mu/M_\eta^2$. Since we are assuming $M_\eta, M_\psi \gg m_\mu$, it follows that the rate in the limit $m_\chi \rightarrow 0$ will depend mostly on the form factors F_2 and G_2 , and can be well approximated by:

$$\Gamma(\mu \rightarrow e\chi) \Big|_{m_\chi \rightarrow 0} \simeq \frac{g_\chi^2 |h_e|^2 |h_\mu|^2 m_\mu^5}{(768)^2 \pi^5 M_\eta^4} \left[q_\eta \mathcal{F}_{2\eta} \left(\frac{M_\psi^2}{M_\eta^2} \right) + q_\psi \mathcal{F}_{2\psi} \left(\frac{M_\psi^2}{M_\eta^2} \right) \right]^2. \quad (29)$$

Nevertheless, the form factors F_1 and G_1 generate a sizable contribution to the rate when $m_\chi/m_\mu \gtrsim 0.1$.

We show in Fig. 4, the branching ratio for $\mu \rightarrow e\chi$ for two representative choices of charges, $q_\eta = 1$ and $q_\psi = 0$ (left plot) and $q_\eta = 0$ and $q_\psi = 1$ (right panel), and three choices of the masses of the particles in the loop: $M_\psi = 750$ GeV and $M_\eta = 500$ GeV (blue line), $M_\psi = M_\eta = 500$ GeV (purple line), and $M_\psi = 500$ GeV and $M_\eta = 750$ GeV (red line). These values are compatible with the current searches for exotic charged particles [49,50]. We have also taken for concreteness $h_e h_\mu = 1$ and $g_\chi = 1$, although the scaling of the rates with the Yukawa couplings is straightforward. The solid lines show the full result calculated using Eq. (28), while the dashed lines assume $F_1 = G_1 = 0$. As apparent for the plot, while for $m_\chi \ll m_\mu$ the form factors F_1 and G_1 can be neglected, they modify the rate when $m_\chi/m_\mu \gtrsim 0.1$, especially close to the threshold. We also show the current 90% C.L. upper limit $Br(\mu \rightarrow e\chi) < 8.1 \times 10^{-6}$ from the TWIST collaboration [19].

The process $\mu^+ \rightarrow e^+ e^- e^+$ is generated in this toy model also at the one loop-level, through χ -penguin and through box diagrams; the former are proportional to $h_e^2 h_\mu^2 g_\chi^4$ and the latter to $h_e^6 h_\mu^2$. Assuming $h_e \ll g_\chi$, the decay will be dominated by the penguin diagrams, with doubly differential rate given by:

$$\begin{aligned} \frac{d^2 \Gamma(\mu \rightarrow 3e)}{ds dt} \simeq & \frac{g_\chi^2}{32\pi^3 m_\mu^5} \left[\frac{1}{(m_\chi^2 - s)^2 + m_\chi^2 \Gamma_\chi^2} \left(|q_e|^2 (m_\mu^2 - s - t) (m_\mu^2 s |F_1(m_\chi^2) - F_2(m_\chi^2)|^2 + \right. \right. \\ & t (|F_1(m_\chi^2)| m_\mu^2 - |F_2(m_\chi^2)| s) + |q_L|^2 t (|F_1(m_\chi^2) m_\mu^2 - F_2(m_\chi^2) s|^2 - \\ & \left. \left. t (|F_1(m_\chi^2)|^2 m_\mu^2 - |F_2(m_\chi^2)|^2 s) \right) + t \leftrightarrow s + \right. \\ & \left. \frac{2|q_e|^2 (m_\mu^2 - s - t) (m_\chi^2 (\Gamma_\chi^2 + m_\chi^2 - s - t) + s t)}{\left((m_\chi^2 - s)^2 + m_\chi^2 \Gamma_\chi^2 \right) \left((m_\chi^2 - t)^2 + m_\chi^2 \Gamma_\chi^2 \right)} \right. \\ & \left. \left. (m_\mu^2 (s + t) (|F_1(m_\chi^2)|^2 - F_1(m_\chi^2) F_2(m_\chi^2)) + |F_2(m_\chi^2)|^2 s t) \right) \right], \quad (30) \end{aligned}$$

where $s \equiv (p_\mu - p_{e_1})^2$ and $t \equiv (p_\mu - p_{e_2})^2$, with kinematic limits given in eq. (19), and Γ_χ the total decay width of χ . Similarly to Section 3, the dominant decay modes are $\chi \rightarrow e^- e^+, \bar{\nu}_{L_1} \nu_{L_1}, \bar{\nu}_{L_2} \nu_{L_2}$, with width:

$$\Gamma_\chi = \frac{g_\chi^2 m_\chi}{24\pi} (|q_e|^2 + 3|q_L|^2). \quad (31)$$

We show in Fig. 5, the branching ratio for $\mu \rightarrow 3e$ for the same choices of q_η and q_ψ as in Fig. 4, and adopting $q_L = 1 + q_e$ (with $q_e = q_\psi + q_\eta$), using the full result Eq. (30) (solid lines) or setting the form factors $F_1 = G_1 = 0$ (dashed lines). As for $\mu \rightarrow e\chi$, the form factors can be neglected when $m_\chi \ll m_\mu$, and only contribute to the rate when $m_\chi/m_\mu \gtrsim 0.1$. In Fig. 6 we show the ratio of rates




Fig. 5. Same as Fig. 4, but for the process $\mu \rightarrow 3e$, assuming $q_L = 1 + q_e$.




Fig. 6. Same as Fig. 4, but for the ratio of rates $\Gamma(\mu \rightarrow e\chi)/\Gamma(\mu \rightarrow 3e)$, assuming $q_L = 1 + q_e$.

$\Gamma(\mu \rightarrow e\chi)/\Gamma(\mu \rightarrow 3e)$ as a function of m_χ for each of the cases. We find that the ratio is ~ 1 . As in Section 3, this result can be understood analytically employing the narrow width approximation. Under this approximation, the decay rate for $\mu \rightarrow 3e$ reads

$$\Gamma(\mu \rightarrow 3e) \simeq \frac{m_\mu}{4\pi} \frac{|q_e|^2 + |q_L|^2}{|q_e|^2 + 3|q_L|^2} \left(1 - \frac{m_\chi^2}{m_\mu^2}\right)^2 \left[\left| F_1(m_\chi^2) \frac{m_\mu}{m_\chi} - F_2(m_\chi^2) \frac{m_\chi}{m_\mu} \right|^2 + 2|F_1(m_\chi^2) - F_2(m_\chi^2)|^2 \right] + \frac{g_\chi^2 |q_e|^2}{16\pi} \frac{m_\chi^2}{m_\mu} \left(1 - 2 \frac{m_\chi^2}{m_\mu^2}\right) \left(2(|F_1(m_\chi^2)|^2 - F_1(m_\chi^2)F_2(m_\chi^2)) + |F_2(m_\chi^2)|^2 \frac{m_\chi^2}{m_\mu^2} \right). \quad (32)$$

Also in this scenario one finds an apparent divergence when $m_\chi \rightarrow 0$, however the factor $1/m_\chi$ in the rate is canceled by the factor m_χ implicitly contained in the form factor F_1 . As a result, in the limit $m_\chi \rightarrow 0$ the rate for $\mu \rightarrow 3e$ is finite and comparable to the rate for $\mu \rightarrow e\chi$, quite independently of the masses and charges of the particles in the loop.

Given that the current limits on the processes $\mu \rightarrow 3e$ and $\mu \rightarrow e\chi$, we expect the former to yield the strongest limits on this scenario. This is apparent from Fig. 5: the three choices of parameters are allowed by the current constraints on $\mu \rightarrow e\chi$, but several orders of magnitude above the SINDRUM limit $\text{Br}(\mu \rightarrow 3e) < 1.0 \times 10^{-12}$.

5. Conclusions

We have studied in detail the lepton flavor violating process $\mu \rightarrow e\chi$, with χ a massive gauge boson arising from the spontaneous breaking of a local $U(1)$ symmetry. We have constructed the most general effective interaction between a muon, an electron and a massive gauge boson, and we have calculated the decay rate in terms of the corresponding form factors. The decay rate presents terms inversely proportional to the inverse of the χ -boson mass, corresponding to the decay into the longitudinal component of the χ -boson, which naively lead to an enhancement of the rate when χ is very light.

We have constructed two gauge invariant and renormalizable models where the decay $\mu \rightarrow e\chi$ is generated either at tree level or at one loop. We have analyzed the behavior of the rate in the limit $m_\chi \ll m_\mu$, and we have explicitly checked that the rate remains finite. We have also calculated the expected rate for the process $\mu \rightarrow 3e$, mediated by an off-shell χ . For these two models, the ratio of rates of $\mu \rightarrow e\chi$ and $\mu \rightarrow 3e$ is $\mathcal{O}(1)$ in the range of χ -masses considered. Correspondingly, and in view of the current limits on $\mu \rightarrow 3e$ from the SINDRUM collaboration, it would be necessary an improvement of experiments searching for $\mu \rightarrow e\chi$ [51] of at least 5-6 orders of magnitude compared to the TWIST sensitivity in order to observe a signal.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Pablo Roig reports financial support was provided by Secretaría de Educación Pública. Pablo Roig reports financial support was provided by Conacyt. Pablo Roig reports financial support was provided by Fundación Marcos Moshinsky. Alejandro Ibarra reports financial support

was provided by German Research Foundation. Alejandro Ibarra reports financial support was provided by Germany Excellence Strategy - EXC-2094 - 390783311. Marcela Marín reports financial support was provided by Conacyt.

Acknowledgements

This work has been supported by the Collaborative Research Center SFB1258, by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-2094 - 390783311, by the projects CB-250628 (CONACYT), Fondo SEP-Cinvestav number 142 and Cátedra Marcos Moshinsky (Fundación Marcos Moshinsky), as well as by the scholarships for Marcela Marín's Ms. Sc. and Ph. D. Theses (CONACYT). P. R. is indebted to Denis Epifanov for suggesting this fascinating research topic. We have benefited from enriching discussions with our experimental colleagues Eduard de la Cruz Burelo and Iván Heredia de la Cruz. We especially thank Xabi Marciano and Julian Heeck for insightful remarks on the manuscript.

References

- [1] S.L. Glashow, Partial symmetries of weak interactions, *Nucl. Phys.* 22 (1961) 579–588.
- [2] S. Weinberg, A model of leptons, *Phys. Rev. Lett.* 19 (1967) 1264–1266.
- [3] A. Salam, Weak and electromagnetic interactions, *Conf. Proc. C 680519* (1968) 367–377.
- [4] Super-Kamiokande Collaboration, Y. Fukuda, et al., Evidence for oscillation of atmospheric neutrinos, *Phys. Rev. Lett.* 81 (1998) 1562–1567, arXiv:hep-ex/9807003.
- [5] SNO Collaboration, Q.R. Ahmad, et al., Measurement of the rate of $\nu_e + d \rightarrow p + p + e^-$ interactions produced by ^8B solar neutrinos at the Sudbury Neutrino Observatory, *Phys. Rev. Lett.* 87 (2001) 071301, arXiv:nucl-ex/0106015.
- [6] SNO Collaboration, Q.R. Ahmad, et al., Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory, *Phys. Rev. Lett.* 89 (2002) 011301, arXiv:nucl-ex/0204008.
- [7] E.P. Hincks, B. Pontecorvo, Search for gamma-radiation in the 2.2-microsecond meson decay process, *Phys. Rev.* 73 (1948) 257–258.
- [8] R.D. Sard, E.J. Althaus, A search for delayed photons from stopped sea level cosmic-ray mesons, *Phys. Rev.* 74 (Nov 1948) 1364–1371.
- [9] MEG Collaboration, A.M. Baldini, et al., Search for the lepton flavour violating decay $\mu^+ \rightarrow e^+ \gamma$ with the full dataset of the MEG experiment, *Eur. Phys. J. C* 76 (8) (2016) 434, arXiv:1605.05081.
- [10] SINDRUM Collaboration, U. Bellgardt, et al., Search for the decay $\mu^+ \rightarrow e^+ e^+ e^-$, *Nucl. Phys. B* 299 (1988) 1–6.
- [11] SINDRUM II Collaboration, W. Honecker, et al., Improved limit on the branching ratio of $\mu \rightarrow e$ conversion on lead, *Phys. Rev. Lett.* 76 (1996) 200–203.
- [12] SINDRUM II Collaboration, J. Kaulard, et al., Improved limit on the branching ratio of $\mu^- \rightarrow e^+$ conversion on titanium, *Phys. Lett. B* 422 (1998) 334–338.
- [13] SINDRUM II Collaboration, W.H. Bertl, et al., A search for muon to electron conversion in muonic gold, *Eur. Phys. J. C* 47 (2006) 337–346.
- [14] L. Willmann, et al., New bounds from searching for muonium to anti-muonium conversion, *Phys. Rev. Lett.* 82 (1999) 49–52, arXiv:hep-ex/9807011.
- [15] BNL Collaboration, D. Ambrose, et al., New limit on muon and electron lepton number violation from $K_L^0 \rightarrow \mu^\pm e^\mp$ decay, *Phys. Rev. Lett.* 81 (1998) 5734–5737, arXiv:hep-ex/9811038.
- [16] LHCb Collaboration, R. Aaij, et al., Search for the lepton-flavor violating decays $B_s^0 \rightarrow e^\pm \mu^\mp$ and $B^0 \rightarrow e^\pm \mu^\mp$, *Phys. Rev. Lett.* 111 (2013) 141801, arXiv:1307.4889.
- [17] L. Calibbi, G. Signorelli, Charged lepton flavour violation: an experimental and theoretical introduction, *Riv. Nuovo Cimento* 41 (2) (2018) 71–174, arXiv:1709.00294.
- [18] J. Jaeckel, A. Ringwald, The low-energy frontier of particle physics, *Annu. Rev. Nucl. Part. Sci.* 60 (2010) 405–437, arXiv:1002.0329.
- [19] TWIST Collaboration, R. Bayes, et al., Search for two body muon decay signals, *Phys. Rev. D* 91 (5) (2015) 052020, arXiv:1409.0638.
- [20] F. Wilczek, Axions and family symmetry breaking, *Phys. Rev. Lett.* 49 (1982) 1549–1552.
- [21] B. Grinstein, J. Preskill, M.B. Wise, Neutrino masses and family symmetry, *Phys. Lett. B* 159 (1985) 57–61.
- [22] Z.G. Berezhiani, M.Y. Khlopov, Cosmology of spontaneously broken gauge family symmetry, *Z. Phys. C* 49 (1991) 73–78.
- [23] J.L. Feng, T. Moroi, H. Murayama, E. Schnapka, Third generation familons, b factories, and neutrino cosmology, *Phys. Rev. D* 57 (1998) 5875–5892, arXiv:hep-ph/9709411.
- [24] M. Hirsch, A. Vicente, J. Meyer, W. Porod, Majoron emission in muon and tau decays revisited, *Phys. Rev. D* 79 (2009) 055023, arXiv:0902.0525, Erratum: *Phys. Rev. D* 79 (2009) 079901.
- [25] J. Jaeckel, A family of WISPy dark matter candidates, *Phys. Lett. B* 732 (2014) 1–7, arXiv:1311.0880.
- [26] A. Celis, J. Fuentes-Martin, H. Serodio, An invisible axion model with controlled FCNCs at tree level, *Phys. Lett. B* 741 (2015) 117–123, arXiv:1410.6217.
- [27] A. Celis, J. Fuentes-Martin, H. Serodio, A class of invisible axion models with FCNCs at tree level, *J. High Energy Phys.* 12 (2014) 167, arXiv:1410.6218.
- [28] I. Galon, A. Kwa, P. Tanedo, Lepton-flavor violating mediators, *J. High Energy Phys.* 03 (2017) 064, arXiv:1610.08060.
- [29] L. Calibbi, F. Goertz, D. Redigolo, R. Ziegler, J. Zupan, Minimal axion model from flavor, *Phys. Rev. D* 95 (9) (2017) 095009, arXiv:1612.08040.
- [30] Y. Ema, K. Hamaguchi, T. Moroi, K. Nakayama, Flaxion: a minimal extension to solve puzzles in the standard model, *J. High Energy Phys.* 01 (2017) 096, arXiv:1612.05492.
- [31] F. Björkeröth, E.J. Chun, S.F. King, Flavourful axion phenomenology, *J. High Energy Phys.* 08 (2018) 117, arXiv:1806.00660.
- [32] M. Bauer, M. Neubert, S. Renner, M. Schnubel, A. Thamm, Axionlike particles, lepton-flavor violation, and a new explanation of a_μ and a_e , *Phys. Rev. Lett.* 124 (21) (2020) 211803, arXiv:1908.00008.
- [33] J. Heeck, H.H. Patel, Majoron at two loops, *Phys. Rev. D* 100 (9) (2019) 095015, arXiv:1909.02029.
- [34] C. Cornella, P. Paradisi, O. Sumensari, Hunting for ALPs with lepton flavor violation, *J. High Energy Phys.* 01 (2020) 158, arXiv:1911.06279.
- [35] L. Calibbi, D. Redigolo, R. Ziegler, J. Zupan, Looking forward to lepton-flavor-violating ALPs, *J. High Energy Phys.* 2021 (Sep 2021).
- [36] L.B. Okun, Y.B. Zeldovich, Paradoxes of unstable electron, *Phys. Lett. B* 78 (1978) 597–600.
- [37] L.B. Okun, M.B. Voloshin, On the electric charge conservation, *JETP Lett.* 28 (1978) 145.
- [38] M. Suzuki, Slightly massive photon, *Phys. Rev. D* 38 (1988) 1544.
- [39] Y. Farzan, I.M. Shoemaker, Lepton flavor violating non-standard interactions via light mediators, *J. High Energy Phys.* 07 (2016) 033, arXiv:1512.09147.
- [40] J. Heeck, Lepton flavor violation with light vector bosons, *Phys. Lett. B* 758 (2016) 101–105, arXiv:1602.03810.
- [41] Y. Farzan, J. Heeck, Neutrinophilic nonstandard interactions, *Phys. Rev. D* 94 (5) (2016) 053010, arXiv:1607.07616.
- [42] S.T. Petcov, The processes $\mu \rightarrow e + \gamma$, $\mu \rightarrow e + \bar{\nu}$, $\nu' \rightarrow \nu + \gamma$ in the Weinberg-Salam model with neutrino mixing, *Sov. J. Nucl. Phys.* 25 (1977) 340, Erratum: *Sov. J. Nucl. Phys.* 25 (1977) 698, Erratum: *Yad. Fiz.* 25 (1977) 1336.
- [43] W.J. Marciano, A.I. Sanda, Exotic decays of the muon and heavy leptons in gauge theories, *Phys. Lett. B* 67 (1977) 303–305.
- [44] B.W. Lee, R.E. Shrock, Natural suppression of symmetry violation in gauge theories: muon - lepton and electron lepton number nonconservation, *Phys. Rev. D* 16 (1977) 1444.
- [45] J.M. Cornwall, D.N. Levin, G. Tiktopoulos, Derivation of gauge invariance from high-energy unitarity bounds on the s matrix, *Phys. Rev. D* 10 (1974) 1145, Erratum: *Phys. Rev. D* 11 (1975) 972.
- [46] C.E. Vayonakis, Born helicity amplitudes and cross-sections in nonabelian gauge theories, *Lett. Nuovo Cimento* 17 (1976) 383.
- [47] B.W. Lee, C. Quigg, H.B. Thacker, Weak interactions at very high-energies: the role of the Higgs boson mass, *Phys. Rev. D* 16 (1977) 1519.
- [48] M.S. Chanowitz, M.K. Gaillard, Multiple production of W and Z as a signal of new strong interactions, *Phys. Lett. B* 142 (1984) 85–90.
- [49] CMS Collaboration, A.M. Sirunyan, et al., Search for supersymmetric partners of electrons and muons in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* 790 (2019) 140–166, arXiv:1806.05264.
- [50] ATLAS Collaboration, G. Aad, et al., Search for electroweak production of charginos and sleptons decaying into final states with two leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector, *Eur. Phys. J. C* 80 (2) (2020) 123, arXiv:1908.08215.
- [51] M. Aiba, et al., Science case for the new high-intensity muon beams HIMB at PSI, arXiv:2111.05788.